



## Original Article

## Complete inhibition of phosphatase and tensin homolog promotes the normal and oxygen-glucose deprivation/reperfusion-injured PC12 cells to cell death

Sohrab Minaei Beyrami<sup>1</sup>, Mohammad Hasan Khadem Ansari<sup>1</sup>, Yousef Rasmi<sup>1</sup>, Nader Shakib<sup>1</sup>, Poursan Karimi<sup>2\*</sup>

<sup>1</sup>Department of Biochemistry, Faculty of Medicine, Urmia University of Medical Sciences, Urmia, Iran

<sup>2</sup>Neurosciences Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

## Article info

## Article History:

Received: 18 February 2018  
Accepted: 26 April 2018  
published: 21 May 2018

## Keywords:

OGD  
Reperfusion Injury  
AKT  
p38  
MAPK  
PC12 Cells

## Abstract

**Introduction:** Lipid phosphatase and tensin homolog deleted from chromosome 10 (PTEN) antagonizes phosphoinositide 3-kinase (PI3K)/AKT cell survival pathway. The effect of PTEN inhibitors has been rarely examined on cell survival following reperfusion injury. In this study, we investigated the neuroprotective effect of SF1670, as a new PTEN inhibitor, on an in vitro stroke-like model.

**Methods:** PC12 cells were exposed to oxygen-glucose deprivation/reperfusion (OGD/R). The cells were treated in five conditions as follows: normoxic normoglycemic (NO/NG); 60 minutes OGD; 60 minutes OGD and 6 h reperfusion (OGD/R); OGD/R treated with 10  $\mu$ M SF1670 (OGD/R-SF), and NO/NG treated with 10  $\mu$ M SF1670 (NO/NG-SF). Then, phosphorylation levels of AKT, P38 in PC12 cells were measured by immunoblotting. The cell viability was also determined by colorimetric assay.

**Results:** The results of immunoblotting revealed that following OGD/R the levels of phospho-AKT (p-AKT) significantly decreased, compared to NO/NG cells ( $P < 0.05$ ). However, the ratio of p-AKT/total AKT significantly increased in the presence of SF1670 in the OGD/R-SF group, compared to the OGD/R condition. On the other hand, SF1670 significantly reduced the p-P38 MAPK and p-JNK levels, compared to OGD/R cells. Moreover, cell viability significantly decreased in the OGD and OGD/R condition compared to NO/NG cells. Surprisingly, SF-treated cells (OGD/R-SF and NO/NG-SF group) showed low cell viability compared to NO/NG condition.

**Conclusion:** Overall, our results demonstrated that complete inhibition of phosphatase activity of PTEN not only did not exhibit neuroprotective effect but also promoted PC12-deprived cells to death.

**Please cite this article as:** Minaei Beyrami S, Khadem Ansari MH, Rasmi Y, Shakib N, Karimi P. Complete inhibition of phosphatase and tensin homolog promotes the normal and oxygen-glucose deprivation/reperfusion-injured PC12 cells to cell death. *J Cardiovasc Thorac Res* 2018;10(2):83-89. doi: 10.15171/jcvtr.2018.13.

## Introduction

Ischemic stroke, as a common life-threatening cerebrovascular disease,<sup>1</sup> causes a high percentage of permanent disabilities all around the world.<sup>2</sup> The fibrinolytic treatment of ischemic stroke or successful recanalization<sup>3</sup> restores the oxygenation but initiates secondary local inflammation after reperfusion<sup>3,4</sup> which in turn exacerbates cerebral tissue injury, the so-called reperfusion injury (I/R).<sup>4,5</sup> I/R injury initiate widespread inflammation, reactive oxidation, excitotoxicity and cell-specific dysregulation of metabolic processes promoting neurodegeneration through specific programmed cell death mechanisms.<sup>6</sup>

studies have proven that the balance between two intracellular signaling pathways including the phosphatidylinositol 3-kinase/AKT (PI3K/AKT) as a cell survival pathway, and mitogen-activated protein kinase (MAPK) as an inflammatory pathway<sup>7</sup> play important role in the determination of cell fate after I/R.<sup>8</sup> Different mechanisms which are involved in the inhibition or enhancement of these pathways could be taken as suitable therapeutic targets to limit I/R damages.<sup>9</sup> MAPKs family is consist of p38 group of protein kinases, c-jun N-terminal (c-JNK), and extracellular signal-regulated kinases (ERKs). P38 MAPK pathway, as a major stress kinase, activates through phosphorylation in responses to cellular

\*Corresponding Author: A Poursan Karimi, Email: pouran.karimi@yahoo.com

© 2018 The Author (s). This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

stress, heat shock, oxidative stress, and inflammation.<sup>10,11</sup> A research has revealed that activation of P38 initiates apoptosis cascade and up-regulates pro-inflammatory cytokines production.<sup>12</sup> In addition, C-JNK activation is also associated with neuronal death.<sup>13-15</sup>

In contrary, phosphorylated AKT (p-AKT) in downstream phosphorylates numerous substrates are involved in cell growth,<sup>16</sup> proliferation,<sup>17</sup> and survival,<sup>18</sup> as well as cellular metabolism, glucose uptake and angiogenesis and protects cells from I/R induced cell injury.<sup>19</sup>

Phosphatase and tensin homolog (PTEN) is a lipid phosphatase and tumor suppressor which plays an important role in the regulation of cell proliferation, differentiation, and apoptosis.<sup>20,21</sup> Up-regulation of PTEN is capable to inhibit AKT activation by degradation of PI3P to PIP2<sup>16,21</sup> and facilitates the p38 MAPK signaling pathway.<sup>22,23</sup> Decreasing PTEN activity through phosphorylation of PTEN<sup>24</sup> or deleting of one copy of its gene increases resistance to apoptotic cell death.<sup>25</sup> Hence, down-regulation of PTEN may lead to a neuroprotective effect in the neurodegenerative disorders with a beneficiary role in I/R injury.<sup>26-28</sup>

The current paper used a recently developed specific PTEN inhibitor, SF1670,<sup>29</sup> for inhibiting PTEN pathway.<sup>30,31</sup> Li et al showed that pretreatment with SF1670 in nanomolar concentration enhances PIP3 signaling in transplanted neutrophils.<sup>29</sup> In the current study, for the first time, we examined the effects of pretreatment with high dose (10  $\mu$ M) of SF1670 for complete inhibition of phosphatase activity of PTEN against I/R injury in oxygen-glucose deprivation (OGD) as an in vitro stroke-like model in PC12 cultured cells.

## Materials and Methods

### Chemicals and antibodies

PC12 cells were obtained from Pasteur Institute (Tehran, Iran), DMEM (Gibco, Grand Island, NY, USA), p-Akt1/2/3 Antibody (Ser 473): sc-7985 (SANTA CRUZ, CA, USA), p-p38 Antibody (Tyr 182): sc-101759 (SANTA CRUZ, CA, USA), p38 $\alpha$  Antibody (N-20): sc-728 (SANTA CRUZ, CA, USA), p-JNK Antibody (14.Thr 183/Tyr 185): sc-293136 (SANTA CRUZ, CA, USA), JNK1/3 Antibody (C-17): sc-474, RIPA Buffer (Sigma-Aldrich, New York, NY, USA), Anti-Protease Cocktail (Sigma-Aldrich, New York, NY, USA), Acrylamide and bisacrylamide (Sigma-Aldrich, New York, NY, USA).

### Cell culture

Rat pheochromocytoma-derived cell line PC12 cells were cultured in normoxic normoglycemic (NO/NG) condition as following: The cells were seeded in six-well plate in Dulbecco's modified Eagle's medium (DMEM) (Gibco, Grand Island, NY, USA) supplemented with 10% horse serum, 5% fetal bovine serum (FBS), 100 kU/L of penicillin, and 100 mg/L of streptomycin (Sigma, St. Louis, MS, USA) and maintained at 37°C in a normoglycemic

(5 mM glucose) and humidified normoxic atmosphere incubator (95% air and 5% CO<sub>2</sub>). All treatments were performed on cells at 80% confluence.

### Oxygen-glucose deprivation/reperfusion model

In order to mimic ischemic-like conditions in vitro, PC12 cells were exposed to OGD/R injury as following: the culture media of the cell, DMEM, was replaced by Hanks Balanced Salt (HBSS; glucose concentration = 0 mg/dl) and then transferred to a hypoxic chamber (95% nitrogen and CO<sub>2</sub> 5%) for 60 minutes. At the end of the OGD phase, the cells reperused for 6 h in normoxic-normoglycemic (NO/NG) condition.<sup>32,33</sup> The pretreatment was performed by 10  $\mu$ M of SF1670 24 hours before OGD induction (OGD/R-SF group). We also pre-treated a batch of normoxic and normoglycemic cells by SF1670 as drug control (NO/NG-SF group).

### Western blotting

PC12 cells were subjected to Western blot analysis for phosphorylation of p38, and AKT proteins as previously described<sup>34</sup> with minor changes. Briefly, 10<sup>6</sup> cells were homogenized in 500  $\mu$ l lysis buffer [0.05 mmol/L Tris-NaOH (pH=8), 150 mmol/L NaCl, 0.01 mmol/L EGTA, 1%SDS, 0.1% anti Protease Cocktails (ROCHE)]. The supernatants were mixed with loading buffer solution containing 60 mM Tris-HCl, 25% glycerol, 2% SDS, 14.4 mmol/L 2-mercaptoethanol, and 0.1% bromophenol blue. Then proteins were separated on a 10% SDS-polyacrylamide gel and transferred onto the nitrocellulose membrane. After incubation in blocking buffer (phosphate buffered saline, 3% (w/v) BSA, 0.1% Tween 20), the membranes were probed overnight at 4°C with the appropriate primary antibody as follows: rabbit polyclonal anti-phospho-Akt, anti-AKT, anti-phospho-p38, anti-p38, anti-phospho-JNK, anti-JNK antibodies. Having washed and exposed to horseradish peroxidase-conjugated secondary antibody for 1 hour at room temperature, antibody-antigen complexes were visualized by enhanced chemiluminescence substrates. The scanned images of the protein bands were analyzed using ImageJ (National Institutes of Health, Bethesda, Maryland, USA) software.

### Cell viability assay

Cell viability was determined using the 3-[4, 5-dimethylthiazol-2-yl]-2, 5-diphenyltetrazolium bromide (MTT). PC12 cells (2-5  $\times$  10<sup>5</sup> cells/well) were seeded in 96-well plates in DMEM medium until confluence 80%. Following the interventions, MTT reagent (20  $\mu$ l, 5 mg/ml in PBS) was added to wells and incubated for 4 h at 37°C. Then the medium was removed, and replaced with 150  $\mu$ l DMSO. Afterward, absorbance was measured at 570 nm by an automatic microplate reader (Awareness Technologies Stat Fax 4200).

### Statistical analysis

Data was analyzed using SPSS software version 16.0 (SPSS, Chicago, IL, USA) and expressed in mean  $\pm$ SD. One-way ANOVA and Tukey's post-hoc tests were used in statistical comparisons. *P* values less than 0.05 were considered as significant.


### Results

#### The effect of SF1670 on AKT phosphorylation in OGD/R-injured PC12 cells


Figure 1 shows that AKT phosphorylation levels (p-AKT) significantly ( $P < 0.05$ ) decreased in the OGD-treated cells, compared to NO/NG cells. Moreover, reperfusion (6 hours) following OGD significantly ( $P < 0.001$ ) decreased p-AKT/total AKT ratio in the OGD/R cells, compared to NO/NG cells. Conversely, SF1670 pretreatment (24 hours before OGD/R) significantly ( $P < 0.01$ ) increased p-AKT/total AKT ratio, compared to the OGD/R condition. Surprisingly, SF1670 ( $P < 0.01$ ) decreased the phosphorylation of AKT in normoxic normoglycemic cells (NO/NG-SF), compared to the NO/NG cells.

#### The effect of SF1670 on p38 MAPK phosphorylation in OGD/R-injured PC12 cells

Figure 2 shows a significant increase in the average levels of phospho-P38 MAPK (p-P38 MAPK) in the OGD ( $P < 0.01$ ) and OGD/R ( $P < 0.001$ ) conditions, compared to the NO/NG condition. Nevertheless, SF1670 administration in OGD/R-SF group significantly ( $P < 0.01$ ) decreased the P38 MAPK phosphorylation, compared to OGD/R cells. No significant change was observed in the levels of p-P38 MAPK in the NO/NG-SF condition, compared to the NO/NG condition.



**Figure 1.** The effect of SF1670 on AKT phosphorylation in PC12 cell exposed to OGD/R. **A)** Immunoblotting images of p-AKT and total AKT proteins in PC12 cells exposed to one hour OGD followed by 6 h of reperfusion and/or 10 $\mu$ M SF1670 pretreatment. **B)** Quantitative densitometric analysis of the p-AKT against total AKT protein bands. Data are presented as means  $\pm$  SEM. \**P* < 0.05, \*\**P* < 0.01, \*\*\**P* < 0.001 vs. NO/NG. ##*P* < 0.01 vs. OGD/R. [NO/NG: normoxic/normoglycemic cells; OGD: Oxygen and glucose deprived cells (60 min); R: Reperused cells for 6 hours; SF: SF1670 (10  $\mu$ M) treated cells with SF1670 one hour before OGD].



**Figure 2.** The effect of SF1670 on phosphorylation of P38 MAPK in PC12 cell exposed to OGD/R. **A)** Immunoblotting images of p-P38 MAPK and total P38 MAPK proteins in PC12 cells exposed to one hour OGD followed by 6 h of reperfusion and/or 10 $\mu$ M SF1670 pretreatment. **B)** Quantitative densitometric analysis of the p-P38 MAPK against total P38 MAPK protein bands. Data are presented as means  $\pm$  SEM. \*\**P* < 0.01, \*\*\**P* < 0.001 vs. NO/NG. ##*P* < 0.01 vs. OGD/R. [NO/NG: normoxic/normoglycemic cells; OGD: Oxygen and glucose deprived cells (60 min); R: Reperused cells for 6 hours; SF: SF1670 (10  $\mu$ M) treated cells with SF1670 one hour before OGD].

#### The effect of SF1670 on c-JNK phosphorylation after being exposed to OGD/R

Our results also demonstrated that OGD/R condition significantly (Figure 3,  $P < 0.05$ ) increased phospho-JNK (p-JNK) levels, compared to the NO/NG. However, SF1670 pretreatment in the OGD/R-SF group decreased the p-JNK when compared to OGD/R cells ( $P < 0.05$ ). No significant change was observed in the levels of p-JNK between NO/NG-SF and NO/NG condition.

#### The effect of SF1670 on cell viability


In addition, MTT assay was carried out to evaluate cellular viability in the PC12 cells. The results showed that cell viability was reduced after exposure to OGD for 1 hour ( $P < 0.05$ ), and OGD-R for 6 hours ( $P < 0.01$ ), compared to NO/NG condition (Figure 4). The SF1670 at the concentration of 10  $\mu$ mol/L decreased ( $P < 0.05$ ) cell viability in OGD/R-SF, compared to the OGD/R. Furthermore, SF1670 decreased the cell viability in NO/NG-SF, compared to NO/NG ( $P < 0.001$ ).

### Discussion

The current paper deprived PC12 cells of oxygen and glucose for 1h followed by 6 hours reperfusion to induce a valid in vitro stroke-like model. The current study aimed at investigating the neuroprotective effect of PTEN inhibitor, SF1670, on OGD/R-induced injury by assessing the phosphorylation of AKT, JNK, and P38 MAPK proteins, and cell viability. The results of the current study showed that SF1670 increased p-AKT, and decreased p-P38, p-JNK, and cell viability in the PC12 cells exposed to OGD/R insult.

The oxygen and glucose deprivation are two common






**Figure 3.** The effect of SF1670 on phosphorylation of JNK in PC12 cell exposed to OGD/R. **A)** Immunoblotting images of p-JNK and total JNK (as loading control) proteins in PC12 cells exposed to one hour OGD followed by 6 h of reperfusion and/or 10 $\mu$ M SF1670 pretreatment. **B)** Quantitative densitometric analysis of the p-JNK against total JNK protein bands. Data are presented as means  $\pm$  SEM. \* $P$  < 0.05, vs. NO/NG; # $P$  < 0.05 vs. OGD/R. [NO/NG: normoxic/normoglycemic cells; OGD: Oxygen and glucose deprived cells (60 min); R: Reperused cells for 6 hours; SF: SF1670 (10  $\mu$ M) treated cells with SF1670 one hour before OGD].

reasons for ischemic insult damages which are deteriorated after reoxygenation.<sup>1,8,10</sup> PTEN is an upstream negative regulator of AKT signaling pathway which inhibits AKT phosphorylation by PI3K, the positive regulator of AKT. Therefore, PTEN activity is associated with down-regulation of PI3K/Akt down-streams involved in cell survival pathways.<sup>35,36</sup> Conversion of Inositol diphosphate (IP<sub>2</sub>) to Inositol 3, 4, 5-triphosphate (IP<sub>3</sub>) is required to Akt phosphorylation which promotes cellular survival and attenuates cell death. The current paper showed that exposure to OGD and reperfusion reduced p-AKT levels in the PC12 cells. As shown in Figures 1 and 4, the parallel changes were detected for p-AKT expression and cell viability during the experiment. Similarly, in the previous report the OGD/R-induced cell viability loss in PC12 cells was associated with diminished Akt phosphorylation.<sup>37</sup> In support of our finding, previous studies also showed that ischemia and reperfusion inhibited PI3K/Akt pathway resulting in cell death.<sup>38,39</sup>

P38 and JNK, stress-activated kinases, play important role in the regulation of apoptosis signals. Previous studies demonstrated that JNK and P38 activities were up-regulated in response to the brain ischemia. In addition, these proteins are involved in cerebral ischemia/reperfusion insult, and inhibition of their phosphorylation attenuates ischemic brain injury.<sup>40-42</sup> Recently, a study also reported that OGD/R increases p-P38 and p-JNK in PC12 cells.<sup>42</sup> Similarly, in the present experiment, the levels of p-JNK and p-P38 expression were significantly amplified following OGD/R. Nevertheless, SF1670 pretreatment prevented these changes induced by OGD/R. Moreover, our study showed that increased p-P38 MAPK and p-JNK levels were accompanied with cell viability loss in OGD/R. In addition, it has been shown that AKT signaling via phosphorylation of apoptosis signal-regulated kinase 1



**Figure 4** The effect of SF1670 on cell viability (MTT) in cultured PC12 cells exposed to OGD/R insult or OGD-R-SF1670. Data are presented as means  $\pm$  SEM. \* $P$  < 0.05, \*\* $P$  < 0.01, \*\*\* $P$  < 0.001. vs. NO/NG; # $P$  < 0.05 vs. OGD/R. [NO/NG: normoxic/normoglycemic cells; OGD: Oxygen and glucose deprived cells (60 min); R: Reperused cells for 6 hours; SF: SF1670 (10  $\mu$ M) treated cells with SF1670 one hour before OGD].

(ASK1), inhibits the JNK and p38-mediated apoptosis.<sup>43,44</sup> In this study, SF1670 treatment attenuated p-AKT, p-p38, and p-JNK accompanied by cell viability loss. Although previous evidence indicated that activation of p38 and JNK is associated with apoptotic cell death, some reports revealed that p38 MAPK plays a critical role in the control of cell survival and proliferation.<sup>45-47</sup> Phong et al have demonstrated that p38 signaling promotes cell survival in response to DNA damage possibly to inhibit the onset of premature apoptosis.<sup>48</sup> Moreover, previous reports linked p38 signaling pathway to increased levels of antiapoptotic protein such as Bcl-2 and Bcl-x<sub>1</sub> following DNA damage and stress.<sup>49,50</sup> It seems that the role of p38 signaling in the control of apoptosis is context dependent and depending on the physiological context of the stress induction it may switch from cell survival to pro-apoptosis. In the present study, SF1670 attenuated p-p38 and p-JNK in PC12 cells accompanied by low cell viability. We suggest that ameliorated JNK and P38 MAPK signaling pathways promote cells toward cell death.

Furthermore, we found that pretreatment of NO/NG cells with SF1670 attenuated p-AKT expression and cell viability as assessed by MTT. We suggest that SF1670 in 10  $\mu$ M concentration has a neurotoxic effect on PC12 cells. In addition, it is likely that other apoptosis-prompting factors override the cytoprotective effects of SF1670 activity. This paradoxical effect of 10  $\mu$ M of SF1670 may be related to the dual roles of PTEN in the cell. Lately, Zhou et al. reported that inhibition of PTEN with bpV(Hopic) aggravates ischemic acute kidney injury via augmenting apoptosis and inflammation.<sup>51</sup> It has also reported that astrocytic PTEN loss exacerbated ischemia damage.<sup>52</sup> Evidence has also shown that PTEN-knockout mice died in early development.<sup>16</sup> Therefore, it is likely that the complete suppression of phosphatase activity of PTEN could not be a good idea in cell protection, particularly in this model which was made by neuroblastoma cells. Probably PTEN has other crucial roles in cell viability that have not been clarified yet which needs further studies. Similar to

other novel protein, the specific substrates of PTEN are a mystery. To identify of this phosphatase, systematic approaches including generation of null mutations, exploration of possible roles in transient overexpression studies, further studies are needed to be given mutants with normal PTEN activities.

In summary, the study demonstrated that complete inhibition of phosphatase activity of PTEN promoted cells toward death, possibly through attenuation p38 signaling pathways in OGD/R PC12 cells.

#### Ethical approval

The ethical approval for this study was obtained from ethics committee of Tabriz University of Medical Sciences.

#### Competing interests

None.

#### References

1. Adams HP Jr, Bendixen BH, Kappelle LJ, Biller J, Love BB, Gordon DL, et al. Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. TOAST. Trial of Org 10172 in Acute Stroke Treatment. **Stroke** 1993;24:35-41. doi: 10.1161/01.STR.24.1.35
2. Adams HP Jr, Brott TG, Furlan AJ, Gomez CR, Grotta J, Helgason CM, et al. Guidelines for Thrombolytic Therapy for Acute Stroke: a Supplement to the Guidelines for the Management of Patients with Acute Ischemic Stroke. A statement for healthcare professionals from a Special Writing Group of the Stroke Council, American Heart Association. **Stroke** 1996;27:1711-8. doi: 10.1161/01.CIR.94.5.1167
3. Tissue plasminogen activator for acute ischemic stroke. The National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group. **N Engl J Med** 1995;333:1581-7. doi:10.1056/NEJM199512143332401
4. Asaithambi G, Tong X, George MG, Tsai AW, Peacock JM, Luepker RV, et al. Acute stroke reperfusion therapy trends in the expanded treatment window era. **J Stroke Cerebrovasc Dis** 2014;23:2316-21. doi: 10.1016/j.jstrokecerebrovasdis.2014.04.023.
5. Chan PH. Reactive oxygen radicals in signaling and damage in the ischemic brain. **J Cereb Blood Flow Metab** 2001;21:2-14. doi:10.1097/00004647-200101000-00002
6. Chen X, Kintner DB, Baba A, Matsuda T, Shull GE, Sun D. Protein aggregation in neurons following OGD: a role for Na<sup>+</sup> and Ca<sup>2+</sup> ionic dysregulation. **J Neurochem** 2010;112:173-82. doi: 10.1111/j.1471-4159.2009.06438.x.
7. Achiwa Y, Hasegawa K, Udagawa Y. Regulation of the phosphatidylinositol 3-kinase-Akt and the mitogen-activated protein kinase pathways by ursolic acid in human endometrial cancer cells. **Biosci Biotechnol Biochem** 2007;71:31-7. doi: 10.1271/bbb.60288
8. Alessandrini A, Namura S, Moskowitz MA, Bonventre JV. MEK1 protein kinase inhibition protects against damage resulting from focal cerebral ischemia. **Proc Natl Acad Sci U S A** 1999;96:12866-9. doi: 10.1073/pnas.96.22.12866
9. Bai L, Ren F, Wang W, Zheng S, Zhang J, Chen Y, et al. [Protective role of PTEN inhibition against liver ischemia-reperfusion injury in mice and its underlying mechanisms]. **Zhonghua Gan Zang Bing Za Zhi** 2014;22:451-5. doi: 10.3760/cma.j.is.sn.1007-3418.2014.06.011
10. Choi EK, Yeo JS, Park CY, Na HI, Lim JA, Lee JE, et al. Inhibition of reactive oxygen species downregulates the MAPK pathway in rat spinal cord after limb ischemia reperfusion injury. **Int J Surg** 2015;22:74-8. doi: 10.1016/j.ijisu.2015.08.016.
11. Coulthard LR, White DE, Jones DL, McDermott MF, Burchill SA. p38(MAPK): stress responses from molecular mechanisms to therapeutics. **Trends Mol Med** 2009;15:369-79. doi: 10.1016/j.molmed.2009.06.005.
12. Ozbek E, Cekmen M, Ilbey YO, Simsek A, Polat EC, Somay A. Atorvastatin prevents gentamicin-induced renal damage in rats through the inhibition of p38-MAPK and NF-kB pathways. **Ren Fail** 2009;31:382-92. doi: 10.1080/08860220902835863
13. Shin W-H, Park S-J, Kim E-J. Protective effect of anthocyanins in middle cerebral artery occlusion and reperfusion model of cerebral ischemia in rats. **Life Sci J** 2006;79:130-7. doi: 10.1016/j.lfs.2005.12.033
14. Sawe N, Steinberg G, Zhao H. Dual roles of the MAPK/ERK1/2 cell signaling pathway after stroke. **J Neurosci Res** 2008;86:1659-69. doi: 10.1002/jnr.21604
15. Jiang M, Li J, Peng Q, Liu Y, Liu W, Luo C, et al. Neuroprotective effects of bilobalide on cerebral ischemia and reperfusion injury are associated with inhibition of pro-inflammatory mediator production and down-regulation of JNK1/2 and p38 MAPK activation. **J Neuroinflammation** 2014;11:167. doi: 10.1186/s12974-014-0167-6
16. Carracedo A, Pandolfi PP. The PTEN-PI3K pathway: of feedbacks and cross-talks. **Oncogene** 2008;27:5527-41. doi:10.1038/onc.2008.247
17. Chappell WH, Steelman LS, Long JM, Kempf RC, Abrams SL, Franklin RA, et al. Ras/Raf/MEK/ERK and PI3K/PTEN/Akt/mTOR inhibitors: rationale and importance to inhibiting these pathways in human health. **Oncotarget** 2011;2:135-64. doi: 10.18632/oncotarget.240
18. Carracedo A, Ma L, Teruya-Feldstein J, Rojo F, Salmena L, Alimonti A, et al. Inhibition of mTORC1 leads to MAPK pathway activation through a PI3K-dependent feedback loop in human cancer. **J Clin Invest** 2008;118:3065-74. doi: 10.1172/JCI34739.
19. Ding J, Wang J, Li QY, Yu JZ, Ma CG, Wang X, et al. Neuroprotection and CD131/GDNF/AKT Pathway of Carbamylated Erythropoietin in Hypoxic Neurons. **Mol Neurobiol**. 2017;54 (7):5051-5060. doi: 10.1007/s12035-016-0022-0
20. Dahia PL. PTEN, a unique tumor suppressor gene. **Endocr Relat Cancer** 2000;7:115-29. doi: 10.1677/erc.0.0070115
21. Chu EC, Tarnawski AS. PTEN regulatory functions in tumor suppression and cell biology. **Med Sci Monit**

- 2004;10:Ra235-41.
22. Duman BB, Sahin B, Acikalin A, Ergin M, Zorludemir S. PTEN, Akt, MAPK, p53 and p95 expression to predict trastuzumab resistance in HER2 positive breast cancer. **J BUON** 2013;18:44-50.
  23. Ghorbani A, Zand H, Jeddi-Tehrani M, Koohdani F, Shidfar F, Keshavarz SA. PTEN over-expression by resveratrol in acute lymphoblastic leukemia cells along with suppression of AKT/PKB and ERK1/2 in genotoxic stress. **J Nat Med** 2015. doi: 10.1007/s11418-015-0915-7
  24. Zhu Y, Hoell P, Ahlemeyer B, Krieglstein J. PTEN: a crucial mediator of mitochondria-dependent apoptosis. **Apoptosis** 2006;11:197-207. doi: 10.1007/s10495-006-3714-5
  25. Li L, Liu F, Salmonsens RA, Turner TK, Litofsky NS, Di Cristofano A, et al. PTEN in neural precursor cells: regulation of migration, apoptosis, and proliferation. **Mol Cell Neurosci** 2002;20:21-9. doi: 10.1006/mcne.2002.1115
  26. Gary DS, Mattson MP. PTEN regulates Akt kinase activity in hippocampal neurons and increases their sensitivity to glutamate and apoptosis. **Neuromolecular Med** 2002;2:261-9. doi:10.1385/NMM:2:3:261
  27. Ning K, Pei L, Liao M, Liu B, Zhang Y, Jiang W, et al. Dual neuroprotective signaling mediated by downregulating two distinct phosphatase activities of PTEN. **J Neurosci** 2004;24:4052-60. doi:10.1523/JNEUROSCI.5449-03.2004
  28. Lee JH, Kim KY, Lee Y-K, Park SY, Kim CD, Lee WS, et al. Cilostazol prevents focal cerebral ischemic injury by enhancing casein kinase 2 phosphorylation and suppression of phosphatase and tensin homolog deleted from chromosome 10 phosphorylation in rats. **J Pharm Exp Ther** 2004;308:896-903. doi: 10.1124/jpet.103.061853
  29. Li Y, Prasad A, Jia Y, Roy SG, Loison F, Mondal S, et al. Pretreatment with phosphatase and tensin homolog deleted on chromosome 10 (PTEN) inhibitor SF1670 augments the efficacy of granulocyte transfusion in a clinically relevant mouse model. **Blood** 2011;117:6702-13. doi: 10.1182/blood-2010-09-309864
  30. Liao J, Marumoto T, Yamaguchi S, Okano S, Takeda N, Sakamoto C, et al. Inhibition of PTEN tumor suppressor promotes the generation of induced pluripotent stem cells. **Mol Ther** 2013;21:1242-50. doi: 10.1038/mt.2013.60
  31. Liu C, Wu J, Xu K, Cai F, Gu J, Ma L, et al. Neuroprotection by baicalein in ischemic brain injury involves PTEN/AKT pathway. **J Neurochem** 2010;112:1500-12. doi: 10.1111/j.1471-4159.2009.06561.x
  32. Guan J, Du S, Lv T, Qu S, Fu Q, Yuan Y. Oxygen-glucose Deprivation Preconditioning Protects Neurons against Oxygen-glucose Deprivation/reperfusion Induced Injury via Bone Morphogenetic Protein-7 Mediated ERK, p38 and Smad Signaling Pathways. **Clin Exp Pharmacol Physiol**. 2016;43(1):125-34. doi: 10.1111/1440-1681.12492
  33. Ma Y, Zhao P, Zhu J, Yan C, Li L, Zhang H, et al. Naoxintong Protects Primary Neurons from Oxygen-Glucose Deprivation/Reoxygenation Induced Injury through PI3K-Akt Signaling Pathway. **Evid Based Complement Alternat Med** 2016;2016:5815946. doi: 10.1155/2016/5815946
  34. Khani M, Motamedi P, Dehkhoda MR, Dabagh Nikukheslat S, Karimi P. Effect of thyme extract supplementation on lipid peroxidation, antioxidant capacity, PGC-1alpha content and endurance exercise performance in rats. **J Int Soc Sports Nutr** 2017;14:11. doi: 10.1186/s12970-017-0167-x
  35. Kim H-A, Kim K-J, Seo KH, Lee H-K, Im S-Y. PTEN/MAPK pathways play a key role in platelet-activating factor-induced experimental pulmonary tumor metastasis. **FEBS Lett** 2012;586:4296-302. doi: 10.1016/j.febslet.2012.10.034
  36. Walker C, Liu N-K, Xu X-M. PTEN/PI3K and MAPK signaling in protection and pathology following CNS injuries. **Front Biol** 2013;8:421-33. doi: 10.1007/s11515-013-1255-1
  37. Kinkade CW, Castillo-Martin M, Puzio-Kuter A, Yan J, Foster TH, Gao H, et al. Targeting AKT/mTOR and ERK MAPK signaling inhibits hormone-refractory prostate cancer in a preclinical mouse model. **J Clin Invest** 2008;118:3051-64. doi: 10.1172/JCI34764
  38. Xu X, Chua CC, Gao J, Chua K-W, Wang H, Hamdy RC, et al. Neuroprotective effect of humanin on cerebral ischemia/reperfusion injury is mediated by a PI3K/Akt pathway. **Brain Res** 2008;1227:12-8. doi: 10.1016/j.brainres.2008.06.018
  39. Zhang L, Qu Y, Tang J, Chen D, Fu X, Mao M, et al. PI3K/Akt signaling pathway is required for neuroprotection of thalidomide on hypoxic-ischemic cortical neurons in vitro. **Brain Res**. 2010 Oct 21;1357:157-65. doi: 10.1016/j.brainres.2010.08.007
  40. Gao Y, Signore AP, Yin W, Cao G, Yin X-M, Sun F, et al. Neuroprotection against focal ischemic brain injury by inhibition of c-Jun N-terminal kinase and attenuation of the mitochondrial apoptosis-signaling pathway. **J Cereb Blood Flow Metab** 2005;25:694-712. doi: 10.1038/sj.jcbfm.9600062
  41. Qi L-L, Fang S-H, Shi W-Z, Huang X-Q, Zhang X-Y, Lu Y-B, et al. CysLT 2 receptor-mediated AQP4 up-regulation is involved in ischemic-like injury through activation of ERK and p38 MAPK in rat astrocytes. **Life Sci** 2011;88:50-6. doi: 10.1016/j.lfs.2010.10.025
  42. Xu J, Hua C, Pan X, Fu X, Wu W. Eupatilin inhibits OGD/R-induced neuronal injury in PC12 cells. **Int J Clin Exp Med** 2017;10:6728-34.
  43. Kim AH, Khursigara G, Sun X, Franke TF, Chao MV. Akt phosphorylates and negatively regulates apoptosis signal-regulating kinase 1. **Mol Cell Biol** 2001;21:893-901. doi: 10.1128/MCB.21.3.893-901.2001
  44. Yang Z-Z, Tschopp O, Baudry A, Duemmler B, Hynx D, Hemmings BA. Physiological functions of protein kinase B/Akt. Portland Press Limited; 2004. doi: 10.1042/bst0320350
  45. Van Laethem A, Van Kelst S, Lippens S, Declercq W, Vandenabeele P, Janssens S, et al. Activation of p38 MAPK is required for Bax translocation to mitochondria, cytochrome c release and apoptosis induced by UVB irradiation in human keratinocytes.

- FASEB J** 2004;18:1946-8. doi: 10.1096/fj.04-2285fje
46. Deacon K, Mistry P, Chernoff J, Blank JL, Patel R. p38 Mitogen-activated protein kinase mediates cell death and p21-activated kinase mediates cell survival during chemotherapeutic drug-induced mitotic arrest. **Mol Biol Cell** 2003;14:2071-87. doi: 10.1091/mbc.e02-10-0653
  47. Sen P, Chakraborty PK, Raha S. RETRACTED: Activation of p38MAPK by repetitive low-grade oxidative stress leads to pro-survival effects. **Biochim Biophys Acta** 2007;1773(3):367-74. doi: 10.1016/j.bbamcr.2006.11.012
  48. Phong MS, Van Horn RD, Li S, Tucker-Kellogg G, Surana U, Ye XS. p38 Mitogen-Activated Protein Kinase Promotes Cell Survival in Response to DNA Damage but Is Not Required for the G(2) DNA Damage Checkpoint in Human Cancer Cells. **Mol Cell Biol** 2010;30:3816-26. doi: 10.1128/MCB.00949-09
  49. Kim M-J, Choi S-Y, Park I-C, Hwang S-G, Kim C, Choi Y-H, et al. Opposing roles of c-Jun NH2-terminal kinase and p38 mitogen-activated protein kinase in the cellular response to ionizing radiation in human cervical cancer cells. **Mol Cancer Res** 2008;6:1718-31. doi: 10.1158/1541-7786.MCR-08-0032
  50. Flacke J-P, Kumar S, Kostin S, Reusch HP, Ladilov Y. Acidic preconditioning protects endothelial cells against apoptosis through p38-and Akt-dependent Bcl-xL overexpression. **Apoptosis** 2009;14:90-6. doi: 10.1007/s10495-008-0287-5
  51. Zhou J, Jia L, Hu Z, Wang Y. Pharmacological Inhibition of PTEN Aggravates Acute Kidney Injury. **Sci Rep** 2017;7:9503. doi: 10.1038/s41598-017-10336-8
  52. Li W, Huang R, Chen Z, Yan L-J, Simpkins JW, Yang S-H. PTEN degradation after ischemic stroke: a double-edged sword. **Neuroscience** 2014;274:153-61. doi: 10.1016/j.neuroscience.2014.05.027