Geography and Environmental Planning Journal 23rd Year, vol. 47, No.3, Autumn 2012 ISSN (Online): 2252-0848 ISSN (Print): 2008-5354 http://uijs.ui.ac.ir/gep

Site selection of the suitable areas for the physical development of Tehran megalopolis based on the climatic elements and geographic factors

F. Mojarrad. S. Hoseinifar Received: August 13, 2011 / Accepted: December 13, 2011, 5-8 P

Extended abstract 1- Introduction

Developing the cities and it's influences on the spatial structure of megalopolises has always been one of the most important factors for the planners. Among the most important factors directing the development of the cities, are natural factors such as climatic ones which have been less considered in the country as yet because of various factors like economic benefits resulting from construction immethodical of the buildings. Since this carelessness has caused an unwise and easy-going development of Tehran megalopolis in unsuitable geographic directions, so this research intends to accomplish the optimum site selection for physical

Author(s)

F. Mojarrad (🖂)

Assistant Professor of Climatology, Razi University, Kermanshah, Iran e-mail: f_moiarrad@vahoo.com

S. Hoseinifar

MA. Of Climatology, Razi University, Kermanshah, Iran

development of Tehran megalopolis based on climatic elements and geographic factors.

2- Methodology

Topographic maps of the region which mainly consist some parts of Tehran and Alborz provinces on a scale of 1:250,000 were prepared and the region boundary was defined on them. In the next step, the climatic data of six meteorological stations was taken from the Iranian Meteorological Organization and, after reconstruction, was considered through the similar time range of 1984 to 2005. Then, using ArcGIS software and on the basis of the climatic and geographic factors, various layers affecting the site selection including topography, digital elevation model (DEM), slope percent and aspect, solar radiation angle, temperature means (minimum, maximum and daily), mean diurnal temperature difference, 24 hrs. maximum precipitation, and mean wind

GEP Journal

speed were made and entered into the software. Pixels' values in the climatic were layers calculated via spline interpolation method regression or equations. Then the resulted raster layers, after the reclassification of their pixels' values, were weighed and overlayed using Spatial Analyst (SA) and Spatial Analytical Hierarchy Process (SAHP) models and thus the final maps of the physical development suitability of these two models were obtained.

On the other hand, two Landsat ETM-7 images prepared from Iranian Space Agency and some geometric corrections were made on them using PCI (Geomatica) software in UTM WGS84 projection. Before geometric corrections, because of temporal differences of images, radiometric corrections were made on them too. radiometric correction or normalization means the reconstruction of image values so that there is a linear and similar relation between pixels and their real radiations in the whole imaging The result was the Normalized area. Difference Vegetation Index (NDVI) map and the land use map. As the last step, the final site selection maps were presented via overlaying the NDVI and the land use maps with final maps of each two SA and SAHP models.

3– Discussion

Based upon the distribution of the development suitability zones in the site selection maps, the most unfavorable areas have been developed in the high slope sections of the north of the region in the SA model. According to this model, about 8500 km² of the region has not any special limitation for the development. In

the SAHP model too, the most important limitation in the way of the development is the high slope sections of north of the region. The most suitable areas with almost 1400 km² area stands in the south, west and the submontanes of north and northwest of the region. Partly suitable areas with the area equal to 5300 km² have been developed in the central and southern parts of the region. The comparison of the final maps of two models reveals that the most suitable areas in the SAHP model have lesser extent than in the SA model.

4- Conclusion

Considering the bare land expansion which is mainly developed in the south of region and overlaying it with suitability zones of the SA and SAHP models and in order to protect the vegetation cover specially in the western parts of the region, among the three suitable zones for physical development of the city, the south direction is the most favorable direction and the western and southwestern regions respectively stand in the next precedences. In order to get a more comprehensive study in this regard, the other natural factors such as geologic, geomorphologic, hydrologic and human factors must be taken into account too.

Key words: site selection, physical development, Tehran, climate, GIS

References

- Alijani, B., (2006), Climate of Iran, Payam-e Noor Univ. Pub., Tehran, pp. 221.
- Badr, R., (2000), the use of GIS and RS in determining the city's physical development direction (Case Study:

Razi city), M. Sc. Thesis, supervisor: Alimohammadi, Abbas, Tarbiat Modarres Univ., Tehran, Dept. of remote sensing.

- Bahraini, S. H., (2011), Process of urban designing, Tehran Univ. Pub., 6th Edition, Tehran, pp. 469.
- Bahram Soltani, K., (1992), Issues and methods of urban planning (environment), urban planning & architecture research center, Ministry of housing and urban development, 1th Edition, Tehran, pp. 240.
- Bahram Soltani, K., (2001), The role of climate in urban environments studies, Green Wave, 5: 18-21.
- Bathrellos, G. D., Gaki-Papanastassiou, K., Skilodimou, H. D., Papanastassiou, D., Chousianitis, K. G., (2011), Potential suitability for urban planning and industry development using natural hazard maps and geologicalparameters. geomorphological Environ Earth Sci, published online 04 August 2011, www. springerlink. com.
- Burrough, P. A., (1986), Principles of Geographic Information System for Land Resources Assessment, Clarendon Press, Oxford. 193 pp.
- Chakhar, S., Mousseau, V., (2008), GISbased multicriteria spatial modeling generic framework, Int J Geogr Inf Sci 22(11–12): 1159–1196.
- Dong, J., Zhuang, D., Xu, X., Ying, L., (2008), Integrated evaluation of urban development suitability based on remote sensing and GIS techniques a case study in Jingjinji Area, China, Sensors, 8:5975–5986.
- Esbah, H., (2007), Land Use Trends During Rapid Urbanization of the City

of Aydin, Turkey, Environ Manage, 39:443-459.

- Faraji Sabokbar, H. A., (2005), Site selection of commercial services units using AHP method (Case Study: Torghabeh District of Mashhad County), Geographical Researches, 51: 125-137.
- Farajzadeh, M., (2010), Principals of geographic information system, Entekhab Pub., 1th Edition, Tehran, pp. 224.
- Ghayoor, H. A., (1996), Flood and floody areas in Iran, Geographical Research, 40: 101-120.
- Khosravi, M., (2004), Determining the physical development direction of Andimeshk city using satellite data (RS) and geographic information system (GIS), M. Sc. Thesis, supervisors: Nazarian, A., Tavallaei S., Tarbiat Moallem Univ., Tehran, Dept. of Geography.
- Liu, Y. G., Zeng, X. X., Xu, L., Tian D.
 L., Zeng, G. M., Hu, X. J., Tang, Y.
 F., (2011), Impacts of land-use change on ecosystem service value in Changsha, China, J. Cent. South Univ. Technol., 18: 420–428.
- Ministry of Power, (2006), Draft of guide of flood damage evaluation, Iranian Water Resources Management Co., Issue No: 296-A, pp. 95.
- Mkhabela, M. S., Bullock, P., Raj, S., Wang, S., Yang, Y., (2011), Crop yield forecasting on the Canadian Prairies using MODIS NDVI data, Agricultural and Forest Meteorology, 151: 385–393.
- Movahedi, S., Soltanian, M. (2011), GIS and climatology, Kankash Press & Pub., 1th Volume, 1th Edition, Esfahan, pp. 188.

GEP Journal

- Over, S., Buyuksarac, A., Bekta, O., Filazi, A., (2011), Assessment of potential hazard and site effect in Antakya (Hatay Province), SE Turkey, Environ Earth Sci, 62: 313– 326.
- Rabiei Dastjerdi, H. R., (2003), Modeling the unreliability of change detection based on the satellite data classification (Case Study: Esfahan city), M. Sc. Thesis, supervisor: Ziaeian, P., Tarbiat Modarres Univ., Tehran, Dept. of remote sensing.
- Rahimion, A., (1998), The analysis of suitability of inside city lands using geographic information system in Baghershahr city in Tehran, M. Sc. Thesis, Tarbiat Modarres Univ., Tehran, Dept. of remote sensing.
- Rahnemaei, M. T., (1999), Spatial limitations of Tehran city, Geographical Researches, 34: 7-19.
- Saaty, T. L., (2004), Decision making–the analytic hierarchy and network processes (AHP/ANP), J Syst Sci Syst Eng, 13(1): 1–35.
- Saeednia, A., (1993), Location of Tehran

city, Environmental researches collection, 15: 43-61.

- Thapa, R. B., Murayama, Y., (2010), Drivers of urban growth in the Kathmandu Valley, Nepal: examining the efficacy of the analytic hierarchy process, Appl Geogr, 30: 70–83.
- Tudes, S., Yigiter, N.D., (2010) Preparation of land use planning model using GIS based on AHP: case study Adana-Turkey, Bull Eng Geol Environ, 69: 235–245.
- Xiao, J., Shen, Y., Ge, J., Tateishi, R., Tang, C., Liang, Y., Huang, Z. (2006), Evaluating urban expansion and land use change in Shijiazhuang, China, by using GIS and remote sensing, Landsc Urban Plan, 75:69–80.
- Yesilnacar, E., Doyuran, V., (2000), Selection of Settlement Areas Using GIS and Statistical Method (Spatial-AHP), Middle East Technical University, Ankara.
- Zooeshtiagh, H., (1992), Abstract of Tehran organizing plan, Ministry of housing and urban development, Tehran.