

Geography and Environmental Planning Journal 24th Year, Vol. 50, No.2, Summer 2013 ISSN (Online): 2252-0848 ISSN (Print): 2008-5354 http://uijs.ui.ac.ir/gep

Neotectonic, morphoclimatic and anthropogenic agents in Appearance and Genesis of alluvial fans (Case study: Garmsar alluvial fan)

S. Shayan, M. R. Sharifikia, Gh. Zare

Received: April 19, 2011/ Accepted: March 10, 2012, 17-20 P



Alluvial fans are common feature in the hillside of mountain, where that sudden change in slope river bed lead to reduce the flow rate and sediments are left according to size. The most appropriate place for the creation of alluvial fans is located output of the mountains and arrival of water network into plain, vast plains and even in the centers of plains in adjacent terminal basins. In this research was tried to investigate the effective factors on creation of Garmsar alluvial fan.

2- Material and method

In this research, first has been collected required data through library research, documents relating to the subject. Then

Author(s)

S. Shayan(🖾)

Assistant Professor of Geomorphology, Tarbiat Modarres University, Tehran, Iran

email: Shayan@Modares.ac.ir

M. Sharifikia

Assistant Professor Remote sensing, Tarbiat Modarres University, Tehran, Iran

Gh. Zare

PhD. Student of Geomorphology, Tarbiat Modarres University, Tehran, Iran

was determined territory of Hablerod basin and Garmsar alluvial fan by 1:50000 topographic maps and satellite image. Information of lithology was extracted of 1:100000 and 1:250000 geological maps. The study of temperature and precipitation has been made based on meteorological data from synoptic stations of Garmsar, Firouzkoh and Pluviometer of Namrood and Simindasht.

The study of sediment and debit rates was performed based on estimation of the sediment and debit stations of Firouzkoh (Hablerood), Namrood. Simindasht (Hablerood), Simindasht (Dalichay), Bonkoh (Hablerood) and Gorsafid. Also the study of neotectonic condition by the 1:50000 topographic maps, based on four indicators of the sinousity geomorphologic Mountain Front (Smf), the valley floor width to height ratio (Vf), the curve Hypsometry (Hc), the asymmetric basin waterways (AF). In order to quicken in the research process was used of Arc GIS software and field survey for Ensure of (human actions).

3- Discussion

The factors affected on creation of Garmsar alluvial fan can be divided into two parts of essentials and basic. The morphology and sediment are essential factors and the climate and hydrology are basic parameters. In Hablerod basin, slope (by the addition of runoff rate, faster delivery of erosion materials to downstream) is one of the important factors that create and transport alluvial materials and sediment, in the creation and extention of Garmsar alluvial fan. Hablerod basin is considered large watersheds, so this area is one of the factors in the creation of Garmsar alluvial fan. Garmsar fault with movement and precession of path Hablerod role plays role in locating on this fan. 22.84% of the Hablerod basin rocks are with low resistance or sensitive, 55.98 percent of rocks have the moderate strength and 21.18 percent of rocks have high strength. Therefore the type of basin rock is also one of the factors in sediment creation. Most basin debit rate occurred in months Mars and April.

4- Conclusion

The various factors affect on the creation and extension of Garmsar alluvial fan. These factors can be divided into two categories: basic and essential factors. The morphology and sediment are essential factors, the climate, hydrology and human are basic factors in creation of Garmsar alluvial fan. Also basin poor lithology causes to weather more rapidly and increases the sedimentation. Basin climate with Temperature and precipitation affects on types of weathering, causing floods and etc. Basin hydrology with transport and deposition of sediment material is effective on creation Garmsar alluvial

fan. On the one hand in recent years, the man efforts have increased sediment and on the other hand, by construct dam on the Hablerod have reduced sedimentation in the Garmsar alluvial fan.

Key Word: River Process, Garmsar Alluvial fan, Neotectonic, Anthropogenic Geomorphology

Refrences

- Abasnejad. A, 1997, Geomorphological studies in Rafsanjan plain, PhD Thesis, Faculty of Humanities and Social Sciences University of Tabriz, 475 p.
- Alaei Taleghani. M, 2003, Geomorphology of Iran, Ghomes Press, 413 p.
- Alizade. A. 2006, Applied Principles of Hydrology, Astan Ghodse Razavi press, 856 p.
- Aram. A, 1988, Science in Islam, Sorosh press, Tehran.
- Beaumont, P, 1972, Alluvial fans along the foothills of the Elburz Mountains, Iran, paleogeography, Paleoclimatology, Paleology, V 12.
- Blair, T.C et al, 1994, Alluvial Fans and Their Natural Distinction from Rivers Based on
- Bull, W. B, 1964, Geomorphology of segmented alluvial fans in western Fresno County, California. U. S, Geological Survey professional paper, 352-E.
- Bull, W.B, 1977, The alluvial fan environment, progress in physical geography, Vol 1, pp 222-270.
- Bull. W.B, McFadden, L.D, 1977,
 Tectonic geomorphology north and south of the Garlock fault, California:
 In: Doehring, D, O. Geomorphology symposium. State university of New York, Binghamton.
- Giles, Philip, 2010, Investigation the use of alluvial fan volume to represent fan size in morphometric studies, Geomorphology 121.

- Harvey, A.M, 1997, The Role of Alluvial fans in arid-zone fluvial systems. Wiley, Chichester.
- Harvey. A. M. 2002, The role of baselevel change in the dissection of alluvial fans" case studies from southeast Spain and Nevada, Journal of Geomorphology, Vol 45, pp 67-87.
- Hassanabadi. D, 2000, Hydrologic and Hydrogeological effects of dispense floods on alluvial fans Hablehrod Garmsar, MA thesis, Department of Earth Sciences, Shahid Beheshti University, 250 p.
- Haug Erik W, 2009, Climatic and geomorphic interactions on alluvial fans in the Atacama Desert, Chile, Master of Science in Geosciences.
- Iranbakhsh. A. R, Hamdi. S. M, Asadi. M, 2008, Flora, life forms and chorotypes of plants of Garmsar region in Semnan province, Journal of Research and development on natural resources, No 79, pp 179-189.
- Jihad Agriculture, 1990, Integrated management of water catchment Hablehrod, Department of Watershed Management, Department of Planning and Budget, 135 p.
- Keller Edward, A, and Nicholas Pinter, 2002, Active Tectonics Earthquake, Uplift, and Morphology, Hydraulic Processes, Sedimentary Processes, and Facies Assemblages: Journal of Sedimentary Research Section a-Sedimentary Petrology and Processes, V. 64.
- Khayam. M, Mokhtari. D, 2004, Performance evaluation of tectonic activity on the morphology of cones Alluvial Fans (case study: the northern slopes Mishodagh Alluvial Fans), Journal of Geographic Research Quarterly, No 44, pp 1-10.
- Maghsodi. M, 2008, Investigation Geomorphological factors in the development of cones Fans (Case Study: Fans Jajroud), Journal of Geographic Research Quarterly, No 65, pp 73-95.
- Mahmodi. F, 2004, Geomorphology of Dynamic, Payamnor press, 309 p.

- Mokhtari. D, 2002, Factors in the development and evolutions of Quaternary alluvial fans of northern range Mishodagh (Azarbaijan Iran) and assessment its potential environmental, PhD Thesis, Faculty of Humanities and Social Sciences University of Tabriz, 510 p.
- Mosaviherami, R, 1990, Sedimentology, Astan Ghodse Razavi press, 410 p.
- Oguchi, T, Ohmori, H, 1994, Analysis of relationships among alluvial fan area, source basin area, basin slope and sediment yield. Zeitschrift fur Geomorphologie 38.
- Pacific Southwest Inter-Agency Committee (PSIAC), 1968, Factors affecting sediment yield in the pacific southwest area and selection and evaluation of measures for reduction of erosion and sediment yield. Report of the Water Management Subcommittee.
- Ramesht, M. H, Seyf. A, Shahzeydi. S.S, Entezari. M, 2009, Tectonic lateral influence on the morphology of alluvial fans Darakhtangan Shahdad in Kerman, Journal of Geographic and
 - Development, No 16, pp 29-46.
- Ramesht. M. H, Shahzeydi. S.S, 2008, Handling faults in the centers of successive and divergent evolution in the Quaternary alluvial fans Darakhtangan, Journal of Geographic and Development Regional, No 10, pp 1-20.
- Rezaei Moghadam. M.H, 1996, Research in the foothills and plains of the southern slopes of cumulative Mishodagh, PhD Thesis, Faculty of Humanities and Social Sciences University of Tabriz, 450 p.
- Rezaei Moghadam. M. H, Maghammoghimi. Gh, Rajabi. M, 2006, Factors affecting the formation and development of alluvial fans on the southern slopes Aladagh River in northeast Iran, Journal of Geographical Research, Vol 79, pp 64-80.
- Roed.M.A et al, 1973, Age of inactive alluvial fan-Bow

River.Alberta.Canadian Journal of Earth Science, 10.

Rostaei. Sh, Rajabi. M, Zomorodian. M. J, Maghammoghimi. Gh, 2009, The role of tectonic activity in the formation and development of the southern slopes of cones Fans Aladagh, Journal of Geographic and Development, No 13, pp 137-156.

Schumm, S.A et al, 1987, Experimental Fluvial Geomorphology, John Wiley and Sons, New York, 413 pp.

Shayan. S, 2000, The density of the outer-Quaternary erosion dynamics and its role in the management of mountain environments, PhD thesis, Tarbiat Modares University, 650 p.

Shayan. S, 2003, Geomorphological features of the basin Gamasiab Alluvial Fans, Journal of Geographic Research Quarterly, No 46, pp. 99-113.

Yamani. M, Maghsodi. M, 2003, Investigation and evaluation channels on the surface of Alluvial Fans (Case Study: Fans Tangoeeh in Sirjan hole), No 45, pp. 103-113.