

Geography and Environmental Planning Journal 24<sup>th</sup> Year, Vol. 50, No.2, Summer 2013 ISSN (Online): 2252-0848 ISSN (Print): 2008-5354 http://uijs.ui.ac.ir/gep

# Synoptic analysis of extensive and persistent frosts in Iran

S.A. Masoodian, M. Darand

Received: January 14, 2012 / Accepted: May 9, 2012, 29-32 P

## Extended Abstract 1- Introduction

In the point of meteorological view, frost occurs when minimum temperature is below  $0^{\circ}$  centigrade degree. In total, frosts classified in two groups: a) Radiative Frosts and b) Advective frosts. Radiative frost is the form most common of the phenomenon. Nocturnal long wave (heat) radiation loss from the surface leads to the progressive reduction of surface temperature (and that of the adjacent air) between sunset and sunrise (Oke, 1987; Linacre and Geerts, 1997). Advective frost occurs under meteorological conditions that typically include the passage across a region of a large mass of very cold, dry air followed by a well-mixed, windy atmosphere and a temperature that is often subzero, even during the daytime(Scholefield, 1994; Davidson, 1997). The combination of factors such those related to atmospheric as

Author(s)

S.A. Masoodian Professor of Climatology, University of Isfahan, Isfahan, Iran M. Darand(⊠) Assistant Professor of Climatology, University of Kordestsn, Sanandaj, Iran email: m.darand@uok.ac.ir circulation, radiative balance, soil humidity, topographic features(Muller et al., 2005) and local factors (for example, latitude and close to the big water resources) are an example of the multiple processes taking part to produce a marked drop in temperature. The frosts have obvious effects on the energy, transportation and agriculture that among the three above-mentioned issues, frost temperature have the most effect on agriculture (Vega et al., 1994). The aim of this study is the synoptic analysis of the extensive and persistent frosts in Iran.

#### 2-Data and Method

In this study the interpolated minimum daily temperature data from Asfazari data base during 11/3/1962 to the 31/12/2004 has been used. The data base by using recorded minimum temperature in 663 synoptic and climatology stations and Kriging geostatistic method has been interpolated for 15\*15 kilometer pixels over Iran. with 15706\*7187 The matrix dimension created that days located on the rows and pixels were on the columns. This matrix was the base of calculation frost days in this study.

For recognition of extensive and persistent frosts, two criteria have been considered. The first one was that frost observed over 50 percent area of Iran. And the other one frost persistence over 3 days. These two criteria discriminate the Advective frost in contrast to Radiative frost type.

#### **3- Discussion**

The results of cluster analysis on the sea level pressure data in the extensive and persistent frost days showed that 5 different synoptic atmospheric circulation patterns result in extensive and persistent frosts in Iran. The persistent extensive and frosts observed during cold seasons so that starts in November and ending in March. The highest frequency occurrence observed in January.

Pattern 1: In this pattern two strong anticyclones observed over Siberia and Europe. The low pressure of sub-polar is very extensive and strong. The displacement of sub-polar low pressure to the southern and eastern part of study region result in displacement of two high pressures of Europe and Siberia to the south and drive to the Iran and occurrence of extensive frost.

Pattern 2: In this pattern the high pressure with the two strong cores between Mongolia and Kazakhstan country. The sub-polar low pressure has been weak rather than pattern 1. Although the Siberian high is very strong in this pattern but the temperature decrease in Iran is weak. The reason is weakness and location sub-polar low pressure in west part.

Pattern 3: In this pattern the geographic location of circulation atmospheric patterns is same as the pattern 1 but the sub polar low pressure is weak. The temperature decrease is accompanied to the occurrence and invasion of the cold air result in two

high pressures over Europe and Siberia.

Pattern 4: The geographic distribution of circulation atmospheric patterns in this pattern is same as the pattern 2 but the sub-polar low pressure is located in western statue. At the point of time view, the highest occurrence of this pattern is in February.

Pattern 5: In this pattern the geographic distribution of atmospheric systems is very different to the other ones. The two high pressures over the Siberia and eastern part of Europe are observed. On the northern part of Russia, a weak low pressure is formed that control the behaviour of high pressures over the Siberia and Eastern Europe.

## **4-Conclusion**

The results of analyzing sea level pressure during the occurrence of extensive and persistent extreme frost over the Iran showed that five atmospheric circulation patterns with geographic different distribution observed. In total in the some patterns the Siberia high presence alone. In the other ones the Siberia and Europe high observed together. If the sub polar low pressure has been strong and displacement to the south, result in displacement of two high pressure of Siberia and Europe to the south and negative temperature anomaly in Iran. Therefore, the sub polar low pressure role is indirect on the occurrence of extensive and persistent extreme frost over the Iran. The extreme extensive frost observed when the Siberia and Europe are presence together. The highest negative temperature decrease is observed over the Zagros, Alborz and north-eastern range mountains. The southern part of Iran and the south of Caspian Sea didn't experience any frost in the study period.

**Keywords:** Frost, Synoptic analysis, Extensive, Persistent, Iran

### References

- Alijani, B & Barati, Gh.R (1996),Synoptic analysis of frost in 1987 spring, Geographical research Quarterly, Vol. 40, 121-135.
- Alijani, B and Tagiloo, M (2010), Synoptic analysis of frost days in Zanjan province of Iran,EMS annual Meeting Abstracts, Vol, 7, EMS2010-167.
- Alijani, B, Mahmmodi, Pyman, Regi Chahi, A.B & Khosravi, P (2009), Analysis of frost days persistence in Iran by using Markov Chain Model, Physical Geography Research Quarterly, Vol.73, 1-20.
- Alijani, B., M. Hoshiar, (2007), Recognition of Synoptic Patterns of Severe Colds of West North of Iran, Physical Geography Research Quarterly, Vol. 65. pp.1-16.
- Barati, G., (1996), Designing and Forecasting the Patterns of Spring Frost Days in Iran, Ph.D. Thesis of physical geography (climatology), Faculty of humanities sciences, Tarbiat Modarres University, p. 213.
- Barati, G., (1999), System Relationships of Migratory High Pressures and Spring Frosts of Iran, geographical Research Quarterly, 14 (3 and 4), pp. 132-150.
- Barati, GH. (1999). Migration systematic relationships and Iranian vernal frosts, quarterly journal of researches for geography, Isfahan, Iran, No. 53, pp 45-57.
- Davidson, D (1997), Frost and ways to minimise the risk, Australian Grapegrower and Winemaker, 403, 21 -22.
- Esazadeh, Sh, Memarian, M.H, Mirrokni, S.M & Taghizadeh,

A(2009), Synoptic analysis of frosts in Eastern Azerbaijan province, 4th Iranian geophysics conference, Tehran 21-23 May, 238-241.

- Fatahi, I., and T. Salehi Pak., (2009), A Synoptic Patterns Analysis of Winter Freezing in Iran, geography and development, 7(13), pp. 127-136.
- Fortune, M.A & Kousky, V.E. (1983), Two severe freezes in Brazil: precursors and synoptic evolution, Monthly weather review, Vol 111, 181-196.
- Hajbarpour, GH & Alijani., B. (2007), Synoptic analysis of frosts in Ardabil province, geography and development magazine,Vol.10, 89-106.
- Hamilton, M.G. & Tarifa, J.R. (1978), Synoptic aspect of a polar outbreak leading to frost in tropical Brazil, july 1972. Monthly weather review, Vol 106, 1545-1556.
- Linacre, ET and Geerts, B, (1997), Climates and Weather Explained, Routledge, London.
- Masoodian, S.A.,(2008), Recognition Isfahan weather Types, project of University of Isfahan.
- Masoodian, S.A, Ghaedi, S and Mohammadi, B., (2007), Regions of humidity of Iran, Literature and humanity Sciences Faculty journal of Tabriz, Geography, Vol.20, 1-13.
- Mojarad Ghareh Bagh, F., (1997), Analysis and Prediction of Frost in Azerbaijan, Ph.D. Thesis of physical geography (climatology), Faculty of humanities sciences, Tarbiat Modarres University, p. 186.
- Mojarad Karabakh, F. (1995). Analysis and forecasting frost in Azerbaijan. PhD thesis, Tarbiat Modarres University, Tehran, Iran.

- Müller GV & Berri GJ (2007), Atmospheric Circulation Associated with Persistent Generalized frosts in Central-Southern South America, Mon. Wea. Rev., 135, 4, 1268-1289.
- Müller GV, Nuñez MN & Seluchi M (2000), Relationship Between ENSO Cycles and Frost events Within the Pampa Humeda Region. Int. Journal of Climatology, 20, 1619-1637.
- Muller, G.V et al., (2003), Surface circulation associated with frost in the wet Pampas, International journal of climatology, Vol.23, 943-961.
- Müller, GV, Ambrizzi, T & Nuñez, MN (2005), Mean atmospheric circulation leading to generalized frosts in Central Southern South America, Theor. Appl. Climat, 82, 95-112.
- Müller GV and Ambrizzi T (2007), Teleconnection Patterns and

TC

Rossby Wave Propagation Associated to Generalized Frosts over Southern South America. Climate Dynamics, 29, 6, 633-645.

- Oke, T, (1987), Boundary Layer Climates, Routledge, London.
- Scholefield, P (1994), Frost is this the year?, Australian Grapegrower and Winemaker, 370, 31-64.
- Snyder, R.L., Melo-Albreu, J.P (2005), Frost Protection: fundamentals, practice and economics, Food and Agriculture Organization of the United Nations, Rome.
- Vassi, M (2008), Synoptic analysis of frost periods in Kurdistan province, MA thesis, climatology department, geography faculty, Tehran University.
- Vega, A.T., Robbons , K.D. & Grymes, J.M. (1994), Frost/Freeze analysis in the Southern climate region. Southern Regional Climate Center.