

Smoking Paradox at Cardiac Rehabilitation

Allahyar Golabchi

Cardiologist, Cardiac Rehabilitation Research Center, Isfahan Cardiovascular Research Institute, Isfahan University Of Medical Sciences, Isfahan, Iran

Date of Submission: Dec 23, 2011

Date of Acceptance: Jan 09, 2012

Correspondence to:

Dr. Allahyar Golabchi, Cardiac Rehabilitation Research Center, Isfahan Cardiovascular Research Institute, Isfahan University Of Medical Sciences, Isfahan, Iran. E-mail: golabchi@edc.mui.ac.ir

How to cite this article: Allahyar G, Smoking paradox at cardiac rehabilitation. Int J Prev Med 2012;3:139-40.

DEAR EDITOR,

Recently, I have read with interest the article by Mosayebi *et al.*^[1] entitled "The Effects of Cardiac Tertiary Prevention Program after Coronary Artery Bypass Graft Surgery on Health and Quality of Life" in your esteemed journal. The cardiac rehabilitation (CR) program significantly improves the functional capacity in postmyocardial infarction patients, especially those revascularized by coronary artery bypass graft surgery (CABG) or percutaneous coronary intervention (PCI).^[2] I would like to attract the authors' and readers' attention to some points that can be helpful in having high quality research.

The study was a prospective observational study with random sampling, carried out on 80 patients undergoing elective CABG. Half of the patients attended the CR program and the others did not. The outcomes evaluated were level of activity, the patients' employment status, and restoration to the previous level of performance in social activities after the surgery.^[1]

As can be observed in Table 1 of the mentioned article^[1], at the beginning of the study, there was a significant difference (P < 0.05) between the two groups in the number of smokers. This can impact on results of this study surely, so I believe they should match the smoking rate in two groups.

Additionally, in my literature review, I came to an interesting entity that I named it "smoking

paradox". Goel et al.^[3] showed that smoking is associated with decreased participation in CR after PCI. Consistently, Kerins et al.[4] reported that smoking was associated with people who do not attend or do not complete CR programs. In contrast, some robust studies such as the one performed by Fitzpatrick *et al.*^[5] demonstrated that the CR program leads to sustained improvements in major risk factors, particularly smoking. They reported that in their 3.5-year follow-up, there were significant absolute reductions from the baseline in smoking prevalence. Furthermore, in another robust study,^[6] 12-16% of patients were smokers at the CR program entry, which decreased to 5-7%at CR program discharge. This decrease had no correlation with the number of sessions attended to the CR program or days enrolled. Colivicchi et al.^[7] showed that participation in a postdischarge CR program was associated with a significant 26% reduction of the relative risk of smoking relapse within 1 year from the acute coronary syndrome (ACS) events. They finally advise that patients with ACS who have quit smoking during hospitalization should be encouraged to participate in the CR program, since they might derive additional benefits from prevention of smoking relapse.

REFERENCES

1. Mosayebi A, Javanmard SH, Mirmohamadsadeghi M, Rajabi R, Mostafavi S, Mansourian M. The

effects of cardiac tertiary prevention program after coronary artery bypass graft surgery on health and quality of life. Int J Prev Med 2011;2:269-74.

- Kargarfard M, Basati F, Sadeghi M, Rouzbehani R, Golabchi A. Effects of a cardiac rehabilitation program on diastolic filling properties and functional capacity in patients with myocardial infarction. J Isfahan Med Sch 2011;29:1-10.
- 3. Goel K, Lennon RJ, Tilbury RT, Squires RW, Thomas RJ. Impact of cardiac rehabilitation on mortality and cardiovascular events after percutaneous coronary intervention in the community clinical perspective. Circulation 2011;123:2344-52.
- 4. Kerins M, McKee G, Bennett K. Contributing factors to patient non-attendance at and non-

completion of Phase III cardiac rehabilitation. Eur J Cardiovasc Nurs 2011;10:31-6.

- Fitzpatrick P, Fitz-Simon N, Lonergan M, Collins C, Daly L. Heartwatch: The effect of a primary care-delivered secondary prevention programme for cardiovascular disease on medication use and risk factor profiles. Eur J Cardiovas Prev Rehabil 2011;18:129-35.
- Miller KM. Relationship between sessions attended and outcomes in outpatient cardiac rehabilitation. Available from: http://www.digital.library.wisc.edu/1793/54372 [Last accessed on 2011]
- Colivicchi F, Mocini D, Tubaro M, Aiello A, Clavario P, Santini M. Effect of smoking relapse on outcome after acute coronary syndromes. Am J Cardiol 2011;108:804-8.

Source of Support: Nil. Conflict of Interest: None declare.