



On the Occasion of World Cancer Day 2015; the Possibility of Cancer Prevention or Treatment with Antioxidants: The Ongoing Cancer Prevention Researches

Mahmoud Rafeian-Kopaie, Hamid Nasri¹

Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran, ¹Department of Nephrology, Division of Nephropathology, Isfahan University of Medical Sciences, Isfahan, Iran

Correspondence to:

Prof. Mahmoud Rafeian-Kopaie, Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran. E-mail: rafeian@yahoo.com

How to cite this article: Rafeian-Kopaie M, Nasri H. On the occasion of world cancer day 2015; the possibility of cancer prevention or treatment with antioxidants: The ongoing cancer prevention researches. *Int J Prev Med* 2015;6:108.

ABSTRACT

On February, 2014 World Cancer Day (WCD) was established to raise alertness of cancer and to encourage its prevention, detection, and treatment. In fact, WCD is celebrated every year on the 4th of February all over the world to commemorate all the accomplishments of the WHO. In this paper, we aimed to present the scientific evidence for the role of antioxidants in cancer. Damage to cells by reactive oxygen species, especially the damage to DNA, has been found to play a crucial role in the development of cancer. Exogenous antioxidants can prevent free radical damage associated with cancer development. However, whether or not taking dietary antioxidants can prevent or reduce the risk of developing cancer in humans is not clear. Some researchers have suggested that antioxidants counteract with drugs or toxins, which induce oxidative stress and hence prevent damage to cells or body organs.

Keywords: Antioxidant, cancer, herbs

INTRODUCTION

On February, 2014 World Cancer Day (WCD) was founded to raise alertness of cancer and to encourage its prevention, detection, and treatment.^[1,2] In fact, WCD is celebrated worldwide every year on the 4th of February to commemorate the cancer-related accomplishments of the WHO.^[3-5] The first WCD celebration was planned in Geneva, Switzerland in 1933 under the direction of Union for International Cancer Control with the support

of various other well-known cancer societies, treatment centers, research institutes, and patient groups.^[3,4] The WCD event was founded to help find the resources needed to fight and control this deadly disease.^[4-6] Moreover, the WCD is celebrated to help inform the public about cancer risk factors preventive measures, the benefits of early detection. In this review, we present the scientific evidence for the preventive role of antioxidants on cancer.^[7,8]

Free radicals and in particular reactive oxygen species (ROS) are formed naturally in the body when an atom or a molecule either loses or gains an electron. Free radicals play an important role in many normal cellular processes.^[9,10] However, at high concentrations, they can damage all major components of cells including proteins, DNA, and cell membranes. These damages to cells, especially the damage to DNA, can play a crucial role in the development of cancer.^[11] Moreover, some

Access this article online	
Quick Response Code: 	Website: www.ijpvmjournal.net/www.ijpm.ir
	DOI: 10.4103/2008-7802.169077

environmental toxins like cigarette smoke, may contain large amounts of free radicals or stimulate the body's cells to produce more free radicals.^[11]

The body makes some antioxidants, which are called endogenous antioxidants. However, the body also relies on exogenous sources of antioxidants for the rest of its protection against ROS and free radicals.^[12-14] If free radicals and ROS can cause cancer, antioxidants, therefore, should be able to prevent or inhibit this process.^[14,15]

Exogenous antioxidants can prevent free radical induced damage associated with cancer development.^[16-19] However, whether or not taking dietary antioxidants can prevent or reduce the risk of developing cancer in humans is not clear.^[20,21] It has been suggested that antioxidants counteract with drugs^[22,23] or toxins,^[20-23] which induce oxidative stress and hence prevent damage to cells or body organs.^[22-25]

In laboratory and preclinical studies in cancer prevention^[26,27] or immune system stimulation^[28] through medicinal plants with antioxidant activity, promising results have been achieved. However, the cohort and case-control studies, which have investigated the use of dietary antioxidants in the risk of cancer in humans have achieved mixed results.

Due to inadequate control for biases, which might influence the results, observational studies must be viewed with much caution. Randomized controlled clinical trials are considered to provide more reliable evidence of the harm or benefit of a health-related intervention. The results of the most important trials are summarized below.

THE LINXIAN GENERAL POPULATION NUTRITION INTERVENTION TRIALS

Effect of synthetic alpha-tocopherol

The first trial was a large-scale randomized trial investigating the effect of synthetic alpha-tocopherol on cancer risk in healthy Chinese people at increased risk of developing gastric and esophageal cancers.^[29] The subjects were randomly assigned to take a combination of 50 µg selenium, 15 mg beta-carotene, and 30 mg synthetic alpha-tocopherol, per day for 5 years or no supplement. The results of the study demonstrated that the risk of developing gastric cancer and/or esophageal cancer was not affected by antioxidant supplementation. However, people who took antioxidants had lower death due to gastric cancer but not for to esophageal cancer.^[29] In this study, 10 years after antioxidants supplementation ended, the reduced risk of gastric cancer death was no longer

found for those who took antioxidants, in comparison to those who did not.^[30]

Effect of alpha-tocopherol and beta-carotene

In the second trial, which was conducted in Finland, 5–8 years consumption of synthetic alpha-tocopherol and/or beta-carotene could reduce the incidence of different cancers in some middle-aged male smokers. The initial results of the study showed an increase in the incidence of lung cancer in the subjects who took 20 mg per day beta-carotene supplementation.^[31] There were no effects of alpha-tocopherol or beta-carotene supplementation on the incidence of renal pelvis, bladder, ureter, pancreas, colorectal, kidney, pharyngeal, laryngeal or esophageal cancers.^[32]

Effects of beta-carotene and Vitamin A on lung cancer

In a trial in the United States, daily supplementation with beta-carotene and Vitamin A on people who were at high risk of lung cancer due to having a history of smoking or exposure to asbestos showed that daily supplementation with both 25,000 IU retinol and 15 mg beta-carotene was associated with increased lung cancer and increased death from all-cause mortality.^[33] A later report in 2004 showed that the adverse effects persisted up to 6 years after the consumption of these supplements ended. However, the enhanced risks of all-cause mortality and lung cancer were no longer statistically significant.^[34]

In another study in 1996, 50 mg beta-carotene administration every other day for 12 years had no effect on cancer mortality, cancer incidence, and all-cause mortality among US male subjects.^[35]

Every other day administration of 50 mg beta-carotene, 600 IU Vitamin E, and 100 mg aspirin has also had no benefit or the incidence of cancer and cardiovascular diseases in US women of over 45 years.^[36] In 2005, similar results were published for Vitamin E consumption.^[37]

Effects of selenium and alpha-tocopherol on cancer in a trial published in 2004, daily supplementation with 100 µg selenium, 20 mg zinc, 6 mg beta-carotene, 30 mg synthetic Vitamin E, and 120 mg Vitamin C for a median of 7.5 years had no effect on the incidence of cardiovascular or cancer diseases or all-cause mortality, too.^[38]

In an international trial reported in 2005, no effect of daily supplementation with 400 IU synthetic alpha-tocopherol which was administered for 7 years was seen in the incidence of major cardiovascular events such as stroke, heart attack or death from heart disease, cancer incidence or death from cancer in people diagnosed with cardiovascular disease or diabetes.^[39]

The next trial was conducted in US, began in 2001 and was stopped in 2008. The trial investigated whether daily supplementation with 200 µg selenium, 400 IU synthetic Vitamin E or both would reduce the incidence of prostate cancer in men over 50 years. The results showed that the use of these supplements for a period of about 5.5 years did not affect the incidence of prostate or other cancers.^[39,40] Updated findings, reported in 2011, demonstrated that, following 1.5 years off supplements, the cases of prostate cancer among men taking Vitamin E alone were 17% more compared to placebo group.^[41] No increase in prostate risk was observed for men assigned to take Vitamin E plus selenium or selenium alone compared with men assigned to take a placebo.^[39]

In a trial which was conducted in 2009 the use of 500 mg Vitamin C, 400 IU Vitamin E or combination of the two, every other day for a median of 7.6 years did not reduce the incidence of prostate cancer or other cancers including leukemia, melanoma, lymphoma, and cancers of the bladder, pancreas, lung, colon, and rectum.^[42]

The above-randomized controlled clinical trials did not show that dietary antioxidant supplements are always beneficial in primary cancer prevention. However, it is possible that the lack of benefit in most of the clinical studies can be related to the effects of the tested antioxidants on cancer. They all were consumed as purified chemicals and their effects might be different when they are consumed as natural foods, which contain complex mixtures of minerals, vitamins, and various antioxidants. This acquires a more complete understanding of the antioxidant content of individual foods, how the various antioxidants interact with each other, and the factors that influence the uptake and distribution of food-derived antioxidants in the body. These are all active areas of ongoing cancer prevention research.^[42-45]

Another question, which might be raised is that whether or not people already diagnosed with cancer should take antioxidant supplements? Recent studies reported that antioxidant supplements during cancer treatment might alter the effectiveness or reduce the toxicity of specific therapies.^[46-53] Other trials have reported mixed results; some of them found that people who took antioxidants during cancer therapy had worse outcomes, especially if they were smokers.^[54-74]

Additional large randomized controlled studies are necessary to provide clear evidence about the benefits or harms of taking antioxidant supplements during cancer treatment.

CONCLUSIONS

In sum, a lot of *in vitro* studies have shown that exogenous antioxidants are able to help prevent the

free radical damage associated with the development of cancer. However, researches in humans have not demonstrated convincingly that taking antioxidant can reduce the risk of developing cancer, and some studies have even shown an increased risk for some cancers, particularly, in smokers.

Received: 01 Jun 14 **Accepted:** 30 Apr 15

Published: 04 Nov 15

REFERENCES

- Toh HC. World Cancer Day 2011: A world without cancer one day? *Ann Acad Med Singapore* 2011;40:65-6.
- Rafeian-Kopaei M, Nasri H. The ameliorative effect of *Zingiber officinale* in diabetic nephropathy. *Iran Red Crescent Med J* 2014;16:e11324.
- McAvoy B. Optimising cancer care in Australia. *Aust Fam Physician* 2003;32:369-72.
- Mortazavi M, Nasri H. Granulomatosis with polyangiitis (Wegener's) presenting as the right ventricular masses: A case report and review of the literature. *J Nephropathol* 2012;1:49-56.
- Nasri H. Atypical presentations of the sarcoidosis with kidney involvement. *J Renal Inj Prev* 2012;1:51-2.
- Hajivandi A, Amiri M. World Kidney Day 2014: Kidney disease and elderly. *J Parathyroid Dis* 2014;2:3-4.
- Nasri H, Baradaran A. The influence of serum 25-hydroxy vitamin D levels on *Helicobacter pylori* Infections in patients with end-stage renal failure on regular hemodialysis. *Saudi J Kidney Dis Transpl* 2007;18:215-9.
- Nasri H. Cisplatin and renal injury; current concepts. *J Renal Inj Prev* 2013;2:89-90.
- Setorki M, Rafeian-Kopaei M, Merikhi A, Heidarian E, Shahinfard N, Ansari R, et al. Suppressive impact of anethum graveolens consumption on biochemical risk factors of atherosclerosis in hypercholesterolemic rabbits. *Int J Prev Med* 2013;4:889-95.
- Nasri H. Reply: Oxytocin ameliorates cisplatin-induced nephrotoxicity in Wistar rats. *Ann Saudi Med* 2013;33:510-1.
- Ardalan MR, Sanadgol H, Nasri H, Baradaran A, Tamadon MR, Rafeian-Kopaei R. Vitamin D therapy in diabetic kidney disease; current knowledge on a public health problem. *J Parathyroid Dis* 2014;2:15-7.
- Valko M, Leibfritz D, Moncol J, Cronin MT, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol* 2007;39:44-84.
- Nasri H. Comment on: A model for prediction of cisplatin induced nephrotoxicity by kidney weight in experimental rats. *J Res Med Sci* 2013;18:1119-20.
- Nasri H, Nematbakhsh M, Ghobadi S, Ansari R, Shahinfard N, Rafeian-Kopaei M. Preventive and curative effects of ginger extract against histopathologic changes of gentamicin-induced tubular toxicity in rats. *Int J Prev Med* 2013;4:316-21.
- Bouayed J, Bohn T. Exogenous antioxidants – Double-edged swords in cellular redox state: Health beneficial effects at physiologic doses versus deleterious effects at high doses. *Oxid Med Cell Longev* 2010;3:228-237.
- Nasri H, Ardalan MR, Rafeian-Kopaei M. Comment on: Prevention of renal damage by treating hyperuricemia. *Int J Prev Med* 2015;6:10.
- Rafeian-Kopaei M, Baradaran A, Rafeian M. Plants antioxidants: From laboratory to clinic. *J Nephropathol* 2013;2:152-3.
- Ardalan MR, Sanadgol H, Nasri H, Baradaran A, Tamadon MR, Rafeian-Kopaei R. Impact of vitamin D on the immune system in kidney disease. *J Parathyroid Dis* 2013;1:17-20.
- Baradaran A, Nasri H, Nematbakhsh M, Rafeian-Kopaei M. Antioxidant activity and preventive effect of aqueous leaf extract of Aloe Vera on gentamicin-induced nephrotoxicity in male Wistar rats. *Clin Ter* 2014;165:7-11.
- Nasri H, Rafeian-Kopaei M. Herbal medicine and diabetic kidney disease. *J Nephropathol* 2013;2:1-2.
- Shirzad M, Kordyazdi R, Shahinfard N, Nikokar M. Does Royal jelly affect tumor cells? *J HerbMed Pharmacol* 2013;2:45-8.

22. Nasri H, Rafeian-Kopaei M. Protective effects of herbal antioxidants on diabetic kidney disease. *J Res Med Sci* 2014;19:82-3.
23. Nasri H. Comment on: A model for prediction of cisplatin induced nephrotoxicity by kidney weight in experimental rats. *J Res Med Sci* 2013;18:1119-20.
24. Tagne RS, Telefo BP, Nyemb JN, Yemele DM, Njina SN, Goka SM, et al. Anticancer and antioxidant activities of methanol extracts and fractions of some Cameroonian medicinal plants. *Asian Pac J Trop Med* 2014;7S1:S442-7.
25. Nasri H, Baradaran A, Shirzad H, Rafeian-Kopaei M. New concepts in nutraceuticals as alternative for pharmaceuticals. *Int J Prev Med* 2014;5:1487-99.
26. Mosavel M, Simon C, Oakar C, Meyer S. Cervical cancer attitudes and beliefs – A Cape Town community responds on World Cancer Day. *J Cancer Educ* 2009;24:114-9.
27. Ardalan MR, Rafeian-Kopaei M. Is the safety of herbal medicines for kidneys under question? *J Nephro pharmacol* 2013;2:11-2.
28. Ardalan MR, Khodaie L, Nasri H, Jouyban A. Herbs and hazards: Risk of aristolochic acid nephropathy in Iran. *Iran J Kidney Dis* 2015;9:14-7.
29. Cannon G, Gupta P, Gomes F, Kerner J, Parra W, Weiderpass E, et al. Prevention of cancer and non-communicable diseases. *Asian Pac J Cancer Prev* 2012;13:3-11.
30. Rafeian-Kopaei M, Nasri H, Baradaran A. *Teucrium polium*: Liver and kidney effects. *J Res Med Sci* 2014;19:478-9.
31. Nasri H, Rafeian-Kopaei M. World cancer day 2014: "Increasing the awareness". *Cell J* 2014;16:383-4.
32. Nematbakhsh M, Nasri H. The effects of vitamin E and selenium on cisplatin-induced nephrotoxicity in cancer patients treated with cisplatin-based chemotherapy: A randomized, placebo-controlled study. *J Res Med Sci* 2013;18:626-7.
33. Nasri H. Cisplatin therapy and the problem of gender-related nephrotoxicity. *J Nephro pharmacol* 2013;2:13-4.
34. Baradaran A, Nasri H, Rafeian-Kopaei M. Oxidative stress and hypertension: Possibility of hypertension therapy with antioxidants. *J Res Med Sci* 2014;19:358-67.
35. Shirzad H, Tajji F, Rafeian-Kopaei M. Correlation between antioxidant activity of garlic extracts and WEHI-164 fibrosarcoma tumor growth in BALB/c mice. *J Med Food* 2011;14:969-74.
36. Shirzad H, Kiani M, Shirzad M. Impacts of tomato extract on the mice fibrosarcoma cells. *J HerbMed Pharmacol* 2013;2:13-6.
37. Shirzad H, Shahrani M, Rafeian-Kopaei M. Comparison of morphine and tramadol effects on phagocytic activity of mice peritoneal phagocytes *in vivo*. *Int Immunopharmacol* 2009;9:968-70.
38. Blot WJ, Li JY, Taylor PR, Guo W, Dawsey S, Wang GQ, et al. Nutrition intervention trials in Linxian, China: Supplementation with specific vitamin/mineral combinations, cancer incidence, and disease-specific mortality in the general population. *J Natl Cancer Inst* 1993;85:1483-92.
39. Nasri H. C-Phycocyanin attenuates cisplatin-induced nephrotoxicity in mice. *Ren Fail* 2013;35:1054-5.
40. Nematbakhsh M, Nasri H. Re: Effect of alcoholic extract of *Nigella sativa* on cisplatin-induced toxicity in rat. *Iran J Kidney Dis* 2013;7:165-6.
41. Nasri H. Protective effects of subchronic caffeine administration on cisplatin induced urogenital toxicity in male mice. *Indian J Exp Biol* 2013;51:406.
42. Qiao YL, Dawsey SM, Kamangar F, Fan JH, Abnet CC, Sun XD, et al. Total and cancer mortality after supplementation with vitamins and minerals: Follow-up of the Linxian General Population Nutrition Intervention Trial. *J Natl Cancer Inst* 2009;101:507-18.
43. Setorki M, Rafeian-Kopaei M, Merikhi A, Heidarian E, Shahinfard N, Ansari R, et al. Suppressive impact of anethum graveolens consumption on biochemical risk factors of atherosclerosis in hypercholesterolemic rabbits. *Int J Prev Med* 2013;4:889-95.
44. Baradaran A, Nasri H, Rafeian-Kopaei M. Oxidative stress and hypertension: Possibility of hypertension therapy with antioxidants. *J Res Med Sci* 2014;19:358-67.
45. Nematbakhsh M, Nasri H. Cisplatin nephrotoxicity may be sex related. *Kidney Int* 2013;83:1201.
46. Baradaran A, Behradmanesh S, Nasri H. Association of body mass index and serum vitamin D level in healthy Iranian adolescents. *Endokrynol Pol* 2012;63:29-33.
47. Nasri H. Hypertension and renal failure with right arm pulse weakness in a 65 years old man. *J Nephro pathol* 2012;1:130-3.
48. The effect of vitamin E and beta carotene on the incidence of lung cancer and other cancers in male smokers. The Alpha-Tocopherol, Beta Carotene Cancer Prevention Study Group. *N Engl J Med* 1994;330:1029-35.
49. Wright ME, Virtamo J, Hartman AM, Pietinen P, Edwards BK, Taylor PR, et al. Effects of alpha-tocopherol and beta-carotene supplementation on upper aerodigestive tract cancers in a large, randomized controlled trial. *Cancer* 2007;109:891-8.
50. Omenn GS, Goodman GE, Thornquist MD, Balmes J, Cullen MR, Glass A, et al. Effects of a combination of beta carotene and vitamin A on lung cancer and cardiovascular disease. *N Engl J Med* 1996;334:1150-5.
51. Goodman GE, Thornquist MD, Balmes J, Cullen MR, Meyskens FL Jr, Omenn GS, et al. The Beta-Carotene and Retinol Efficacy Trial: Incidence of lung cancer and cardiovascular disease mortality during 6-year follow-up after stopping beta-carotene and retinol supplements. *J Natl Cancer Inst* 2004;96:1743-50.
52. Hennekens CH, Buring JE, Manson JE, Stampfer M, Rosner B, Cook NR, et al. Lack of effect of long-term supplementation with beta carotene on the incidence of malignant neoplasms and cardiovascular disease. *N Engl J Med* 1996;334:1145-9.
53. Lee IM, Cook NR, Manson JE, Buring JE, Hennekens CH. Beta-carotene supplementation and incidence of cancer and cardiovascular disease: The Women's Health Study. *J Natl Cancer Inst* 1999;91:2102-6.
54. Lee IM, Cook NR, Gaziano JM, Gordon D, Ridker PM, Manson JE, et al. Vitamin E in the primary prevention of cardiovascular disease and cancer: The Women's Health Study: A randomized controlled trial. *JAMA* 2005;294:56-65.
55. Hercberg S, Galan P, Preziosi P, Bertrais S, Mennen L, Malvy D, et al. The SU.VI.MAX Study: A randomized, placebo-controlled trial of the health effects of antioxidant vitamins and minerals. *Arch Intern Med* 2004;164:2335-42.
56. Lippman SM, Klein EA, Goodman PJ, Lucia MS, Thompson IM, Ford LG, et al. Effect of selenium and vitamin E on risk of prostate cancer and other cancers: The Selenium and Vitamin E Cancer Prevention Trial (SELECT). *JAMA* 2009;301:39-51.
57. Nasri H. Re: Effect of silymarin on streptozotocin-nicotinamide-induced type 2 diabetic nephropathy in rats. *Iran J Kidney Dis* 2013;7:414-5.
58. Klein EA, Thompson IM Jr, Tangen CM, Crowley JJ, Lucia MS, Goodman PJ, et al. Vitamin E and the risk of prostate cancer: The Selenium and Vitamin E Cancer Prevention Trial (SELECT). *JAMA* 2011;306:1549-56.
59. Gaziano JM, Glynn RJ, Christen WG, Kurth T, Belanger C, MacFadyen J, et al. Vitamins E and C in the prevention of prostate and total cancer in men: The Physicians' Health Study II randomized controlled trial. *JAMA* 2009;301:52-62.
60. Ghafari M, Taheri Z, Hajivandi A, Amiri M. Parathyroid carcinoma; facts and views. *J Parathyroid Dis* 2015;3:37-40.
61. Nasri H. Sudden onset of renal failure requiring dialysis associated with large B-cell lymphoma of colon. *J Nephro pathol* 2012;1:202-6.
62. Maghsoudi AR, Baradaran-Ghahfarokhi M, Ghaed-Amini F, Nasri H, Dehghani Mobarakeh M, Rafeian-Kopaei M. Renal failure and submental lymphadenopathy in a 68 years old woman. *J Nephro pathol* 2012;1:198-201.
63. Nasri H, Tavakoli M, Ahmadi A, Baradaran A, Nematbakhsh M, Rafeian-Kopaei M. Ameliorative effect of melatonin against contrast media induced renal tubular cell injury. *Pak J Med Sci* 2014;30:261-5.
64. Baradaran A, Nasri H, Rafeian-Kopaei M. Comment on: Anti-oxidative stress activity of *Stachys lavandulifolia* aqueous extract in humans. *Cell J* 2013;15:272-3.
65. Nasri H, Madihi Y, Marikhi A. Commentary on: Effects of cinnamon consumption on glycemic status, lipid profile and body composition in type 2 diabetic patients. *Int J Prev Med* 2013;4:618-9.
66. Nasri H. Effect of garlic extract on blood glucose level and lipid profile in normal and alloxan diabetic rabbits. *Adv Clin Exp Med* 2013;22:449-50.
67. Wiseman M. The second World Cancer Research Fund/American Institute for Cancer Research expert report. Food, nutrition, physical activity, and the prevention of cancer: A global perspective. *Proc Nutr Soc* 2008;67:253-6.
68. Ardalan MR. Parathyroid carcinoma in hemodialysis patients; it should not be diagnosed as a thyroid nodule. *J Parathyroid Dis* 2013;1:25-6.
69. Ardalan MR, Nasri H, Rafeian-Kopaei M. Comment on: Protective

- role of recombinant human erythropoietin in kidney and lung injury following renal bilateral ischemia-reperfusion in Rat Model. *Int J Prev Med* 2013;4:1226-7.
70. Nasri H. Antioxidants for prevention of gentamicin-induced nephrotoxicity. *Iran J Kidney Dis* 2014;8:1-2.
71. Nasri H, Rafeian-Kopaei M. Oxidative stress and aging prevention. *Int J Prev Med* 2013;4:1101-2.
72. Nasri H, Madihi Y, Marikhi A. Commentary on: Effects of cinnamon consumption on glycemic status, lipid profile and body composition in type 2 diabetic patients. *Int J Prev Med* 2013;4:618-9.
73. Nasri H, Rafeian-Kopaei M. Protective effects of herbal antioxidants on diabetic kidney disease. *J Res Med Sci* 2014;19:82-3.
74. Nasri H, Nematbakhsh M, Rafeian-Kopaei M. Ethanol extract of garlic for attenuation of gentamicin-induced nephrotoxicity in Wistar rats. *Iran J Kidney Dis* 2013;7:376-82.

Source of Support: Nil, Conflict of Interest: None declared.

Archive of SID