

Simplified Algorithm for Evaluation of Proteinuria in Clinical Practice: How should A Clinician Approach?

Abstract

Background: Proteinuria is a common laboratory finding among children and adolescents. It can be identified as either a transient or a persistent finding and can represent a benign condition or a serious disease. **Methods:** Pertinent medical literature for asymptomatic proteinuria in children and adolescents published in English was searched between January 1980 and May 2017 using PubMed, MEDLINE, EMBASE, and Google Scholar research databases. Of the 64 reviewed articles, 24 studies were eligible for inclusion. **Results:** Random spot urine protein-to-creatinine (PCR) ratio is widely used to reliably detect proteinuria. The normal value for the spot PCR in children aged 2 years or older is less than 0.3. In children aged below 2 years, the PCR can be as high as 0.5. Orthostatic proteinuria is defined as urine PCR greater than 0.3 detected in a urine specimen during the daytime activity but less than 0.3 on the first morning void specimen. PCR above 3.0 signifies heavy proteinuria as seen in nephrotic syndrome. Orthostatic proteinuria is a frequent cause of proteinuria in asymptomatic children and adolescents, which require no specific therapy except for health maintenance follow-up. Pediatric nephrologist referral is indicated when the proteinuria is constant and persists over 6 months or is associated with hematuria, hypertension, or renal dysfunction. **Conclusions:** We provide a simplified diagnostic algorithm for evaluation of proteinuria in primary care adolescents who appear well and in whom proteinuria is incidentally discovered during a routine examination.

Keywords: Adolescents, algorithm, asymptomatic proteinuria, children

Introduction

Proteinuria is a frequent laboratory finding both in the outpatient clinic and inpatient settings. The finding of protein in the urine may reflect either a benign finding or significant renal functional disorders or structural abnormalities.^[1-4] This distinction is possible by carefully taking the patient's medical history, performing through physical examination, and ordering appropriate laboratory investigations.

Evaluation of proteinuria should begin with a careful history and thorough physical examination, urine microscopic examination, and determination of the amount of protein excretion rate (PER). The PER has been traditionally measured using 24-h urine collections. However, the collection of 24-h urine is often cumbersome, and spot urinary protein-to-creatinine ratio (PCR), expressed in g/g or mg/mg, has become a simple and attractive yet reliable alternative.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

A spot urine PCR has been found to have a significant linear correlation with a 24-h urine PCR.^[5-9] Furthermore, because the PCR compares urinary protein concentration with urinary creatinine concentration, urinary dilution or concentration does not influence this value.

We present a simple practical diagnostic approach to differentiate benign proteinuria from proteinuria resulting from glomerular disease in children and adolescents who present with asymptomatic isolated proteinuria.

Methods

Pertinent medical literature for proteinuria published in English was searched from January 1980 to May 2017 using PubMed, MEDLINE, EMBASE, and Google Scholar research databases, and then the search was extended as linked citations indicated. The search terms included *asymptomatic proteinuria, isolated proteinuria, children, adolescents, evaluation, and management*. Of the 64 reviewed articles, 24 studies were eligible for inclusion.

How to cite this article: Mazaheri M, Assadi F. Simplified algorithm for evaluation of proteinuria in clinical practice: How should a clinician approach?. *Int J Prev Med* 2019;10:35.

Mojgan Mazaheri, Farahnak Assadi¹

Departments of Pediatrics, Section of Nephrology, Semnan University of Medical Sciences, Semnan, Iran, ¹Department of Pediatrics, Section of Nephrology, Rush University Medical Center, Chicago, Illinois, USA

Address for correspondence:

Dr. Farahnak Assadi,
445 E. North Water Street,
Suite 1804, Chicago, IL USA.
E-mail: fassadi@rush.edu

Access this article online

Website:
www.ijpvmjournal.net/www.ijpvm.ir

DOI:
10.4103/ijpvm.IJPVM_557_18

Quick Response Code:



Archive of SID

Results

The initial evaluation of proteinuria should include a complete history, including a family history of renal disease, recent upper respiratory infections, gross hematuria, and changes in urine output. The physical examination should include urinalysis and examination of urine sediment; measurements of height, weight, and blood pressure; identification of edema, ascites, and skin pallor; and palpation of the kidneys.

Orthostatic proteinuria is a benign medical condition with excellent long-term prognosis, and it does not warrant an extensive workup. Patients with constant proteinuria, which persists beyond 6 months, or proteinuria associated with hematuria, hypertension, or abnormal renal function may require referral to a pediatric nephrologist.

Discussion

We aimed to provide an organized practical approach for the primary care adolescents to evaluate a child with proteinuria that emphasizes common conditions and stepwise laboratory and radiologic investigations [Figure 1].

Urinary PER in the normal child is less than 100 mg/m² per day or a total of 150 mg per day. Healthy children excrete small amounts of protein in their urine, which varies with age and the size of the child.^[2-4] In neonates, normal urinary protein excretion is higher, up to 300 mg/m² per day, because of reduced reabsorption of filtered protein. In children, PER decreases progressively with age until late childhood till it reaches the adult levels of less than 150 mg/m² per day [Table 1].^[2-4]

The urine dipstick method is widely used in routine office evaluation of proteinuria. The urine protein dipstick is sensitive to the albumin concentration of the urine. It can yield both false-negative and false-positive results in the presence of dilute (specific gravity less than 10.10) or alkaline urine (pH greater than 7.0), respectively.^[7]

The 24-h urine collection is traditionally used for the urinary protein quantitation. The 24-h urine collection can be done as a single collection or two split collections during the daytime activity and overnight recumbent positions.^[6,7] However, the 24-h timed urine collections, particularly, in small children, cumbersome and often are flawed with collection errors.

Furthermore, PER can vary significantly from day to day and is also influenced by diet, posture, exercise, and body size. Does a 10-kg patient seem as “nephrotic” at 0.5 g per day protein excretion compared with the 80-kg patient at 4.0 g per day? Does the 80-kg patient have eight times worse glomerular disease and is this person eight times as nephrotic [Table 2].

The answer is “NO!” These value by themselves, while each a eightfold difference, are similar in each of the two

Table 1: Normal urinary protein excretion in infants and children 90

	Total protein mg per day	Total protein mg/m ² per day	Range mg per day
Premature <1 month	29	182	88-377
Full term <1 month	32	145	68-309
1-12 months	38	109	48-244
1-4 years	49	91	7-223
4-10 years	71	85	1-234
10-16 years	83	63	2-181

Table 2: Comparison of creatinine excretion, protein excretion, and protein-to-creatinine ratio in relation to increasing weight*

Weight	Timed (24-h) urine collection		
	CrE, mg per day	PE, mg per day	Spot P/Cr, mg/mg
10 kg	200	500	2.5
20 kg	400	1000	2.5
40 kg	800	2000	2.5
80 kg	1600	4000	2.5

*Assuming that the patient excretes 20 mg/kg of creatinine a day. CrE=Creatinine excretion, PE=Protein excretion, P/Cr=Protein-creatinine ratio

patients and have the same clinical significance as their PCR values are similar, 2.5 mg/mg [Table 2]. The ease and simplicity of obtaining first morning spot urine PCR value allows one to exactly estimate the 24-h PER and automatically adjust values for patient size.^[10]

A random spot PCR is a good representative of 24-h urine PCR, and is now widely used in children in lieu of a 24-h urine collection because of its convenience and simplicity.^[5-9] One of the advantages of measuring the spot untimed urine PCR over a 24-h urine protein measurement is that the urinary concentration or dilution does not affect its values as the PCR compares urinary protein concentration with urinary creatinine concentration, and as a result, the urinary concentration or dilution does not affect its value.^[9,11]

The KDIGO guidelines (Kidney Disease: Improving Global Outcomes) reports that there are insufficient data to recommend 24-h or spot urine for PER.^[12]

More recent studies have compared the first morning void PCR and random single void PCR with the PCR of a timed urine collection ranging from 4 to 24 h to determine collection accuracy in PER, documenting the accuracy and reliability of random single void for determination of PER.^[10,13,14] Because a false-positive result may occur if urine is highly diluted, the PCR should be performed on a first-voided morning specimen rather than the one collected randomly later in the day.^[9,10,12]

The normal PCR in children and adolescents is less than 0.3. In infants and younger children, the PCR is higher

Archive of SID

with the upper normal limit of 0.5. PCR above 3.0 is found in patients with nephrotic syndrome.^[9] The daily PER can

be determined from spot urine PCR, based on sex, age, and weight using the following equations.^[9,10,13-15]

$$\text{PER (g/m}^2 \text{ per day)} = 0.63 \times (\text{PCR})$$

Many clinical disorders are associated with proteinuria [Table 3]. Transient proteinuria is benign and non-pathologic and is usually after infantile febrile seizure, exposure to rigid cold, and strenuous exercise.^[16,17] These conditions require no specific therapy and need only health maintenance follow-up.

Orthostatic proteinuria is diagnosed when the PCR is greater than 0.3 in a urine specimen tested during the daytime activity but less than 0.3 when the urine is collected after the nighttime recumbent position [Table 3].^[16,17]

Isolated persistent proteinuria lasting more than 6 months or proteinuria complicated with hematuria, hypertension, or abnormal renal function usually associated with glomerular lesions or congenital kidney and urinary tract anomalies such as unilateral kidney agenesis, obstructive hydronephrosis, and reflux nephropathy, which often require further evaluations including renal ultrasonography and voiding cystourethrogram (VCUG) [Table 4].^[18-23] If proteinuria is associated glomerulonephritis, then referral to a pediatric nephrologist is warranted for possible renal biopsy indication [Figure 1].^[22,23] Further therapeutic regimen with immunosuppressive medications, inhibition of angiotensin converting enzyme (ACE) or angiotensin receptor blockade (ARB), may be indicated to slowing the progression of glomerular disease.^[24]

Conclusions

We have developed a simple and yet cost-effective diagnostic algorithm that is based on determinations of random urine PCR to differentiate the multiple causes of proteinuria in children and adolescents, in a step-by-step fashion [Figure 1]. The ease and simplicity of obtaining first morning spot urine PCR value allows one to exactly estimate the 24-h PER. Utilization of this approach brings

Table 3: Etiologic classification of proteinuria among children and adolescents

Transient proteinuria
Fever
Strenuous exercise
Extreme cold exposure
Epinephrine administration
Emotional stress
Congestive heart failure
Seizures
Abdominal surgery
Isolated asymptomatic proteinuria
Orthostatic proteinuria
Persistent fixed proteinuria
Proteinuria secondary to renal disease
Minimal change nephrotic syndrome
Focal segmental glomerulosclerosis (FSGS)
Acute postinfectious glomerulonephritis
Membranoproliferative glomerulonephritis
Membranous glomerulonephritis
Lupus nephritis
Henoch-Schönlein purpura (HSP)
Human immunodeficiency virus (HIV)-associated nephropathy
Hemolytic uremic syndrome
Vasculitis
Chronic interstitial nephritis
Renal structural abnormalities
Hydronephrosis
Cystic kidney disease
Reflux nephropathy
Renal dysplasia
Unilateral kidney agenesis

Table 4: Clinical correlations in proteinuria

Likely diagnosis	History/physical examination	Cr ^a	Blood Albumin	C3 ^b	Pr/Cr ^c	Other
Orthostatic proteinuria	>10 years of age	Normal	Normal	Normal	<1.0	-
Nephrotic syndrome	Edema <6 years of age	Normal	Low	Normal	>3.0	High cholesterol
MCD ^d						
Acute GN ^e	Edema, gross hematuria	High or low	Normal or low	Low	<3.0	High ASO ^f titer, HTN ^g
FSGS ^h , MGN ⁱ	Hematuria, HTN	High or low	Normal or low	Normal	>1.0	High cholesterol
MPGN ^j	Hematuria, HTN	High or low	Normal or low	Low	>1.0	
Lupus nephritis	Rash, arthritis	Normal or low	Normal or low	Low	>1.0	High ANA ^k
HSP ^l	Hematuria	Normal	Normal	Normal	>1.0	-
Tubulointerstitial disease	Polyuria	Normal	Normal	Normal	<1.0	-

^acreatinine, ^bcomplement-3, ^cprotein/creatinine ratio, ^dminimal change disease, ^eglomerulonephritis, ^fantistreptolysin-O, ^ghypertension, ^hfocal and segmental glomerulosclerosis, ⁱmembranous glomerulonephritis, ^jmembranoproliferative glomerulonephritis, ^kantinuclear antibody, ^lHenoch-Schönlein purpura




Figure 1: Simplified algorithm for the evaluation of proteinuria. ^aProtein/creatinine ratio, ^bantistreptolysin O, ^chypertension, ^dantinuclear antibody, ^ehypertension, ^fglomerular filtration rate, ^gminimal change nephrotic syndrome, ^hmembranoproliferative glomerulonephritis, ⁱfocal segmental glomerulosclerosis, ^jsystemic lupus erythematosus, ^kimmunoglobulin-A glomerulonephritis, ^lvesicoureteral reflux, ^mhuman immunodeficiency virus, ⁿHenoch-Schönlein Purpura, ^opost-infectious glomerulonephritis, ^pmembranous glomerulonephritis, ^qcystic kidney disease

a greater clarity and simplicity for evaluation of patients with asymptomatic isolated proteinuria.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Received: 08 Nov 18 Accepted: 14 Dec 18

Published: 05 Mar 19

References

1. Quigley R. Evaluation of hematuria and proteinuria: How should

a pediatrician proceed? *Curr Opin Pediatr* 2008;20:140-4.

2. Leung KA, Wong AH. Proteinuria in children. *Am Fam Physician* 2010;82:645-51.

3. Thoai LH, Cachat F, Guignard JP. Proteinuria in children: Practical approach. *Rev Med Suisse Romande* 2000;120:245-50.

4. Ariceta G. Clinical practice: Proteinuria. *Eur J Pediatr* 2011;170:15-20.

5. Mori Y, Hiraoka M, Sganuma N, Tsukahara H, Yoshida H, Mayumi M. Urinary creatinine excretion and protein/creatinine ratios vary by body size and gender in children. *Pediatr Nephrol* 2006;21:683-7.

6. Schwab S, Christensen RL, Dougherty K, Klahr S. Quantitation of proteinuria by the use of protein-to-creatinine ratios in single urine samples. *Arch Intern Med* 1987;147:943-4.

Archive of SID

7. Shidham G, Hebert L. Timed urine collections are not needed to measure urine protein excretion in clinical practice. *Am J Kidney Dis* 2006;47:8-14.
8. Assadi F. Quantitation of microalbuminuria using random urine sample. *Pediatr Nephrol* 2002;17:107-10.
9. Ginsberg JM, Chang BS, Matarese RA, Garella S. Use of single voided urine samples to estimate quantitative proteinuria. *N Engl J Med* 1983;309:1543-6.
10. Rodby RA. Timed urine collections for albumin and protein: The king is dead, long live the king. *Am J Kidney Dis* 2016;68:836-8.
11. Guedes Marques M, Cotovio P, Ferrer F, Silva C, Botelho C, Lopes K, *et al.* Random spot urine protein/creatinine ratio: A reliable method for monitoring lupus nephritis? *Clin Kidney J* 2013;6:590-4.
12. KDIGO clinical practice guidelines for glomerulonephritis. *Kidney Int Suppl* 2012;2:157.
13. Hirano D, Fujinaga S, Shinozaki T, Endo A, Watanabe T, Murakami H, *et al.* Optimal urinary protein-to creatinine ratio as a renal biopsy criterion in children with asymptomatic proteinuria. *Clin Nephrol* 2014;82:115-21.
14. Fotheringham J, Campbell MJ, Fogarty DG, EL Nahas M, Ellam T. Estimated albumin excretion rate versus urine albumin creatinine ratio for the estimation of measured albumin excretion rate: Derivation and validation of an estimated albumin excretion equation. *Am J Kidney Dis* 2014;63:405-14.
15. Teo BW, Loh PT, Wong WK, Ho PJ, Choi KP, Toh QC, *et al.* Spot urine estimations are equivalent to 24-hour urine assessments of urine protein excretion for predicting clinical outcomes. *Int J Nephrol* 2015;2015:156484.
16. Rytand DA, Spreiter S. Prognosis in postural (orthostatic) proteinuria: Forty to fifty-year follow-up of six patients after diagnosis by Thomas Addis. *N Engl J Med* 1981;305:618-21.
17. Sebestyn JF, Alom US. The teenager with asymptomatic proteinuria: Think orthostatic first. *Clin Pediatr (Phila)* 2011;50:179-82.
18. Hogg RJ, Portman RJ, Milliner D, Lemley KV, Eddy A, Ingelfinger J. Evaluation and management of proteinuria and nephrotic syndrome in children: Recommendations from a pediatric nephrology panel established at the National Kidney Foundation conference on proteinuria, albuminuria, risk, assessment, detection, and elimination (PARADE). *Pediatrics* 2000;105:1242-9.
19. Ardissino G, Testa S, Dacco V, Viganò S, Taioli E, Claris-Appiani A, *et al.* Proteinuria as a predictor of disease progression in children with hypoplastic nephropathy. Data from Ital Kid Project. *Pediatr Nephrol* 2004;19:172-7.
20. Springberg PD, Farrett LE Jr, Thompson AL Jr, Collins NF, Lordon RE, Robinson RR. Fixed and reproducible orthostatic proteinuria: Results of a 20-year follow-up study. *Ann Intern Med* 1982;97:516-9.
21. Assadi F. Value of urinary excretion of microalbuminuria in predicting glomerular lesions in children with isolated microscopic hematuria. *Pediatr Nephrol* 2005;20:1131-5.
22. Trachtman H, Bergwerk A, Gauthier B. Isolated proteinuria in children. Natural history and indications for renal biopsy. *Clin Pediatr* 1994;33:468-72.
23. Coppo R. How early renal biopsy has to be performed in children with isolated asymptomatic proteinuria? *Nephrol Dial Transplant* 2012;27:3016-7.
24. Ruggenetti P. Dual renin-angiotensin system blockade for nephroprotection. *Nephro Ther Suppl* 2017;1:S43-5.

