



Efficacy of Pre-Synchronization and CIDR on the Outcome of Short-Term Synchronization Program in Zandi Ewes During the Breeding Season

Reza Arjmandi*, Mahdi Vodjgani, Amir Niasari Naslaji, Faramarz Gharagozlou, Vahid Akbarinejad

Department of Theriogenology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

Abstract

BACKGROUND: Timed breeding programs are essential to implementing extensive artificial insemination (AI) programs in sheep.

OBJECTIVES: This study aimed to evaluate whether application of pre-synchronization and controlled internal drug releasing (CIDR) before and during fixed time AI protocol, respectively, could enhance estrus synchronization and fertility of ewes.

METHODS: A total of 120 ewes were randomly assigned into four experimental groups (n=30 in each group) considering age, weight, and body condition score (BCS). All ewes received GnRH (25 µg of alarelin acetate), and five days afterwards, PGF₂α (75 µg d-cloprostenol) plus eCG (400 IU). In the control group, ewes received no additional treatment. In Pre-synch group, ewes received two injections of PGF₂α at 9-day interval three days before GnRH administration of main estrus synchronization protocol. In CIDR group, ewes received 5-day CIDR between GnRH and PGF₂α of main estrus synchronization protocol. In Pre-synch-CIDR group, ewes received both two injections of PGF₂α at 9-day interval and 5-day CIDR. Blood serum progesterone concentrations were measured in all ewes prior to injection of PGF₂α (day 5). All ewes were subjected to fixed time laparoscopic AI 48 hours after administration of the last PGF₂α.

RESULTS: No interaction was found between CIDR and pre-synchronization protocols ($P>0.05$). Progesterone concentration was higher in the CIDR groups than in groups without CIDR ($P<0.0001$). Estrous cycle was not affected by pre-synchronization and CIDR ($P>0.05$). The estrus was earlier in ewes with pre-synchronization compared to ewes without pre-synchronization following the last injection of prostaglandin ($P=0.022$). Pregnancy rate, lambing rate, prolificacy rate, fecundity rate, lamb weight at birth, and lamb gender were not significantly different between the treatment groups ($P>0.05$).

CONCLUSIONS: In our study, estrus rate and reproductive parameters showed no significant differences between different groups, although pre-synchronization advanced onset of estrus expression.

KEYWORDS: Breeding season, CIDR, Ewe, Pre-synchronization, Short-term synchronization

Correspondence

Reza Arjmandi, Department of Theriogenology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran
Tel: +98 (011) 43232386, Fax: +98 (011) 43232386, Email: r.arjomandi@ut.ac.ir

Received: 2020-08-04

Accepted: 2020-10-28

Copyright © 2021. This is an open-access article distributed under the terms of the Creative Commons Attribution- 4.0 International License which permits Share, copy and redistribution of the material in any medium or format or adapt, remix, transform, and build upon the material for any purpose, even commercially.

How to Cite This Article

Arjmandi, R., Vodjgani, M., Niasari Naslaji, A., Gharagozlou, F., Akbarinejad, V., A. (2021). Efficacy of Pre-Synchronization and CIDR on the Outcome of Short-Term Synchronization Program in Zandi Ewes During the Breeding Season. *Iranian Journal of Veterinary Medicine*, 15(1), 46-55.

Introduction

In recent years, there has been an increasing demand for the use of various breeds in domestic sheep herds to implement breeding programs. The high price of imported rams, the sensitivity and mortality of these rams due to the various diseases in domestic herds, as well as the presence of rump in the most Iranian breeds as a cause of inefficient mating prevent the extensive use of such breeds of rams in the Iranian herds (Vodjgani *et al.*, 2017). Artificial insemination (AI) can be applied as a useful method in breeding programs (Sathe, 2018). Due to the special anatomy of ewes' cervix, it is difficult to access inside the uterus and thus, insemination of ewes with frozen semen has been encountered with a big challenge (Sathe, 2018; Noakes *et al.*, 2019).

Laparoscopic AI is currently one of the well-known intrauterine methods of insemination with frozen sperm in sheep (Casali *et al.*, 2017). Successful synchronization of estrus and ovulation is one of the major contributing factors affecting conception rate following laparoscopic AI in sheep (Olivera–Muzante *et al.*, 2019).

In this context, various estrus synchronization protocols have been used in sheep including application of 12- or 14-day progestogen-dependent protocols (Abecia *et al.*, 2012; Ávila-Castillo *et al.*, 2019; Martemucci and D'Alessandro, 2010). Yet, synchronization of estrus by protocols merely including progestogens or prostaglandin F₂α (PGF₂α) during the breeding season may not lead to the desired outcomes (Martemucci and D'Alessandro, 2010) since prolonged exposure with progestogens (≥ 9 days) reduces the quality of oocytes, and in turn, conception rate, and impairs embryonic development (Fierro *et al.*, 2016). It also affects sperm transport and survival in the female reproductive tract (Martemucci and D'Alessandro, 2010). Moreover, devices containing progestogens could escalate the risk of

environmental and tissue contamination (Ávila-Castillo *et al.*, 2019). To overcome these issues, some studies have shown short-term exposure to progesterone; short-term synchronization protocols have shown acceptable results in sheep during breeding as well as non-breeding seasons because they not only enhance estrus synchronization but also culminate in oocytes of higher quality (Titi *et al.*, 2010).

Administration of a single dose of PGF₂α during the breeding season does not influence synchrony of estrus and ovulation, because it can induce luteolysis but not synchrony in follicle development. (Martemucci and D'Alessandro, 2011). Nevertheless, two injections of PGF₂α at 7-16-day intervals have been reported to improve synchrony of estrous and ovulation (Fierro *et al.*, 2016; Carvalho *et al.*, 2018), but the interval between two administrations of PGF₂α could impact fertility (Ataman and Akoz, 2006). In this sense, administration of GnRH was incorporated at the beginning of the protocol to control follicular growth prior to PGF₂α injection (Ataman and Akoz, 2006; Titi *et al.*, 2010; Vallejo *et al.*, 2019). As a result, short-term GnRH-PGF₂α-based protocol (5-day interval) has been recommended as a practical alternative to common progestogen-based estrous synchronization protocols during the breeding season (Ataman and Akoz, 2006; Titi *et al.*, 2010; Martemucci and D'Alessandro, 2010; Hashem *et al.*, 2015).

Studies on ewes (Bartlewski *et al.*, 2017; Titi *et al.*, 2010) and cows (Simões *et al.*, 2018) have shown that the use of exogenous progesterone between GnRH and PGF₂α injections prevents the spontaneous luteolysis resulted from irresponsiveness of the dominant follicle to GnRH injection. Response to GnRH depends on the stage of the estrous cycle in which the injection is made. Furthermore, exogenous progesterone also increases pregnancy rates and boosts follicular growth (Martinez and Gonzalez, 2019).

In cows, the highest efficiency of Ovsynch program is achieved when there is a dominant responsive follicle at the time of GnRH injection and a suitable corpus luteum at the time of PGF 2α injection (Youngquist and Threlfall, 2007). Studies have shown that starting Ovsynch at the beginning and at the end of the estrous cycle has not good results due to the presence of new follicles that are not responsive to GnRH, and premature luteolysis and onset of estrus before the second GnRH administration, respectively (Youngquist and Threlfall, 2007). Therefore, pre-synchronization using two PGF 2α injections 14 days apart was used 12 days before the start of the Ovsynch program in cows, until the cows were in ideal phase of estrus cycle at the beginning of the Ovsynch program (days 5-12) (Zarkouny *et al.*, 2004).

The use of pre-synchronization to increase the efficiency of GnRH-PGF 2α based programs in sheep is a new practice and only has been carried out sparingly in one study, with two PGF 2α injections nine days apart (Ávila-Castillo *et al.*, 2019).

To increase the efficiency of short-term protocols based on GnRH-PGF 2α , Ferdowsi *et al.* (2018) evaluated eCG performance at different time intervals to PGF 2α . The results of the study showed that the concomitant use of eCG with PGF 2α can increase the effectiveness of short-term GnRH-PGF 2α -based programs.

This study aimed to increase the efficiency short-term protocols to of GnRH-PGF 2α - eCG achieve the appropriate program for increasing the fertility rate resulting from laparoscopic insemination. Therefore, the effects of pre-synchronization with two PGF 2α injections nine days apart, three days before the start of GnRH-PGF 2α - eCG-based short-term

programs, and the effect of using 9-day exogenous progesterone were evaluated in this study.

Materials and Methods

Location of Study

The study was carried out in an ewe flock located in Alborz province of Iran in Eshtehard District with a longitude of 50 degrees and 22 minutes and latitude of 35 degrees and 43 minutes during the breeding season.

Animals

In this study, 120 healthy ewes of Zandi breed were used (age range: 2-6 years). Average weight and body condition score (BCS) were 57.3 ± 1.4 kg and 3.1 ± 0.1 , respectively. All ewes received the same diet and had ad libitum access to fresh water.

Experimental Design

Ewes were randomly assigned into four experimental groups (n=30 in each group) considering age, weight, and BCS. All ewes received GnRH (25 μ g of alarelin acetate, Vetaroline, Aboureihan, Iran), and five days afterwards, PGF 2α (75 μ g d-cloprostenol, Vetaglandin, Aboureihan, Iran) plus eCG (400 IU; Gonaser, Hipra, Spain) (Ferdowsi *et al.*, 2018). In controls (Group 1), ewes received no additional treatment. In controlled internal drug releasing (CIDR) group (Group 2), ewes received 5-day CIDR between GnRH and PGF 2α of main estrus synchronization protocol. In Pre-synch group (Group 3), ewes received two injections of PGF 2α at 9-day interval three days before GnRH administration of main estrus synchronization protocol. In Pre-synch-CIDR group (Group 4), ewes received both injections of PGF 2α at 9-day interval and 5-day CIDR ([Figure 1](#)).




Figure 1. Experimental design including blood sampling (BS), injection and timing in different experimental groups (30 ewes in each group)

Estrus Detection

Estrus detection using teaser rams equipped with aprons and crayons was started six hours after PGF₂α injection and lasted for 48-hour after that (Ferdowsi *et al.*, 2018).

Laparoscopic AI

All the ewes were inseminated based on fixed time laparoscopic AI at 48 hours after administration of the last PGF₂α (Youngquist and Threlfall, 2007). To this end, the ewes were deprived of food and water for 12 h prior to laparoscopy (Sathe, 2018). Animals were sedated with 0.3 ml acepromazine maleate (Neurotanq; 10 mg/mL, Alfasan, Holland) 10–30 minutes before insemination (Sathe, 2018). Ewes were placed in laparoscopic cradles and kept in dorsal recumbency. Local anesthesia was performed by administration of 2 mL lidocaine hydrochloride (Luracaine; 20 mg/mL; Vétaquinol S.A., France) in the site of abdominal puncture 5 min prior to insemination (Sathe, 2018). Two trocars were inserted about 3 cm on either side of the midline and about 3 cm cranial to udder to allow for introducing telescope and insemination pipette (Sathe, 2018). Following observation of uterine horns, half of the volume of thawed frozen semen (100×10^6 sperm; Animal Breeding Center, Karaj, Iran) was inseminated in each uterine horn. After insemination, all trocar puncture wounds were sprayed with a combined antiseptic and insect

repellant (Vetaque, Animal Health Division, Iran). All laparoscopic operations were performed by an experienced operator.

Evaluation of Blood Serum Progesterone Concentration

Blood serum progesterone concentrations were measured in all ewes prior to injection of PGF₂α (day 5). Blood samples (10 mL) were collected from jugular vein, poured into vacuum tubes without anticoagulants (Sunphoria, Mediplus, China; Figure 1), and centrifuged at 3000 rpm for 15 min. Serum was isolated and transferred to 2-mL tubes and maintained at -20°C until measurement of progesterone (Ferdowsi *et al.*, 2018). Progesterone concentration was measured by ELISA kit (DRG Instruments GmbH, Germany) with detection limit of 0.045 ng to 40 ng progesterone per mL. Intra-assay and inter-assay coefficients of variation were calculated as 5.40% and 9.96%, respectively.

Pregnancy Diagnosis

Diagnosis of pregnancy was performed by transabdominal ultrasonographic examination using a B-mode ultrasound machine (V9, Emperor, China) equipped with a 3.5 MHz convex probe, 50 days after laparoscopic AI (Youngquist and Threlfall, 2007).

Evaluation of Reproductive Parameters

The following reproductive parameters were assessed: time of estrus onset, estrus detection rate (number of estrous ewes divided by total

number of ewes × 100), pregnancy rate (number of pregnant ewes divided by number of inseminated ewes × 100), lambing rate (number of lambed ewes divided by number of inseminated ewes × 100), fecundity rate (number of lambs divided by number of inseminated ewes), and prolificacy rate (number of lambs divided by number of lambed ewes × 100). Weight and gender of lambs were also recorded (Vodjgani et al., 2017).

Statistical Analysis

Continuous data (i.e., progesterone concentration, time of estrus onset, and birth weight) were analyzed by General Linear Model (GLM) procedure. Binary data (i.e., estrus detection rate, pregnancy rate, lambing rate, fecundity, prolificacy, and sex ratio) were analyzed using GENMOD procedure including function link logit in the model. Multiple comparisons were performed using LSMEANS statement. All analyses were conducted in SAS version 9.4 (Statistical Analysis Systems, Cary, NC, USA). Data were presented as mean ± SEM or percentage. Differences were considered significant at P-value<0.05.

Results

Pre-synchronization using two administrations of PGF2α nine days apart had no influence on the concentration of progesterone at the time of PGF2α administration in the main synchronization protocol (P>0.05) (Table 1).

However, application of CIDR for five days between GnRH and PGF2α injections in the main protocol culminated in higher concentration of circulating progesterone at the time of PGF2α in the main synchronization protocol (P<0.0001) (Table 1), and in turn, progesterone concentration at this time point was greater in CIDR and Pre-synch-CIDR groups as compared with the control and Pre-synch groups (P<0.0001) (Table 1).

Estrus detection rate was not affected by pre-synchronization using two administrations of PGF2α nine days apart, five-day usage of CIDR, and their interaction (P>0.05) (Table 1).

Time to commencement of behavioral estrus expression was advanced by 3.65 hours by the main effect of pre-synchronization using two administrations of PGF2α nine days apart (30.44±1.19 hours in non-pre-synchronized ewes versus 26.79±1.02 hours in pre-synchronized ewes; P=0.022; Table 1). Yet, five-day CIDR and interaction of pre-synchronization using two PGF2α injections nine days apart and five-day CIDR did not impact the estrus onset time (P>0.05) (Table 1).

Pregnancy rate, lambing rate, fecundity, prolificacy, birth weight of lambs, and sex ratio of offspring were not influenced by two administrations of PGF2α nine days apart, five-day usage of CIDR, and their interaction (P>0.05) (Table 1).

Table 1. Reproductive indices of ewes prepared for timed breeding using GnRH- prostaglandin F2α5 days apart, in association with pre-synchronization (two prostaglandin F2 ,α9 days apart) and CIDR during breeding season

Parameter	Experimental group				P-value		
	Control	CIDR	Presynch	Presynch -CIDR	Pre-synch	CIDR	Presynch-CIDR
Progesterone (ng/ml)	2.6 ± 0.33 ^a	4.6 ± 0.23 ^b	2.4 ± 0.22 ^a	4.7 ± 0.15 ^b	0.38	< 0.0001	0.62
Estrus detection rate (%)	93.3 (28/30)	93.3 (28/30)	96.7 (29/30)	96.7 (29/30)	0.39	1.00	1.00
Time of estrus onset (hour)	31.1 ± 1.54	29.8 ± 1.82	27.1 ± 1.48	26.5 ± 1.48	0.02	0.53	0.81

		Experimental group			P-value		
Pregnancy rate (%)	50 (15/30)	46.7 (14/30)	56.7 (17/30)	60.0 (18/30)	0.27	0.99	0.71
Lambing rate (%)	50 (15/30)	46.7 (14/30)	56.7 (17/30)	60.0 (18/30)	0.27	0.99	0.71
Fecundity	60 ± 12	47 ± 9	70 ± 12	70 ± 11	0.24	0.55	0.55
Prolificacy	120 ± 11	100 ± 0	120 ± 10	110 ± 8	0.60	0.12	0.42
Birth weight (kg)	4.2 ± 0.19	4.7 ± 0.18	4.4 ± 0.26	4.4 ± 0.22	0.91	0.36	0.21
Sex ratio (%)	50 (9/18)	42.9 (6/14)	55.0 (11/20)	30.0 (6/20)	0.71	0.17	0.44

^{a,b}Various letters indicate significant difference ($P < 0.05$).

Discussion

The present study aimed to evaluate whether application of pre-synchronization using two PGF_{2α} injections nine days apart and usage of 5-day CIDR before and during fixed time AI protocol, respectively, could enhance estrus synchronization and fertility of ewes. The efficient estrus and ovulation synchronization is considered as one of the cornerstones of fixed time AI programs in sheep industry (Noakes *et al.*, 2019; Sathe, 2018; Youngquist and Threlfall, 2007).

Pre-synchronization protocol with two injections of PGF_{2α} with 9-day interval and three days before GnRH administration of main estrus synchronization protocol (Group 3) increased fertility rate by 56.7 %. No significant difference was found between the third group and the controls. While fertility and fecundity rates (60%) improved in the pre-synchronization technique, the progesterone was used in the original protocol for five days; but no statistically significant difference was found between the groups, which might be due to small sample size. These results were consistent with the findings of Ávila *et al.* (2019) who assessed pre-synchronization utilizing two injections of PGF_{2α} within 9-day intervals. Although no significant difference was found between the pregnancy and normal mating

rates in the studied groups, the highest rates belonged to the pre-synchronization protocol with 5-day progesterone administration (90%) (Ávila-Castillo *et al.*, 2019). An explanation for this improvement in fertility could be the fact that pre-synchronization prepares the females to initiate the main protocol at a more appropriate stage of estrous cycle so that the GnRH could act as an agent to eliminate the current follicular wave to pave the way for emergence of the subsequent follicular wave at the favorable time (Navanukraw *et al.*, 2004).

Martemucci and D'Alessandro (2011) assessed the efficacy of short-term 5-day GnRH-FGA-PGF_{2α}-eCG-based synchronization protocol for natural mating and laparoscopic AI. Fertility rates in laparoscopic AI in different treatment groups were reported as 60% and 40%, respectively, within 60 and 52 hours. These results were consistent with the findings of the present study regarding fertility rates (46-60%).

Pre-synchronization with two injections of PGF_{2α} within 9-day interval prior to five-day GnRH-PGF_{2α}-eCG-based protocol advanced estrus manifestation, although it had no effect on progesterone concentration at the last PGF_{2α} injection, estrus rate, and other reproductive parameters in ewes. It has been reported that pre-synchronization with two injections of PGF_{2α}

prior to Ovsynch in cattle can enhance the fertility rate in artificial insemination by synchronizing estrus and ovulation (Nowicki *et al.*, 2017). Moreover, El-Zarkouny *et al.* (2004) assessed the effect of pre-synchronization technique with two injections of PGF_{2α} within 14-day interval prior to Ovsynch protocol; they also assessed the efficacy of CIDR. The results showed that a single use of pre-synchronization and CIDR increased the fertility rate, but combination of CIDR and pre-synchronization had no effect on fertility rate (El-Zarkouny *et al.*, 2004).

Use of progesterone caused a sudden increase in progesterone level followed by decreased secretion of pituitary LH (Noakes *et al.*, 2019), which resulted in follicular atresia and generation of new waves of follicles (Martinez and Gonzalez, 2019). A 5-day progesterone protocol generated regular waves of follicles. Adequate progesterone level during follicular development helps to create healthy oocytes (Martinez and Gonzalez, 2019; Nowicki *et al.*, 2017). However, the 5-day progesterone in the synchronization protocol significantly increased the progesterone level at the time of PGF_{2α} injection in the original protocol (on day 5); but no significant difference was found in reproductive parameters. This finding was consistent with the findings obtained in the study on cyclic cows (Nowicki *et al.*, 2017). This might be due to the fact that in the main estrus synchronization protocol, GnRH injection induced ovulation and luteinization or luteinization without ovulation in some ewes (Youngquist and Threlfall, 2007), which would lead to increased endogenous progesterone. Therefore, it is possible that elevation of endogenous progesterone by GnRH overrode the potential beneficial effects of exogenous progesterone provided by CIDR. However, Titi *et al.* (2010) used the 5-day progesterone in the GnRH-PGF_{2α} protocol. The results showed that exogenous progesterone

improved estrus and fertility. The difference between the present study and the former study was simultaneous injection of eCG and PGF_{2α}, which did not necessitate progesterone protocol. Furthermore, Martinez *et al.* (2019) showed that 5-day progesterone administration without eCG (the same as 14-day conventional protocol) was more effective than 6- or 7-day progesterone administration in ovulation and fertility. This might be due to ovulation of a dominant follicle induced by progesterone on day 5 without eCG administration. The eCG should be added to the protocol for accurate synchronization of ovulation and estrus. Martinez *et al.* (2019) used the short-term progesterone administration in GnRH-PGF_{2α}-based protocol without eCG in Group 3 and reported the fertility rate in normal mating as 68.4%, which did not significantly differ from the other studied groups.

Conclusion

Given the findings of the present study and the certain length of estrus in ewes (24 to 36 hours, Abecia *et al.*, 2012), estrus might have long passed in some ewes due to early onset of estrus cycle following the last PGF_{2α} injection at the time of AI. Therefore, AI was delayed in pre-synchronized ewes with early estrus cycle. A certain AI time should be determined in pre-synchronized ewes in future studies.

Considering the fertility rate in the different groups in this study (46%-60%), short-term 5-day GnRH-PGF_{2α}-eCG-based protocols can be used during the breeding season to synchronize estrus and ovulation (50% fertility rate in the control group). However, estrus rate and reproductive parameters (e.g., pregnancy rate, lambing rate, fecundity, and prolificacy) did not show significant differences between different groups in this study. Yet, pre-synchronization improved the onset of estrus expression.

Acknowledgments

The authors appreciate the cooperation of Seyed Javad Hosseini and personnel of the

Doab Agricultural and Industrial Company. We are also grateful for the help of Hamidreza Ferdowsi and other PhD students in the Department of theriogenology, Faculty of Veterinary medicine, University of Tehran.

Conflict of Interest

The authors declared no conflict of interest.

References

- Abecia, J.A., Forcada, F., Gonzalez-Bulnes, A. (2012). Hormonal control of reproduction in small ruminants. *Anim Reprod Sci*, 130,173-179. [DOI:10.1016/j.anireprosci.2012.01.011] [PMID]
- Ataman, M.B., Akoz, M. (2006). GnRH-PGF₂α and PGF₂α-PGF₂α synchronization in akkaraman cross-bred sheep in the breeding season. *B Vet I Pulawy*, 50, 101-104.
- Ávila-Castillo, B. R., García-Flores, E. O., Molina-Mendoza, P., Peralta-Ortiz, J., & Sánchez-Torres-Esqueda, M. T. (2019). Sincronización del estro en ovejas de pelo mediante protocolo basado en prostaglandinas+ GnRH. *CienciaUAT*, 13(2), 141-151. [DOI:10.29059/cienciauat.v13i2.1012]
- Bartlewski, P. M., Sohal, J., Paravinja, V., Baby, T., Oliveira, M. E. F., Murawski, M., ..., and Keisler, D. H. (2017). Is progesterone the key regulatory factor behind ovulation rate in sheep? *Domest Anim Endocrinol*. 58, 30-38. [DOI:10.1016/j.domaniend.2016.06.006] [PMID]
- Casali, R., Pinczak, A., Cuadro, F., Guillen-Muñoz, J.M., Mezzalira, A. and Menchaca, A. (2017). Semen deposition by cervical, transcervical and intrauterine route for fixed-time artificial insemination (FTAI). in the ewe. *Theriogenology*, 103, 30-35. [DOI:10.1016/j.theriogenology.2017.07.021] [PMID]
- Carvalho, S.F., Souza-Fabjan, J.M.G., Balaro, M.F.A., Bragança, G.M., Pinto, P.H.N., Almeida, J.G., Moura, A.B.B., Fonseca, J.F. and Brandão, F.Z. (2018). Use of two doses of cloprostenol in different intervals for estrus synchronization in hair sheep under tropical conditions. *Trop Anim Health Prod*, 50(2): 427-432. [DOI:10.1007/s11250-017-1454-x] [PMID]
- El-Zarkouny, S, Z., Cartmill, J.A., Hensley, B.A., Stevenson, J.S., (2004). Pregnancy in dairy cows after synchronized ovulation regimens with or without presynchronization and progesterone. *J Dairy Sci*, 87,1024–1037. [DOI:10.3168/jds.S0022-0302(04)73248-8]
- Fierro, S., Vinales, C., Olivera-Muzante, J. (2016). Concentrations of steroid hormones, estrous, ovarian and reproductive responses in sheep estrous synchronized with different prostaglandin-based protocols. *Anim Reprod Sci*, 167,74–82. [DOI:10.1016/j.anireprosci.2016.02.009] [PMID]
- Ferdowsi, H.R., Vodjgani, M., Gharagozloo, F., Talebkhan Garoussi M., Niasari Naslaji, A., Akbarinejad, V. (2018). The effects of eCG injection time on the reproductive performance in Shal ewes treated with short-term synchronization program during the breeding season. *J Vet Res*, (in press).
- Hashem, N.M., El-Zarkouny, S.Z., Taha, T.A., Abo-Elezz Z.R. (2015). Oestrous response and characterization of the ovulatory wave following oestrous synchronization using PGF₂α alone or combined with GnRH in ewes. *Small Rumin Res*, 129, 84-87. [DOI:10.1016/j.small-rumres.2015.06.003]
- Martemucci, G., D'Alessandro, A.G. (2010). Estrous and fertility responses of dairy ewes synchronized with combined short-term GnRH, PGF₂α and estradiol benzoate treatments. *Small Rumin Res*, 93, 41–47. [DOI:10.1016/j.small-rumres.2010.05.001]
- Martemucci, G., D'Alessandro, A.G. (2011). Synchronization of oestrus and ovulation by short time combined FGA, PGF₂α, GnRH, eCG

- treatments for natural service or AI fixed-time. *Anim Reprod Sci*, 123, 32–39. [DOI:10.1016/j.anireprosci.2010.11.007] [PMID]
- Martinez P. R, Gonzalez A.B. (2019). Efficiency of CIDR-Based Protocols Including GnRH Instead of eCG for Estrus Synchronization in Sheep. *J Animals*, 9, 146. [DOI:10.3390/ani9040146] [PMID] [PMCID]
- Navanukraw, C., D. A. Redmer, L. Reynolds, J. D. Kirsch, A. T. Grazul-Bilska, and P. M. Fricke. (2004). A modified presynchronization protocol improves fertility to timed artificial insemination in lactating dairy cows. *J Dairy Sci*. 87,1551–1557. [DOI:10.3168/jds.S0022-0302(04)73307-X]
- Noakes, D.E., Parkinson, T.J., England, G.C.W. (2019). Pharmacological Agents in the Control of Reproduction. In: *Veterinary reproduction and obstetrics*. (10th ed). Saunders Elsevier Philadelphia, USA.
- Nowicki. A, Baranski. W, Baryczka. A, Janowski, T. (2017). OvSynch protocol and its modifications in the reproduction management of dairy cattle herds an update. *J Vet Res*, 61, 329-336. [DOI:10.1515/jvetres-2017-0043] [PMID] [PMCID]
- Olivera-Muzante, J., Fierro, S., Alabart, J. L., Claramunt, M., Minteguiaga, M. A., Aunchayna, G., ... & Banchemo, G. (2019). Short-term dietary protein supplementation improves reproductive performance of estrous-synchronized ewes when there are long intervals of prostaglandin or progesterone-based treatments for timed ai. *Anim Reprod Sci*, 206, 78-84. [DOI:10.1016/j.anireprosci.2019.05.011] [PMID]
- Risco, C., & Melendez, P. (Eds.). (2011). *Dairy Production Medicine*. John Wiley & Sons. [DOI:10.1002/9780470960554]
- Sathe, S. R. (2018). Laparoscopic Artificial Insemination Technique in Small Ruminants—A Procedure Review. *Frontiers in veterinary science*, 5, 266. [DOI:10.3389/fvets.2018.00266] [PMID] [PMCID]
- Simões, L. M. S., Orlandi, R. E., Massoneto, J. P. M., Scandiuizzi Jr, L. A., Freitas, B. G., Bastos, M. R., ... & Sales, J. N. S. (2018). Exposure to progesterone previous to the protocol of ovulation synchronization increases the follicular diameter and the fertility of suckled *Bos indicus* cows. *Theriogenology*, 116, 28-33. [DOI:10.1016/j.theriogenology.2018.04.031] [PMID]
- Titi HH, Kridli RT and Alnimer MA. (2010). Estrus Synchronization in Sheep and Goats Using Combinations of GnRH, Progestagen and Prostaglandin F2a. *Reprod Domest Anim* 45, 594–599.
- Vallejo, D.A., Londoño, J.D., Yepes, Y.A., V. Tamayo, V., Mejia, A.F., Maldonado, J.G. (2019). Pregnancy rates in hair sheep after Ovsynch synchronization and a combined intracervical fixed-time artificial insemination and 10-day mating period. *Vet World*, 12(11), 1779-1783. [DOI:10.14202/vetworld.2019.1779-1783] [PMID] [PMCID]
- Vodjgani, M., Gharagozloo, F., Mahmoodzadeh, H., Assadi Moghaddam, B., Ferdowsi, H.R., Hosseini, A. (2017). *Sheep and goat obstetrics and reproductive diseases*. (2nd ed). Tehran University Press, Tehran, Iran.
- Youngquist, R. S., & Threlfall, W. R. (2006). *Current Therapy in Large Animal Theriogenology-E-Book*. Elsevier Health Sciences.

تأثیر پیش همزمان سازی و CIDR روی نتایج حاصل از برنامه های کوتاه مدت همزمان سازی در میش های نژاد زندی طی فصل تولید مثل

رضا ارجمندی*، مهدی وجگانی، امیر نیاسری نسلجی، فرامرز قراگوزلو، وحید اکبری نژاد

گروه مامایی و بیماری های تولیدمثل، دانشکده دامپزشکی، دانشگاه تهران، تهران، ایران

(دریافت مقاله: ۱۴ مرداد ماه ۱۳۹۹، پذیرش نهایی: ۷ آبان ماه ۱۳۹۹)

زمینه مطالعه: برنامه های همزمان سازی تولید مثل برای اجرای گسترده برنامه های تلقیح مصنوعی در گوسفند ضروری است.

هدف فرضیه ما در مطالعه حاضر این بود که، آیا کاربرد پیش همزمان سازی و CIDR، به ترتیب قبل و در طول برنامه اصلی تلقیح در زمان ثابت، می تواند همزمان سازی فحلی و باروری را در میش افزایش دهد.

روش کار: میش ها به طور تصادفی به چهار گروه آزمایشی (۳۰ راس در هر گروه)، با توجه به سن، وزن و نمره وضعیت بدنی تقسیم شدند. همه میش ها GnRH (۲۵ میکروگرم آلارلین استات) و پنج روز پس از آن، PGF2α (۷۵ میکروگرم دی-کلوپرستول) به همراه eCG (۴۰۰ واحد بین المللی) دریافت کردند. در گروه کنترل، میش ها درمان اضافی دریافت نکردند. در گروه CIDR، میش ها ۵ روز CIDR بین GnRH و PGF2α برنامه اصلی دریافت کردند. در گروه Presynch، میش ها دو تزریق PGF2α به فاصله ۹ روز، ۳ روز قبل از تجویز GnRH برنامه اصلی دریافت کردند. در گروه CIDR، میش ها هم دو تزریق PGF2α به فاصله ۹ روز و هم CIDR ۵ روزه دریافت کردند. غلظت پروژسترون سرم خون قبل از تزریق PGF2α (روز ۵) در کلیه میش ها اندازه گیری شد. ۴۸ ساعت بعد از تجویز آخرین PGF2α تلقیح مصنوعی برای همه میش ها به روش لاپاروسکوپی انجام شد.

نتایج: هیچ اثر متقابل بین CIDR و برنامه های پیش همزمان سازی یافت نشد ($P > 0.05$). غلظت پروژسترون در گروه CIDR نسبت به گروه های بدون CIDR بیشتر بود ($P < 0.001$). چرخه فحلی تحت تأثیر پیش همزمان سازی و CIDR قرار نگرفت ($P > 0.05$). بعد از آخرین تزریق پروستاگلاندین، فحلی در میش های با پیش همزمان سازی در مقایسه با میش های بدون پیش همزمانی سریع تر شروع شد ($P = 0.022$). سایر شاخص های مورد ارزیابی تفاوت معنی داری بین گروه های درمانی نشان نداد ($P > 0.05$).

نتیجه گیری نهایی: میزان فحلی و پارامترهای تولید مثل اختلاف معنی داری بین گروه های مختلف در این مطالعه نشان نداد. با این حال پیش همزمان سازی زمان شروع فحلی را تسریع کرد.

واژه های کلیدی: فصل تولید مثلی، سیدر، میش، پیش همزمان سازی، روش های کوتاه مدت همزمان سازی.