Published online 2016 May 31.

Research Article

Antioxidant Expression Response to Free Radicals in Active Men and Women Fallowing to a Session Incremental Exercise; Numerical Relationship Between Antioxidants and Free Radicals

Behrouz Baghaiee,^{1,*} Mohammad Reza Aliparasti,² Shohreh Almasi,² Marefat Siahkuhian,¹ and Behzad Baradaran³

¹Department of Physical Education and Sport Science, Mohaghegh Ardabili University, Ardabil, IR Iran ²Department of Immunology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, IR Iran ³Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, IR Iran

Corresponding author: Behrouz Baghaiee, Department of Physical Education and Sport Science, Mohaghegh Ardabili University, Ardabil, IR Iran. Tel: +98-4533512085, Fax: +98-4533512081, E-mail: behrouz_phsport@yahoo.com

Received 2015 May 31; Revised 2015 December 18; Accepted 2015 December 28.

Abstract

Background: Energy production is a necessary process to continue physical activities, and exercise is associated with more oxygen consumption and increase of oxidative stress. what seems important is the numerical relationship between antioxidant and free radicals. Although the activity of some enzymes increases with physical activities, but it is possible that gene expression of this enzyme is not changed during exercise.

Objectives: The aim of the present study is to investigate the antioxidant enzymes gene expression and changes in malondialdehyde (MDA) and total antioxidant capacity (TAC) levels in men and women affected by a session of incremental exercise and to carefully and numerically assess the relationship between MDA changes and gene expression and activity of antioxidant enzymes.

Materials and Methods: 12 active men and 12 active women (21-24 years old) participated voluntarily in this study. Peripheral blood samples were taken from the subjects in three phases, before and after graduated exercise test (GXT) and 3 hours later (recovery). **Results:** The gene expression of manganese superoxide dismutase (MnSOD) enzyme increased significantly in women in the recovery phase (P < 0.05). Catalase gene expression significantly increased in men in both phases (immediately & recovery) (P < 0.05). But the changes in active women were only significant immediately after the exercise. TAC levels increased significantly in men in the recovery phase and in active women immediately after the exercise (P < 0.05). MDA activity also increased significantly in men in both phases (P < 0.05). However, in women the increase was significant only in the recovery phase (P < 0.05). There was a reverse relationship between changes in MnSOD and copper- and zinc-containing superoxide dismutase (Cu/ZnSOD) levels and MDA in men (P < 0.05). In active women there was also a significant relationship between changes in MDA and gene expression of Cu/ZnSOD and TAC (P < 0.05).

Conclusions: The increase in free radicals during incremental exercises challenges gene expression and activity of antioxidant enzymes. However, despite the negative effects of free radicals, in women, activity and gene expression of antioxidant enzymes respond appropriately to free radicals.

Keywords: Antioxidant Enzyme, Free Radicals, Exercise

1. Background

Energy production is a necessary process to continue physical activities, but energy production leads to the generation of oxidative stress (1-3). Thus, exercise is associated with more oxygen consumption and increase of the oxidative stress (4-6). Oxidative stress includes definitions that have undergone some changes with the development of science. However oxidative stress reflects the imbalance between antioxidant activity and free radicals (7-9). Studies have indicated that the activity of antioxidants is influenced by the amount of free radicals that are produced during physical activity, also gender has effect on this antioxidant response (10-13). It has been reported that estrogen in women leads to increase of antioxidant enzymes activity in the immune system and other cells through estrogen receptors (14, 15). The estrogen receptors are divided into ERa and ERb, that ERb receptors are more predominant in lymphocytes and by which the activity and gene expression of antioxidant enzymes are more affected (16-18).

But what seems important is the numerical relationship between antioxidant and free radicals. Although the activity of some enzymes increases with physical activities, but it is possible that gene expression of the this enzyme is not changed during exercise (11, 19). Some studies suggest the increased activity of antioxidant enzymes during in-

Copyright © 2016, Sports Medicine Research Center. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

tense physical activities (20), but the activity of an enzyme does not necessarily mean the associated gene expression. Also, it is not clear that to what extent changes in the level of free radicals impact MnSOD, Cu/ZnSOD and catalase enzymes expression and also total antioxidant capacity (TAC) and how gender can affect this process. Therefore, it seems that physical fitness levels and gender can produce different responses in terms of gene expression and activity of antioxidant enzymes (20-22). On the other hand, the quantitative (numerical) relationship between gene expression of antioxidant enzymes and their activity associated with increased level of free radicals is not defined. As previous studies have not reported whether the relationship is direct (positive) or indirect (negative), so it is not clearly defined that for every one unit increase in the level of MDA, how gene expression and activity of antioxidant enzymes changes and how gender affects the relationship.

2. Objectives

Thus, the aim of the present study was to examine the gene expression of mitochondrial and cytosolic antioxidant enzymes and changes in MDA and TAC level in active men and women affected by an incremental exercise session and to carefully and numerically investigate the relationship between changes in MDA and gene expression and activities of antioxidant enzymes.

3. Materials and Methods

3.1. Subjects

Twenty four active and healthy men and women (aged 21 - 24 years old, BMI: 22 - 25 Kg/m²) volunteered in this study. All volunteers attended the exercise venue and received information about the protocols and procedures, as well as the possible risks and benefits involved in the study. Informed written consent was obtained from all the participants. The ethics committee of Tabriz University of Medical Sciences, Iran, approved the study design. To determine their inclusion in the study, volunteers completed medical history and physical activity questionnaires (PAS) (23), and took part in physical examinations. The volunteers also were advised to avoid the use of caffeine, nicotine, alcohol, antihypertensive medications, and an exhaustive workout (which could influence the immune response) in the previous hours. The subjects were then randomly assigned to the men (n = 12) and women (n = 12) groups.

3.2. Measurements

Each volunteer was asked to empty their bladder before laboratory examinations and tests. In basal state we measured age, height, weight, heart rate (HR), fat %, body mass index (BMI), maximal oxygen uptake (VO_{2max}) and lipid profile. Gene expression of MnSOD, Cu/ZnSOD, catalase enzyme and TAC, MDA levels measured in three stages; pre exercise test, immediately and 3 hours after exercise (recovery). Body weight (kg) and height (cm) were measured using the standard digital floor scale and precision stadiometer (Seca, Germany.) Body fat percent and BMI (kg/m²) were measured using the portable bioelectrical impedance system (body composition, Omron, China). Maximal oxygen uptake (VO_{2max}, mL.kg⁻¹.min⁻¹) was measured by performing a test on an electronically braked bicycle ergometer (Monark 839 E, Sweden), with initial resistance set at 60 W and 20 W step-wise increments every 3 minutes. Heart rate and rhythm were continuously recorded by 12-lead electrocardiogram and subjects reported their rate of perceived exertion (RPE) on the 15-point Borg scale at the end of each stage (Noble et al. 1983). The volume of oxygen consumed (VO_2) and carbon dioxide produced (VCO₂) during exercise were calculated from minute ventilation (V), measured using mass flow ventilometry, and simultaneous mixing chamber analysis of expired gas fractions (V max E, Sensor Medics, Yorba Linda, CA). Gas analyzers and flow probes were calibrated before each test. VO_2 and VCO_2 were recorded during the final 40 seconds of each stage of the test and at peak exercise and expressed in L min⁻¹ and also relative to body weight (mL.kg⁻¹.min⁻¹) (Andrew et al. 2001). The fix amount of food for volunteers was determined and all volunteers received a detailed verbal explanation and written instructions. Any questions regarding the type and amount of food and beverages consumed were resolved individually (24).

3.3. Exercise Protocol

Subjects started walking on the treadmill (EXE T600, European exercise equipment, EC, TUV) for 3 minutes (Grad: 0%, Speed: 3 Km.h⁻¹) According to the graduated exercise test (GXT). Then up to 5 minutes, g rad of the treadmill was increased to 5% and then did not change. But treadmill speed was increased to 13 Km.h⁻¹ until the end of the training period, and at the end of the training, volume and intensity was increased to 25 minutes and 80% of maximal exercise heart rate. The exercise training was performed at the ambient temperature, 23 - 24°C and relative humidity of 42%. Subjects were supervised by trained physical education teachers. All exercise sessions began between 09.00 and 11.00 hours to avoid circadian rhythm effects.

3.4. RNA

For RNA extraction and biochemical assays, 4 mL blood samples, following a 12-hour overnight fast, were taken between 08.00 and 09.00 hours. Samples were taken at baseline, immediately and 3 hours after exercise. All samples were taken 24 hours after the last exercise bout. Total RNA extraction from peripheral blood (lymphocyte cell) was performed using RNX-PLUS reagent (Cinagen, Iran) following the manufacturer's protocol. Clean-up procedure was carried out to avoid any genomic DNA contamination during qRT-C with a specific wipeout buffer (Quanti Tect Reverse Transcription, Qiagen, Germantown, USA). The RNA quantity and quality were assessed by spectrophotometry. The reverse transcription reaction was performed using revert aid first standard cDNA kit (Fermentas). For cDNA synthesis, RNA was treated with DNaseI (Invitrogen) and a 20 Rl reaction solution containing: 4Rl 5X reaction buffer, 1 Rl RNase inhibitor (20u/Rl), 2 Rl dNTP mix (10mM), 1 Rl random hexamer primer, 1 Rl oligo (dt) 18 primer, and placed 5 minutes at 65°C and 60 minutes at 42°C. Reverse transcription activity was terminated by heating at 70°C for 5 minutes. For RNA extraction 1 mL of RNX solution was mixed with 5 \times 10⁶ cells and incubated at room temperature for 5 minutes. After adding 200 Rl chloroform it was centrifuged at 12000 RPM at 40 3 C for 15 minutes and then transferred to a tube containing an equal volume of isopropanol. After centrifugation at 12000 RPM, 40 C, 15 minutes the supernatant was discarded and mixed with 1 mL of 75% ethanol. Then the pellet was dissolved and centrifuged for 8 minutes at 7500 RPM. Finally the supernatant was discarded and the pellet after drying was dissolved in 20 Rl DEPC-treated water. Approximately 2 Rl of RNA were used to determine RNA yield and quality by optical density reading at 260 nm and 280 nm. Finally, for each sample, real-time PCR reactions were carried out: 20 Rl volume/well containing 10 Rl of 2 \times master mix (10 Rl of 2 \times SYBER Green PCR master mix reagent kit (ABI, USA), 0.3 Rl primer, 2 Rl cDNA and 7.4 Rl DEPC-water mixture) for the tested gene and β -actin RNA. Then the plate placed into the thermocycler (Corbett-gene 6000). The thermal cycle conditions were 95°C for 10 minutes followed by 40 cycles that were run for 15 seconds at 95°C (denaturation) and for 35 seconds at 600 C (Annealing). Expression of MnSOD, Cu/ZnSOD and catalase were standardized with β -actin RNA as endogenous control by melting curve analysis and by 1% agarose gel electrophoresis (Salemi et al. 2011). The primers used for the MnSOD cDNA F-5'-GAGAAGTACCAGGAGGCGTTGamplification were R-5'-CAAGCCAACCCCAACCTGAGC-3'. 3 and for F-5'-AAGGCCGTGTGCGTGCTGAA-3 Cu/ZnSOD and R-5'-CAAGTCTCCAACATGCCTCT-3' and for cata-R-5'-F-5'-TTTGGCTACTTTGAGGTCAC-3 and lase

TCCCCATTTGCATTAACCAG-3'. The primers used for β actin were F-5'-CAGGTCATCACCATTGGCAAT-3' and R-5'-TCTTTGCGGATGTCCACGT-3. To estimate mRNA expression levels, Real-time PCR reaction efficiency was calculated using the standard curve slope. Standard curve drawing was based on serial dilution of the genes of interest against the logarithm concentration of template DNA. MnSOD, Cu/ZnSOD and catalase and β -actin RNA efficiencies were 98% and 97%.

3.5. MDA

Plasma MDA levels were determined by using the methods of Draper and Hadley based on thiobarbituric acid (TBA) reactivity (25).

3.6. TAC

The total antioxidant capacity of plasma (TAC) to scavenge ABTS radicals was measured by a chromogenic method with commercially available kit (Cat. No. NX 2332, Randox, Crumlin, UK) (26).

3.7. Lipid Profile

Total cholesterol, TG, LDL-cholesterol and HDL-cholesterol were determined by enzymatic method (27).

3.8. Statistical Analysis

Kolmogorov-Smirnov test was used to check the normality of data, based on these results the appropriate test was chosen. To compare the values between active and women, we employed the t-test (or Mann-Whitney). The effect of exercise on different parameter intra group was evaluated by repeated measure and Bonferroni test. Also, we used linear mixed model for the relation between groups. All statistical analysis was performed using SPSS 22 (Chicago, IL- USA). P values less than 0.05 were considered as statistically significant.

4. Results

Physiological markers and lipid profile of active men and women have been identified in Table 1. Statistical analysis indicated that gene expression of MnSOD enzyme in active men had increased as a result of incremental physical activities and the increase was also observed in the recovery phase, however, these changes were not significant (P = 0.99). Statistical analysis on gene expression of Cu/ZnSOD enzyme in active men showed that the enzyme mRNA levels have increased immediately after exercise and in the recovery phase, but the changes were not significant (P = 0.346 and P = 0.99, respectively). However, catalase gene expression in active men significantly increased immediately after exercise (P = 0.005). The changes also continued to the recovery phase and catalase mRNA level increased significantly (P=0.003). For total antioxidant capacity (TAC), our studies showed that TAC levels have increased immediately following exercise and also 3 hours later (during recovery), in active men. However, only changes in the recovery phase (3 hours after exercise) were significant (P=0.437 and P=0.009, respectively) (Figure 1).

Furthermore, repeated measures analysis and post hoc Bonferroni test revealed that MDA levels have significantly increased immediately after exercise in men, but in spite of a slight decrease in the recovery phase, the levels were still above the ground state and the difference was significant (P = 0.012 and P = 0.014, respectively) (Figure 1).

Furthermore, mixed model and linear regression analysis showed there was a reverse relationship between changes in MnSOD and Cu/ZnSOD levels and MDA in men (P = 0.001 and P = 0.05, respectively). So that for every one unit increase in the level of MDA, there was a decrease of -3.69 and -2.62 units in gene expression of MnSOD and Cu/ZnSOD, respectively. However, there was not a significant relationship between gene expression of catalase and TAC and MDA in active men (P = 0.456 and P = 939, respectively). As per one unit increase in the level of MDA, catalase gene expression and TAC levels decreased 0.93 and 0.01 units, respectively (Table 2).

The statistical analysis in the group of active women showed that there has not been a significant increase in MnSOD gene expression immediately after exercise (P = 0.421). Yet gene expression of the enzyme increased significantly in the recovery phase (P = 0.006). For the changes in Cu/ZnSOD our studies also showed that gene expression of the enzyme in women has not significantly increased immediately after exercise and during recovery (P = 0.99). Catalase gene expression in active women significantly increased immediately after exercise (P = 0.03), but decreased three hours later and in the recovery phase and was not significantly different from the ground state (P = 0.06). Statistical analysis using the Bonferroni test also showed that TAC level in women has increased significantly immediately after exercise (P = 0.031), but decreased slightly in the recovery phase (P = 0.065) (Table 2).

The MDA concentration was not significantly increased immediately after exercise in women (P = 0.255), but 3 hours after exercise, level of this blood index increased significantly (P = 0.029) (Table 2). Statistical analysis using mixed model test and regression analysis showed that there is not a significant relationship between changes in MDA and gene expression of MnSOD and catalase (P = 0.836 and P = 0.901, respectively), in active women. As per one unit increase in MDA level, the gene expression of Mn-SOD decreased -0.14 units and catalase level increased 0.16 units. However, the relationship between changes in MDA and gene expression of Cu/ZnSOD and TAC was significant (P = 0.014 and P = 0.004, respectively). As per one unit increase in MDA level, the gene expression of Cu/ZnSOD and TAC increased 2.9 and 0.11 units, respectively (Table 2).

Further statistical analysis showed that in all phases of the study (baseline, immediately after exercise and three hours later) active women had higher levels of gene expression of antioxidant enzymes and also TAC. However, in the case of Cu/ZnSOD in the phase immediately after exercise, difference between the two groups was P = 0.017, and for MnSOD difference between the two groups was P = 0.033, P = 0.001 and P = 0.005 at all stages, the difference was reported significant. However, no significant difference was observed in the case of catalase and TAC (Table 3).

5. Discussion

Some research studies emphasize that active men compared to active women have higher activity of the electron transport chain and the xanthine oxidase enzyme that leads to the production of high levels of free radicals (28-30). As our statistical findings show, despite significant increase in MDA level in both active men and women as a result of physical activity, men have higher levels of MDA. In this regard, cardiovascular and respiratory system plays an important role (31, 32). Since athletic men have an effective cardiovascular system and also have highly developed capillary and mitochondrial density, thus the production of O2 and the duration of its presence in tissues is very high (28, 33, 34). So, further increase in free radicals in men will not be unexpected. But research has shown that estrogen prevents the increase of cell damage by increasing the activity of antioxidant enzymes (14, 35).

In this context, statistical review of the present study also showed that women compared to men have higher levels of gene expression of antioxidant enzymes and TAC levels. Other studies in this regard show that lymphocytes and other cells of women have estrogen receptors (36, 37). These receptors can be divided into classes of ERa and ERb, of which ERb receptors are more predominant in the cells of the immune system by which the activity of lymphocytes and antioxidant enzymes are more affected (17). The same studies suggest that the gene expression of antioxidant enzymes, including MnSOD and catalase may be influenced by estrogen (14, 38).

In their findings, the researchers reported that in addition to its effective role in the expression of the estrogen receptor gene, estrogen has a direct relationship with Table 1. Physiological Markers and Lipid Profile in Rest State in Active Men and Women

Variables	Active Men	Active Women	P ^a
Age, y	23 ± 1	21 ± 1	0.168
Height, cm	177.59 ± 7.02	160.8 ± 4.35	0.329
Weight, Kg	70.54 ± 6.1	53.1 ± 5.78	0.947
Vo _{2max} , mL/kg/min	55.97 ± 1.35	52.6 ± 1.83	0.713
Body mass index, kg/m ²	22.5 ± 1.5	25.6 ± 1.83	0.122
Heart rate in basal state, beat/min	78 ± 7.52	67.3 ± 8.67	0.766
TG, mg/dL	86 ± 36.65	75.3 ± 18.97	0.701
Cholesterol, mg/dL	131 ± 14.24	134.3 ± 22.72	0.708
HDL-C, mg/dL	35.11 ± 5.49	46 ± 10.02	0.821
LDL-C, mg/dL	102.10 ± 5.12	96.60 ± 4.26	0.512
^a Based on independent t test ($P \le 0.05$).			

Table 2. Relationship Between MDA Change and Antioxidant Enzyme mRNA Fallowing to Incremental Exercise

Response	Variable	Active Men		Active Women	
		B (95% CI)	Р	B(95% CI)	P ^a
Cu/znSOD	MDA	-2.62 (-5.26 to 0.02)	0.050	2.9 (0.66 to 5.14)	0.014
MnSOD	MDA	-3.69 (-5.57 to -1.8)	0.001	-0.14 (-1.5 to 1.23)	0.836
Catalase	MDA	-0.93 (-3.57 to 1.71)	0.456	0.16 (-2.46 to 2.77)	0.901
TAC	MDA	-0.01	0.939	0.11 (0 to 0.22)	0.051

^aBased on mixed model statical analysis (P \leq 0.05).

gene expression of antioxidant enzymes (17). Therefore, estrogen increases gene expression of antioxidant enzymes by increasing gene expression of ERa receptors, as well (17). Our findings also showed that MnSOD enzyme gene expression had not significantly increased in active men, however, MnSOD gene expression changes were significant in women. These findings are consistent with those obtained by Ademoglu et al. The investigators reported that MnSOD activity is higher in women than in men (39). Therefore, it is likely to indicate the higher influence of estrogen on MnSOD (37). Investigations by Gottipati et al. showed that estrogen leads to a further increase in MnSOD in women (40).

On the other hand, catalase is an enzyme that causes to produce free radicals through reaction with H2O2 and preventing Fenton reaction (41, 42). In our studies, we found that catalase gene expression in both groups of active men and women had a significant increase. In this context, the role of epinephrine seems important. Studies have shown that epinephrine in addition to driving energy mechanisms to maintain the activity of the body, leads to increased NO production through beta-adrenergic receptors (43). However, there is a significant relationship between NO and catalase. It has been recently reported that NO leads to increased activity of catalase in order to deal with H_2O_2 (44). Thus, it seems that one of the mechanisms to increase catalase in the present study in both groups of men and women, may be related to the increased level of NO in sporting activities.

On the other hand, in the present study gene expression of Cu/ZnSOD did not significantly increase as a result of incremental exercise. These findings are consistent with the research by Sachdev et al. (45). Studies show that active individuals have adapted in part to an increase in free radicals (45, 46). Thus, in this study the level of TAC in both active men and women significantly increased. It also seems that MnSOD prevents initial increase of free radicals to some extent and Cu/ZnSOD enzyme deals with the less challenges to fight free radicals. However, these cannot be accepted with certainty. As the relationship between free radicals and antioxidant enzyme gene expression is also important and should be examined quantitatively.

Our results indicate that increased levels of MDA resulted in a -3.69-unit decrease of MnSOD gene expression

5



Figure 1. MDA (μ m) change in all stages of incremental exercise (A) MnSOD, Cu/ZnSOD and catalase enzymes mRNA (fold) in all stages of incremental exercise (B, C and D). (All statical analysis performed based on repeated measure and Bonferroni test (P \leq 0.05)).

in men and a -0.14-unit decrease in women. However, the numerical relationship is different for Cu/Zn SOD and catalase, and TAC. Thus, our findings suggested that there is an inverse relationship between changes in antioxidant parameters and MDA in men and increase in MDA level is associated with a decrease in the Cu/ZnSOD (-2.62 units), catalase (-0.93 units) and TAC (-0.01 units). But there is positive relationship between changes in these parameters and MDA in women. Thus, in line with changes in MDA, gene expression of Cu/Zn SOD (0.051 units), catalase (0.456 units) and TAC (0.11 units) increased, that seems important immunologically. Since the more the response of the immune system is weakened, the more the injuries will be. In our findings despite the fact that gene expression of catalase, Cu/ZnSOD and Mn SOD enzymes and also TAC level increased in both active men and women as a result of incremental physical activity, but regarding the negative or positive relationships between these indicators MDA, it seems that mitochondria and cytosol of lymphocytic cells of men are more susceptible to damage caused by free radicals compared with active women.

5.1. Conclusions

MnSOD gene expression in active women, despite weakening by the increase in MDA, is still an appropriate response to free radicals and partially prevents the spread

Var			Men	Women		Between Group P
		Mean \pm SD	Median (Range)	Mean \pm SD	Median (Range)	
Cu/Z	n					
	Baseline	3.77 ± 1.88	4.54 (0.5 to 5.62)	5.38 ± 2.09	5.55 (0.33 to 7.72)	0.079 ^a
	After	4.04 ± 0.86	3.87 (3.17 to 6.27)	5.89 ± 2.17	6.73 (2.45 to 9.08)	0.017 ^a
	3 hours after	5.03 ± 2.37	4.8 (0.4 to 8.79)	6.35 ± 2.42	6.08 (3.55 to 10.47)	0.222 ^a
Mn						
	Baseline	1.89 ± 1.77	1.29 (0.02 to 5.99)	3.41 ± 1.16	3.36 (1.77 to 6.25)	0.033 ^a
	After	2.19 ± 1.15	1.91 (0.86 to 4.15)	4.09 ± 1.11	3.89 (2.78 to 5.86)	0.001 ^a
	3 hours after	2.58 ± 2.24	1.82 (0.41 to 8.3)	4.94 ± 0.66	5.22 (3.75 to 5.84)	0.005 ^a
Cat						
	Baseline	5.37 ± 2.33	5.43 (1.12 to 9.73)	7.01 ± 1.91	6.89 (5.01 to 10.76)	0.095 ^a
	After	8.21 ± 1.32	7.98 (5.98 to 10.28)	9.91 ± 3.62	10.34 (3.35 to 15.71)	0.161 ^a
	3 hours after	8.35 ± 1.95	8.17 (5.82 to 13.31)	9.58 ± 2.72	8.87 (6.2 to 14.22)	0.242 ^a
MDA						
	Baseline	2.95 ± 0.84	3.2 (1.2 to 3.8)	2.5 ± 1.24	2.15 (1.2 to 4.2)	0.512 ^b
	After	3.37 ± 0.99	3.6 (1.7 to 4.6)	2.84 ± 1.38	2.75 (1.3 to 4.9)	0.349 ^b
	3 hours after	3.36 ± 0.85	3.6 (1.8 to 4.4)	3.04 ± 1.16	3.35 (1.4 to 4.3)	0.605 ^b

Table 3. Difference Between Antioxidant Enzyme mRNA and MDA Levels in Two Groups

^aBased on t-test ($P \le 0.05$)

^bBased on Mann-Whitney test ($P \le 0.05$)

of the resulting damage. On the other hand, about cytosol protection it is important to note that the immune system of active men will respond to changes in free radicals by increasing catalase. However, in active women the response occurs through increased gene expression of catalase and Cu/ZnSOD. Thus, it appears that the increased free radical during incremental exercise challenges gene expression and activity of antioxidant enzymes. However, despite the negative effects of free radicals on active women, activity and gene expression of antioxidant enzymes is an appropriate response to the free radical and their gene expression and activity increases in line with the increase in MDA.

Acknowledgments

The authors are grateful to the subjects who participated in the study.

Footnotes

Authors' Contribution: Behrouz Baghaiee, contributed to data collection, implementation of the incremental exercise test, laboratory analysis, statical data analysis, reviewed literature and discussed; Behzad Baradaran, Mohammad Reza Aliparasti and Shohreh Almasi; contributed to laboratory analysis; Marefat Siahkuhian, contributed to reviewed literature, discussed and suggested the contents as well as edited the manuscript; Also, all the authors read and approved the final manuscript.

Funding/Support: This study was supported by Mohaghegh Ardabili University.

References

- Bendahan D, Mattei JP, Ghattas B, Confort-Gouny S, Le Guern ME, Cozzone PJ. Citrulline/malate promotes aerobic energy production in human exercising muscle. *Br J Sports Med.* 2002;**36**(4):282–9. [PubMed: 12145119].
- 2. Sanz A, Stefanatos RK. The mitochondrial free radical theory of aging: a critical view. *Curr Aging Sci.* 2008;1(1):10–21. [PubMed: 20021368].
- Brand K. Aerobic glycolysis by proliferating cells: protection against oxidative stress at the expense of energy yield. *J Bioenerg Biomembr.* 1997;29(4):355–64. [PubMed: 9387096].
- Seo AY, Hofer T, Sung B, Judge S, Chung HY, Leeuwenburgh C. Hepatic oxidative stress during aging: effects of 8% long-term calorie restriction and lifelong exercise. *Antioxid Redox Signal.* 2006;8(3-4):529–38. doi: 10.1089/ars.2006.8.529. [PubMed: 16677097].
- Ellis GR, Anderson RA, Lang D, Blackman DJ, Morris RH, Morris-Thurgood J, et al. Neutrophil superoxide anion-generating capacity, endothelial function and oxidative stress in chronic heart failure: effects of short- and long-term vitamin C therapy. J Am Coll Cardiol. 2000;36(5):1474–82. [PubMed: 11079645].

- Roberts CK, Won D, Pruthi S, Kurtovic S, Sindhu RK, Vaziri ND, et al. Effect of a short-term diet and exercise intervention on oxidative stress, inflammation, MMP-9, and monocyte chemotactic activity in men with metabolic syndrome factors. *J Appl Physiol (1985)*. 2006;**100**(5):1657-65. doi: 10.1152/japplphysiol.01292.2005. [PubMed: 16357066].
- 7. Mittler R. Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci.* 2002;7(9):405-10. [PubMed: 12234732].
- 8. Betteridge DJ. What is oxidative stress?.*Metabolism*. 2000;**49**(2 Suppl 1):3-8. [PubMed: 10693912].
- Sies H, Cadenas E. Oxidative stress: damage to intact cells and organs. *Philos Trans R Soc Lond B Biol Sci.* 1985;311(1152):617–31. [PubMed: 2869521].
- Gomez-Cabrera MC, Domenech E, Vina J. Moderate exercise is an antioxidant: upregulation of antioxidant genes by training. *Free Radic Biol Med.* 2008;44(2):126–31. doi: 10.1016/j.freeradbiomed.2007.02.001. [PubMed: 18191748].
- Lambertucci RH, Levada-Pires AC, Rossoni LV, Curi R, Pithon-Curi TC. Effects of aerobic exercise training on antioxidant enzyme activities and mRNA levels in soleus muscle from young and aged rats. *Mech Ageing Dev.* 2007;**128**(3):267-75. doi: 10.1016/j.mad.2006.12.006. [PubMed: 17224177].
- Vollaard NB, Shearman JP, Cooper CE. Exercise-induced oxidative stress:myths, realities and physiological relevance. *Sports Med.* 2005;**35**(12):1045–62. [PubMed: 16336008].
- Sverko V, Sobocanec S, Balog T, Marotti T. Age and gender differences in antioxidant enzyme activity: potential relationship to liver carcinogenesis in male mice. *Biogerontology*. 2004;5(4):235–42. doi: 10.1023/B:BGEN.0000038024.58911.6e. [PubMed: 15314273].
- Strehlow K, Rotter S, Wassmann S, Adam O, Grohe C, Laufs K, et al. Modulation of antioxidant enzyme expression and function by estrogen. *Circ Res.* 2003;93(2):170–7. doi: 10.1161/01.RES.0000082334.17947.11. [PubMed: 12816884].
- Moorthy K, Sharma D, Basir SF, Baquer NZ. Administration of estradiol and progesterone modulate the activities of antioxidant enzyme and aminotransferases in naturally menopausal rats. *Exp Gerontol.* 2005;**40**(4):295–302. doi: 10.1016/j.exger.2005.01.004. [PubMed: 15820610].
- Mann V, Huber C, Kogianni G, Collins F, Noble B. The antioxidant effect of estrogen and Selective Estrogen Receptor Modulators in the inhibition of osteocyte apoptosis in vitro. *Bone.* 2007;**40**(3):674–84. doi: 10.1016/j.bone.2006.10.014. [PubMed: 17174166].
- Baltgalvis KA, Greising SM, Warren GL, Lowe DA. Estrogen regulates estrogen receptors and antioxidant gene expression in mouse skeletal muscle. *PLoS One*. 2010;5(4):e10164. doi: 10.1371/journal.pone.0010164. [PubMed: 20405008].
- Lim W, Park Y, Cho J, Park C, Park J, Park YK, et al. Estrogen receptor beta inhibits transcriptional activity of hypoxia inducible factor-1 through the downregulation of arylhydrocarbon receptor nuclear translocator. *Breast Cancer Res.* 2011;13(2):R32. doi: 10.1186/bcr2854. [PubMed: 21435239].
- Gavrilovic L, Stojiljkovic V, Kasapovic J, Popovic N, Pajovic SB, Dronjak S. Treadmill exercise does not change gene expression of adrenal catecholamine biosynthetic enzymes in chronically stressed rats. *Anais Academia Brasileira Ciencias*. 2013;85(3):999–1012.
- Berzosa C, Cebrian I, Fuentes-Broto L, Gomez-Trullen E, Piedrafita E, Martinez-Ballarin E, et al. Acute exercise increases plasma total antioxidant status and antioxidant enzyme activities in untrained men. *J Biomed Biotechnol.* 2011;2011:540458. doi: 10.1155/2011/540458. [PubMed: 21436993].
- Clarkson PM, Thompson HS. Antioxidants: what role do they play in physical activity and health?. *Am J Clin Nutr.* 2000;**72**(2 Suppl):637S– 46S. [PubMed: 10919970].
- 22. Elosua R, Molina L, Fito M, Arquer A, Sanchez-Quesada JL, Covas MI, et al. Response of oxidative stress biomarkers to a 16-week aerobic physical activity program, and to acute physical activity, in healthy

young men and women. *Atherosclerosis*. 2003;**167**(2):327-34. [PubMed: 12818416].

- Ainsworth BE, Haskell WL, Leon AS, Jacobs JDR, Montoye HJ, Sallis JF, et al. Compendium of physical activities: Classification of energy costs of human physical activities. *Med Sci Sports Exerc.* 1993;25(1):71-80.
- 24. Manini TM, Everhart JE, Patel KV, Schoeller DA, Colbert LH, Visser M, et al. Daily activity energy expenditure and mortality among older adults. *JAMA*. 2006;**296**(2):171–9. doi: 10.1001/jama.296.2.171. [PubMed: 16835422].
- Draper HH, Hadley M. Malondialdehyde determination as index of lipid peroxidation. *Methods Enzymol.* 1990;186:421–31. [PubMed: 2233309].
- Jowko E, Dlugolecka B, Makaruk B, Cieslinski I. The effect of green tea extract supplementation on exercise-induced oxidative stress parameters in male sprinters. *Eur J Nutr.* 2015;**54**(5):783-91. doi: 10.1007/s00394-014-0757-1. [PubMed: 25120110].
- Gordon L, McGrowder DA, Pena YT, Cabrera E, Lawrence-Wright M. Effect of exercise therapy on lipid parameters in patients with endstage renal disease on hemodialysis. *J Lab Physicians*. 2012;**4**(1):17–23. doi: 10.4103/0974-2727.98665. [PubMed: 22923917].
- McCully JD, Rousou AJ, Parker RA, Levitsky S. Age- and gender-related differences in mitochondrial oxygen consumption and calcium with cardioplegia and diazoxide. *Ann Thorac Surg.* 2007;83(3):1102–9. doi: 10.1016/j.athoracsur.2006.10.059. [PubMed: 17307466].
- Misiak M, Beyer C, Arnold S. Gender-specific role of mitochondria in the vulnerability of 6-hydroxydopamine-treated mesencephalic neurons. *Biochim Biophys Acta*. 2010;**1797**(6-7):1178–88. doi: 10.1016/j.bbabio.2010.04.009. [PubMed: 20416276].
- Ide T, Tsutsui H, Ohashi N, Hayashidani S, Suematsu N, Tsuchihashi M, et al. Greater oxidative stress in healthy young men compared with premenopausal women. *Arterioscler Thromb Vasc Biol.* 2002;22(3):438– 42. [PubMed: 11884287].
- Ochs-Balcom HM, Grant BJ, Muti P, Sempos CT, Freudenheim JL, Browne RW, et al. Oxidative stress and pulmonary function in the general population. *Am J Epidemiol.* 2005;**162**(12):1137–45. doi: 10.1093/aje/kwi339. [PubMed: 16269582].
- Barreiro E, Coronell C, Lavina B, Ramirez-Sarmiento A, Orozco-Levi M, Gea J, et al. Aging, sex differences, and oxidative stress in human respiratory and limb muscles. *Free Radic Biol Med.* 2006;41(5):797-809. doi:10.1016/j.freeradbiomed.2006.05.027. [PubMed: 16895800].
- 33. Constantin-Teodosiu D, Young S, Wellock F, Short AH, Burden RP, Morgan AG, et al. Gender and age differences in plasma carnitine, muscle strength, and exercise tolerance in haemodialysis patients. *Nephrol Dial Transplant.* 2002;**17**(10):1808–13. [PubMed: 12270989].
- 34. Tarnopolsky MA, Rennie CD, Robertshaw HA, Fedak-Tarnopolsky SN, Devries MC, Hamadeh MJ. Influence of endurance exercise training and sex on intramyocellular lipid and mitochondrial ultrastructure, substrate use, and mitochondrial enzyme activity. *Am J Physiol Regul Integr Comp Physiol.* 2007;**292**(3):R1271-8. doi: 10.1152/ajpregu.00472.2006. [PubMed: 17095651].
- Bellanti F, Matteo M, Rollo T, De Rosario F, Greco P, Vendemiale G, et al. Sex hormones modulate circulating antioxidant enzymes: impact of estrogen therapy. *Redox Biol.* 2013;1:340–6. doi: 10.1016/j.redox.2013.05.003. [PubMed: 24024169].
- Cunningham M, Gilkeson G. Estrogen receptors in immunity and autoimmunity. *Clin Rev Allergy Immunol.* 2011;40(1):66–73. doi: 10.1007/s12016-010-8203-5. [PubMed: 20352526].
- Kovats S. Estrogen receptors regulate an inflammatory pathway of dendritic cell differentiation: mechanisms and implications for immunity. *Horm Behav.* 2012;62(3):254–62. doi: 10.1016/j.yhbeh.2012.04.011. [PubMed: 22561458].
- Stirone C, Duckles SP, Krause DN, Procaccio V. Estrogen increases mitochondrial efficiency and reduces oxidative stress in cerebral blood vessels. *Mol Pharmacol.* 2005;68(4):959–65. doi: 10.1124/mol.105.014662. [PubMed: 15994367].

- Ademoglu E, Ozcan K. Age-related changes in the activity and expression of manganese superoxide dismutase, and mitochondrial oxidant generation in female and male rats. *Turk J Biochem.* 2013;38(4):445–50.
- Gottipati S, Cammarata PR. Mitochondrial superoxide dismutase activation with 17 beta-estradiol-treated human lens epithelial cells. *Mol Vis.* 2008;14:898–905. [PubMed: 18490963].
- Nagy K, Pasti G, Bene L, Nagy I. Involvement of Fenton reaction products in differentiation induction of K562 human leukemia cells. *Leuk Res.* 1995;19(3):203–12. [PubMed: 7700081].
- Tauler P, Aguilo A, Gimeno I, Fuentespina E, Tur JA, Pons A. Response of blood cell antioxidant enzyme defences to antioxidant diet supplementation and to intense exercise. *Eur J Nutr.* 2006;45(4):187–95. doi: 10.1007/s00394-005-0582-7. [PubMed: 16365696].
- Kou R, Michel T. Epinephrine regulation of the endothelial nitricoxide synthase: roles of RAC1 and beta3-adrenergic receptors in endothelial NO signaling. *J Biol Chem.* 2007;**282**(45):32719–29. doi: 10.1074/jbc.M706815200. [PubMed: 17855349].
- 44. Yoshioka Y, Kitao T, Kishino T, Yamamuro A, Maeda S. Nitric oxide protects macrophages from hydrogen peroxide-induced apoptosis by inducing the formation of catalase. *J Immunol.* 2006;**176**(8):4675– 81. [PubMed: 16585560].
- Sachdev S, Davies KJ. Production, detection, and adaptive responses to free radicals in exercise. *Free Radic Biol Med.* 2008;44(2):215–23. doi: 10.1016/j.freeradbiomed.2007.07.019. [PubMed: 18191757].
- Fisher-Wellman K, Bloomer RJ. Acute exercise and oxidative stress: a 30 year history. *Dyn Med.* 2009;8:1. doi: 10.1186/1476-5918-8-1. [PubMed: 19144121].