Veterinary Research Forum

Journal Homepage: vrf.iranjournals.ir

# Effect of curcumin on rat sperm morphology after the freeze-thawing process

Ali Soleimanzadeh, Adel Saberivand\*

Department of Clinical Sciences, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran.

Article Info	Abstract		
Article history:	Reactive oxygen species (ROS) generation, induced by the cryopreservation process, can be		
	responsible for mammalian sperm damage. Curcumin is known as an effective antioxidant		
Received: 02 March 2013	against oxidative stress. The aim of this study was to evaluate the effects of curcumin on sperm		
Accepted: 10 June 2013	count, motility and viability, semen total antioxidant capacity and DNA integrity of rat		
Available online: 15 September 2013	spermatozoa during semen freeze-thawing process. Sperm collected from 10 adult rats were		
	divided into two groups (n=10 for each group): control and a test group supplemented with 2.5		
Key words:	mM curcumin. After freezing-thawing, the number of spermatozoa, motility, viability, total		
	antioxidant capacity (TAC), and DNA integrity of the sperm were analyzed. Motility, viability		
Curcumin	and DNA integrity of sperm were significantly preserved in treatment groups compared to the		
Rat	control ( $p < 0.05$ ) after freezing-thawing. Following cryopreservation, TAC was significantly		
Sperm parameters	preserved in thawing semen supplemented with curcumin compared to the control group		
	(p<0.05). Based on our results, it is concluded that curcumin addition during freezing resulted		
	in positive effects on sperm parameters after thawing in adult rats.		
	© 2013 Urmia University. All rights reserved.		

# اثر کورکومین بر ریخت شناسی اسپرم موش صحرایی بعد از فرایند انجماد و ذوب

#### چکیدہ

تولید گونه های اکسیژن فعال ، ناشی از فرایند انجماد، می تواند باعث آسیب دیدن اسپرم پستانداران باشد. کورکومین به عنوان آنتی اکسیدان موثر در برابر استرس اکسیداتیو شناخته شده است. هدف از این مطالعه بررسی اثرات آنتی اکسیدانی کورکومین در تعداد اسپرم، تحرک و زنده ماندن، ظرفیت آنتی اکسیدانی تام منی و یکپارچگی DNA اسپرم موش صحرایی طی فرآیند انجماد مایع منی می باشد. اسپرم جمع آوری شده از ۱۰ قطعه موش صحرایی بالغ به دو گروه (۱۰ قطعه برای هر گروه) تقسیم شدند: گروه شاهد و یک گروه آزمون با افزودن ۲/۵ میلی مول کورکومین. پس از فرایند انجماد و ذوب، تعداد اسپرم، تحرک، زنده ماندن، توان آنتی اکسیدانی تام منی و ریک گروه آزمون با افزودن ۲/۵ میلی مول کورکومین. پس از نوریند انجماد و ذوب، تعداد اسپرم، تحرک، زنده ماندن، توان آنتی اکسیدانی تام منی و یکپارچگی DNA اسپرم موش صحرایی طی قرابل توجهی در تحرک، زنده ماندن و یکپارچگی DNA اسپرم در گروه درمان نسبت به گروه شاهد و جود داشت (۲۰۰۵). پس از انجماد منجر به اثرات مول توجهی در اسپرم قابل توجهی در کورکومین در مقایسه با گروه شاهد بیشتر بود (۲۰۰ مان نتیج به دست آمده، می توان نتیجه گرفت که افزودن کور کومین بر پر اسپرم و با افزودن د/۲

**واژه های کلیدی:** پارامترهای اسپرم، کورکومین ، موش صحرایی

\*Correspondence:

Adel Saberivand. DVM, PhD

Department of Clinical Sciences, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran. **E-mail:** adelsaberivand@yahoo.ca

## Introduction

Semen freezing and thawing procedures produce membrane damage that reduce longevity of sperm in the female genital tract, with a concomitant decrease in fertility.<sup>1</sup> The success of semen freezing relies upon an understanding of internal (inherent characteristics of spermatozoa) and external (composition of diluents, type and concentration of cryoprotective agents, rates of dilution and cooling, equilibration, and method of freezing and thawing of semen) factors and their interactions which can influence the capacity of sperm to survive during freezing and thawing.<sup>2</sup>

Reactive oxygen species (ROS) generation, induced by cryopreservation process, can be responsible for mammalian sperm damage.<sup>3-5</sup> The ROS production has been associated with reduction of sperm motility, decreased capacity for spermoocyte fusion and infertility. Against ROS attack, sperm cells are well equipped with a powerful defense system of antioxidants, but an imbalance between the production of ROS and the available antioxidant-defenses results in oxidative stress.<sup>6</sup> A variety of antioxidants have been tested to either scavenge ROS directly or effects of counter ROS toxicity in semen of a variety of mammalian and avian species.<sup>7</sup>

Antioxidants are the frontline of defense against free radicals.8 Curcumin (1,7-bis [4-hydroxy-3-methoxyphenyl]-1,6-heptadiene-3,5-dione) is the principal curcuminoid found in turmeric and is generally considered as its most active constituent.910 Turmeric exhibits anti-tumor, anti-infectious and anti-inflammatory activities with low toxicity.<sup>11-13</sup> The most important feature of curcumin is that it has no side effects despite being a therapeutic agent with multiple beneficial functions.<sup>14</sup> Curcumin is considered to be an effective antioxidant against oxidative tissue damage. The antioxidant mechanism of curcumin is due to its specific conjugated structure of two methoxylated phenols and an enol form of β-diketone. This structure is responsible for free radical trapping ability as a chain breaking antioxidant.<sup>15</sup> It can significantly inhibit generation of ROS both in vitro and in vivo.16 Protective effects of curcumin on the testis has been reported in oxidative damage induced by cisplatin, sodium arsenite, aflatoxin, ischemia-reperfusion injury, di-nbutyl phthalate in laboratory animals.12,17-20

Several numbers of studies have established the effectiveness of antioxidants like curcumin against oxidative stress.<sup>21-24</sup> Curcumin is reported to protect male reproductive system in chromium-induced oxidative damage.<sup>25</sup> The aims of this study were to evaluate the effects of curcumin on sperm count, motility, viability, semen TAC and DNA integrity of rat spermatozoa supplemented with curcumin during semen freeze-thawing process.

## **Materials and Methods**

All chemicals used in the experiment were purchased from Sigma-Aldrich, Munich, Germany unless otherwise stated.

**Experimental animals.** Ten adult sexually mature male (4 months of age weighing 200-230 g) albino Wistar rats were obtained from animal house of the Faculty of Veterinary Medicine, Urmia University. They were housed in a specific pathogen-free environment under standard conditions of temperature ( $23 \pm 1 \degree$ C), relative humidity ( $55 \pm 10\%$ ), and 12/12 hr light/dark cycle, and fed with a standard pellet diet and water. All ethical themes of the studies on animals were considered carefully.

**Semen collection**. Epididymal sperms were collected by slicing the cauda region of the epididymis in 5 mL of HTF and incubated for 5 min at 37 °C in an atmosphere of 5% CO<sub>2</sub> to allow sperm to swim out of the epididymal tubules. After collection, semen samples were divided into two groups (n=10 for each group). One control group without antioxidants and one test group supplemented with 2.5 mM curcumin<sup>26</sup>. All samples were packed in 0.5 mL French straws and were frozen in liquid nitrogen vapor, (4 cm above the liquid nitrogen), for 15 min and then plunged into liquid nitrogen for storage. The frozen straws were then thawed individually (37 °C for 20 sec) in a water bath. In frozen thawed semen samples, the sperm motility, count, live percent, DNA integrity and semen total antioxidant capacity (TAC) were evaluated using conventional methods.

**Sperm motility.** Assessment of sperm motility was done according to WHO laboratory manual protocol for the examination of human semen and sperm-cervical mucus interaction.<sup>27</sup> In brief, 10  $\mu$ L of the sperm suspension was placed on semen analysis chamber. A minimum of five microscopic fields were assessed to evaluate sperm motility on at least 200 sperm for each animal.

**Sperm viability.** Eosin-nigrosin staining was used to assess sperm viability according to WHO protocol. Briefly, eosin (Merck, Darmstadt, Germany) and nigrosin (Merck, Darmstadt, Germany) was prepared in distilled water. One volume of sperm suspension was mixed with two volume of 1% eosin. After 30 sec, an equal volume of nigrosin was added to this mixture. Thin smears were then prepared and observed under a light microscope (Model CHT, Olympus optical Co. Ltd, Tokyo, Japan) at 1000× magnification. Viable sperm remained colorless while nonviable sperm was stained red.

**Sperm morphology.** For the analysis of morphological abnormalities, sperm smears were drawn on clean and grease-free slides, and allowed to air-dry overnight. The slides were stained with 1% eosin-Y/5% nigrosin and examined at 400× for morphological abnormalities in each sample.

Assessment of DNA integrity. Thin smears were prepared from the sperm solution and allowed to air-dry. To test sperm DNA integrity, the smears were stained with acridine orange (AO). The AO staining was performed according to a protocol described by Tejada *et al.*<sup>28</sup> In brief, the smears were fixed for 14 hr in methanol/acetic acid (3:1) at 4 °C and stained with AO solution (0.19% in phosphate citrate buffer, pH = 2.5) for 10 min. The slides were gently

washed by distilled water for 5 min and air dried. The stained smears were then observed under fluorescence microscope (Model GS7, Nikon Co, Tokyo, Japan) at 1000× magnification. Three types of staining patterns were considered in sperm head; green spermatozoa (double-stranded DNA), yellow and red spermatozoa (single-stranded DNA). At least 100 spermatozoa per slide were counted to evaluate the percentage of double-stranded DNA in the spermatozoa.

Total antioxidant capacity evaluation. The TAC content of the seminal plasma was determined by the ferric reducing antioxidant power (FRAP) assay.<sup>29</sup> In brief, 300 mmol L<sup>-1</sup> acetate buffer of pH 3.6, 10 mmol L<sup>-1</sup> 2,4,6-tri-(2-pyridyl)-1,3,5-triazine, 98.00% and 20 mmol L<sup>-1</sup> FeCl<sub>3</sub>·6H<sub>2</sub>O were mixed together in the ratio of 10:1:1, respectively, to give the working FRAP reagent. A 50 µL aliquot of sperm samples was added to 1 mL of FRAP reagent in a semi-micro plastic cuvette. Absorbance measurement was taken at 593 nm (A593) exactly 10 min after mixing using 50 µL of water as the reference. To standardize, 50 µL of the standard (FeSO<sub>4</sub>·7H<sub>2</sub>O, 1 mmol L<sup>-1</sup>) was added to 1 mL of FRAP reagent.

**Statistical analysis.** The data analysis was performed by analysis of variance (ANOVA) and Duncan tests using SPSS (Version 17 SPSS Inc., Chicago, IL, USA). A value of p < 0.05 was considered as statistically significant.

#### Results

**Sperm count.** There was no significant difference between sperm number in curcumin group compared to that of the control group (Table 1).

**Sperm motility**. The sperm motility was significantly (p < 0.05) preserved in curcumin group compared to that of the control group (Table 1).

**Sperm viability.** The percentage of viable sperm was significantly (p < 0.05) preserved in curcumin group compared to that of the control group (Fig. 1 and Table 1).

**Sperm morphological anomalies.** Spermatozoa stained with eosin-nigrosin showed that addition of curcumin had no significant effect on the sperm morphology compared to that of the control group (Table 1).

**Sperm DNA integrity.** Thawed spermatozoa stained with AO showed that addition of curcumin had a significant effect on the sperm DNA integrity compared to that of the control group (Fig. 2 and Table 1).

**Table 1.** Sperm number, viability, morphological anomalies, motility, TAC and DNA integrity (AO staining) in frozen-thawed rat semen. Data are presented as mean  $\pm$  SD (n = 10).

<u> </u>		
Sperm characteristics	Control	Curcumin
Number (10 <sup>6</sup> )	20.92 ± 0.14	20.96 ± 0.24
Viability (%)	66.07 ± 0.51	72.96 ± 0.09 *
Morphological anomalies (%)	$1.21 \pm 0.01$	$1.20 \pm 0.01$
Motility (%)	67.58 ± 0.34	73.61 ± 0.44 *
DNA integrity (%)	$80.08 \pm 0.27$	99.19 ± 0.19 *
TAC (µmol L-1)	$62.32 \pm 0.17$	70.16 ± 0.26 *

\* indicates significantly different from controls (*p* < 0.05).

**Sperm Total antioxidant capacity (TAC).** The TAC of sperm was significantly (p < 0.05) preserved in curcumin group compared to that of the control (Table 1).



Fig. 1. Sperm viability in curcumin group; A) Dead sperm (red), B) Viable sperm (colorless), (Black arrows); (Eosin/nigrosin, 1000×).



**Fig. 2.** Rat spermatozoa; **A)** Normal sperm (green), **B)** Damaged DNA (yellow), (White arrows); (A0, 1000×).

# Discussion

Cryopreservation increases the percentage of dead or membrane damaged sperm.<sup>30</sup> The process of cryopreservation is known to cause more production of reactive oxygen species, as antioxidant defenses are reduced in the process.<sup>31</sup> Hence, addition of antioxidants may have beneficial effects on spermatozoa function during the cryopreservation. Curcumin has been shown to have strong antioxidant activity and reduces oxidative stress.<sup>23,24</sup> The present study reports the ameliorative effects of curcumin on the motility, viability, TAC and DNA integrity of rat frozen-thawed sperm.

The spermatozoon motility and viability generally declines following cryopreservation procedure. There is a remarkable decrease in viability and motility caused by ROS resulting in low conception rate following the use of frozen-thawed sperm for artificial insemination.<sup>32</sup> Based on our results, in frozen-thawed semen, curcumin addition had a positive impact on both motility and viability. One of the possible ameliorative mechanisms of curcumin on the above mentioned parameters is to scavenge the free radicals and thereby act as good antioxidants. Another reason for enhancement of sperm motility in frozen semen observed in this study may be due to the increasing of TAC level of semen in curcumin supplemented group and the correlation between total antioxidant capacity in seminal plasma and sperm motility.<sup>33</sup> These results were in agreement with the findings of earlier studies on the ameliorative effect of curcumin on the spermatozoa motility in metronidazoletreated mice.<sup>34</sup>

The integrity of sperm DNA is an important factor for the success of fertilization as well as normal development of the embryo, fetus and newborn.<sup>35</sup> Several studies have shown the effect of cryopreservation procedure on DNA damages through induction of oxidative stress and the generation of ROS.<sup>36,37</sup> In this study, a significant preservation of DNA integrity in post-thaw in curcumin supplemented group compared to the control, was most probably due to increase in its antioxidant capacity.

The TAC in seminal plasma is closely related to male fertility, appropriate TAC provides a favorable environment for sperm swimming. The decreased level of TAC in seminal plasma may be one of the causes of male infertility.<sup>33</sup> In addition, Contri *et al.* reported a positive correlation between sperm parameters and total antioxidant capacity in seminal plasma.<sup>38</sup>Semen processing and cryopreservation decrease the antioxidant defense capacity of semen. Based on our results addition of curcumin to fresh sperm of rats before freezing, significantly (p < 0.05) preserved the TAC level in thawing seminal plasma.

Based on our results, it is concluded that addition of curcumin during cooling, resulted in positive effects on rat sperm motility, viability, DNA integrity and TAC level after thawing .Such data could help to develop and improve semen handling and storage techniques. Future research should address molecular mechanisms underlying the protective effects of curcumin on rat sperm.

## Acknowledgements

Authors would like to gratefully thank Dr. Abbas Ahmadi for his assistance in laboratory work. We also thank Mr. Reza Aliyari for his technical assistance.

#### References

- 1. Satorre MM, Breininger E, Beconi MT. Cryopreservation with  $\alpha$ -tocopherol and Sephadex filtration improved the quality of boar sperm. Theriogenology 2012; 78:1548-1556.
- 2. Johnson LA, Weitze KF, Fiser P, et al. Storage of boar semen. Anim Reprod Sci 2000; 62:143-172.
- Griveau JF, Le Lannou D. Reactive oxygen species and human spermatozoa: Physiology and pathology. Int J Androl 1997; 20:61-69.
- 4. Calamera JC, Neild DM, Fernandez PJ, et al. Effects of long-term *in vitro* incubation of human spermatozoa: Functional parameters and catalase effect. Andrologia 2001; 33:79-86.
- 5. Gabella BM, Chaves MG, Miragaya MH, et al. Membrane changes during different stages of a freeze-thaw protocol for equine semen cryopreservation. Theriogenology 2003; 59:1693-1705.

- 6. Sikka SC. Oxidative stress and role of antioxidants in normal and abnormal sperm function. Front Biosci 1996; 1:78-86.
- 7. Aitken RJ, Clarkson JS, Fishel S. Generation of reactive oxygen species, lipid peroxidation, and human sperm function. Biol Reprod 1989; 141:183-197.
- 8. Osawa T, Kato Y. Protective role of antioxidative food factors in oxidative stress caused by hyperglycemia. Ann N Y Acad Sci 2005; 1043:440-451.
- 9. Duvoix A, Blasius R, Delhalle S, et al. Chemopreventive and therapeutic effects of curcumin. Cancer Lett 2005; 223:181-190.
- 10. Sharma RA, Gescher AJ, Steward WP. Curcumin: The story so far. Eur J Cancer 2005; 541: 1955-1968.
- Cohly HH, Taylor A, Angel MF, et al. Effect of turmeric, turmerin and curcumin on H<sub>2</sub>O<sub>2</sub>-induced renal epithelial (LLC-PK1) cell injury. Free Radic Biol Med 1998; 24:49-54.
- 12. Farombi EO, Abarikwu SO, Adedara IA, et al. Curcumin and kolaviron ameliorate di-n-butyl phthalate-induced testicular damage in rats. Basic Clin Pharmacol Toxicol 2007; 100:43-48.
- Mathuria N, Verma RJ. Curcumin ameliorates aflatoxininduced toxicity in mice spermatozoa. Fertil Steril 2008; 90:775-780.
- 14. Khanna NM. Turmeric Nature's precious gift. Curr Sci 1999; 76(10):1351-1356.
- 15. Masuda T, Maekawa T, hidaka K, et al. Chemical studies on antioxidant mechanisms of curcumin: analysis of oxidative coupling products from curcumin and linoleate. J Agric Food Chem 2001; 49:2539-2547.
- 16. Joe B, Lokesh BR. Role of capsaicin, curcumin and dietary n-3 fatty acids in lowering the generation of reactive oxygen species in rat peritoneal macrophages. Biochem Biophys Acta 1994; 1224: 255-263.
- 17. Mathuria N, Verma RJ. Curcumin ameliorates aflatoxin-induced lipid peroxidation in liver, kidney and testis of mice an *in vitro* study. Acta Pol Pharm 2007; 64:413-416.
- 18. El-Demerdash FM, Yousef MI, Radwan FM. Ameliorating effect of curcumin on sodium arsenite-induced oxidative damage and lipid peroxidation in different rat organs. Food Chem Toxicol 2009; 47:249-254.
- 19. Ilbey YO, Ozbek E, Cekmen M, et al. Protective effect of curcumin in cisplatin-induced oxidative injury in rat testis: mitogen-activated protein kinase and nuclear factor-kappa B signaling pathways. Hum Reprod 2009; 24: 1717-1725.
- 20. Wei SM, Yan ZZ, Zhou J. Curcumin attenuates ischemiareperfusion injury in rat testis. Fertil Steril 2009; 91: 271-277.
- 21. Subudhi U, Das K, Paital B, et al. Alleviation of enhanced oxidative stress and oxygen consumption of l-thyroxine induced-hyperthyroid rat liver mitochondria by vitamin E and curcumin. Chem Biol Interact 2008; 173(2): 105-114.

- 22. Naito M, Wu X, Nomura H, et al. The protective effects of tetrahydrocurcumin on oxidative stress in cholesterol-fed rabbits. J Atheroscler Thromb 2002; 9: 243-250.
- 23. Igbal M, Sharma SD, Okazaki Y, et al. Dietary supplementation of curcumin enhances antioxidant and phase II metabolizing enzymes in ddY male mice: Possible role in protection against chemical carcinogenesis and toxicity. Pharmacol Toxicol 2003; 92:33-38.
- 24. Cheng AL, Hsu CH, Lin JK, et al. Phase I clinical trial of curcumin, a chemopreventive agent, in patients with high-risk or pre-malignant lesions. Anticancer Res 2001; 21:2895-2900.
- 25. Chandra AK, Chatterjee A, Ghosh R, et al. Effect of curcumin on chromium-induced oxidative damage in male reproductive system. Environ Toxicol Pharmacol 2007; 24:160-166.
- 26. Tin Oo Khor, Ying Huang, Tien-Yuan Wu, et al. Pharmaco-dynamics of curcumin as DNA hypomethylation agent in restoring the expression of Nrf2 via promoter CpGs demethylation. Biochem Pharmacol 2011; 82: 1073-1078.
- 27. World Health Organization (WHO). WHO laboratory manual for the examination of human semen and sperm-cervical mucus interaction. Cambridge, UK: Cambridge University Press. Available at: http://whqlib doc.who.int/publications/2010/9789241547789\_eng. pdf. Accessed: Dec 12, 2012.
- Tejada RI, Mitchell JC, Norman A, et al. A test for the practical evaluation of male fertility by acridine orange (AO) fluorescence. Fertil Steril 1984; 42:87-91.
- 29. Benzie IF, Strain JJ. The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": The FRAP assay. Anal Biochem 1996; 239(1):70-76.

rC

- 30. Saacke R. Semen quality: importance of and influencing factors. In proceedings: The 10<sup>th</sup> NAAB technical conference on artificial insemination reproduction, Milwaukee, WI, Columbia, MO: National association of animal breeders.1984; 30-36.
- 31. Bilodeau JF, Blanchette S, Gagnon C, et al. Thiols prevent H<sub>2</sub>O<sub>2</sub>-mediated loss of sperm motility in cryopreserved bull semen. Theriogenology 2001; 56:275-286.
- 32. Marcus-Braun N, Braun G, Potashnik G, et al. Effect of cryopreservation on quality and fertilization capacity of human sperm. Eur J Obstet Gynecol Reprod Biol 2004; 116:63-66.
- 33. Shi YC, Shang XJ, Wang XL, et al. Correlation of total antioxidant capacity in seminal plasma with sperm motility of infertile men. Natl J Androl 2006; 12 (8):703-705.
- 34. Karbalay-Doust S, Noorafshan A. Ameliorative effects of curcumin on the spermatozoon tail length, count, motility and testosterone serum level in metronidazoletreated mice. Prague Med Rep 2011; 112(4): 288-297.
- 35. Zhu P, Ma Y, Huang Y. Role of sperm DNA integrity in male infertility. Natl J Androl 2004; 10:222-226.
- 36. Waterhouse KE, Gjeldnes A, Tverdal A, et al. Alterations of sperm DNA integrity during cryopreservation procedure and in vitro incubation of bull semen. Anim Reprod Sci 2010; 117(1-2): 34-42.
- Gandini L, Lombardo F, Lenzi A, et al. Cryopreservation and sperm DNA integrity. Cell Tissue Bank 2006; 7(2):91-98.
- 38. Contri A, De Amicis I, Molinari A, et al. Effect of dietary antioxidant supplementation on fresh semen quality in stallion. Theriogenology 2011; 75:1319-1326.