

Original article**The prevalence of bacterial contamination of table eggs from retail markets by *Salmonella* spp., *Listeria monocytogenes*, *Campylobacter jejuni* and *Escherichia coli* in Shahrekord, Iran**

Hajieh Ghasemian Safaei, PhD¹
Mohammad Jalali, PhD²
Ahad Hosseini, DVM³
Tahmineh Narimani, MSc¹
Ali Sharifzadeh, PhD³
Ebrahim Raheimi, PhD³

¹Department of Microbiology,
Isfahan University of Medical
Sciences, Isfahan, Iran

²Public Health School, Isfahan
University of Medical Sciences,
Isfahan, Iran

³Department of Food Hygiene,
Azad University of Shahrekord,
Shahrekord, Iran

Address for correspondence:

Dr. Hajieh Ghasemian Safaei,
Department of Microbiology,
Isfahan University of Medical
Sciences, Isfahan, Iran
Tel: +98311 7922469
Fax: +98311 6688597
Email: ghasemian@med.mui.ac.ir

How to cite this article:

Ghasemian Safaei H, Jalali M,
Hosseini A, Narimani T,
Sharifzadeh A, Raheim E. The
prevalence of bacterial
contamination of table eggs from
retails markets by *Salmonella* spp.,
Listeria monocytogenes,
Campylobacter jejuni and
Escherichia coli in Shahrekord,
Iran. Jundishapur J Microbiol.
2011; 4(4): 249-253.

Received: August 2010

Accepted: February 2011

Abstract

Introduction and objective: Contaminated egg and its products are increasing the risks of illness in humans. The significance of these diseases in humans can vary from mild symptoms to life threatening conditions. This study was conducted to determine the contamination of egg to food borne pathogens; *Salmonella* spp., *Listeria monocytogenes*, *Campylobacter jejuni* and *Escherichia coli* in Shahrekord.

Materials and methods: One hundred normal eggs randomly purchased in the spring and summer of 2008 from small and big supermarkets of Shahrekord, and delivered to the food microbiology lab to be tested. The contents of eggs cultured for those bacteria on selective agar and standard microbiological tests performed to identify the isolated organism.

Results: The result showed that there was no contamination by *Salmonella* spp., *L. monocytogenes* and *C. jejuni* in all 100 eggs. However, 19 samples were contaminated by *E. coli*, four samples by *Proteus* spp. and one sample by *Klebsiella* spp. Average colony count of coli form bacteria was 20cfu/g and *E. coli* was 12/6cfu/g.

Conclusion: We concluded that *Salmonella* spp., *L. monocytogenes* and *C. jejuni* contamination of eggs does not make up a serious health hazard in this area. *E. coli* are known to contaminate the surface of egg while mechanical process can spread the bacteria through eggs.

Significance and impact of the study: It is important to remember that control is required at all levels in the food chain and by separating cooked and raw.

Keywords: Egg; *Salmonella* spp.; *Listeria monocytogenes*; *Campylobacter jejuni*; *Escherichia coli*

Introduction

Microbial contamination of egg has important outcome to the poultry industry and illness from contaminated egg is a serious public health problem around the world. The significance of these diseases in humans can vary from mild symptoms to life threatening situation [1]. The egg and its products are an important component source of necessary nutrients and a major food within the human diet. In spite of the antibacterial factors, it can be infected with different bacteria such as *Salmonella* spp., *Listeria monocytogens*, *Campylobacter jejuni* and *Escherichia coli*. *Campylobacter* is the most common identified cause of food borne disease.

It has been found mostly in poultry, egg, red meat, unpasteurized milk and untreated water. The egg can act as a vector in the transmission of food poisoning organism. Although it doesn't grow in food, it spreads easily, so only a few bacteria in a piece of undercooked chicken could cause illness [2,3]. *Salmonella* spp. is the second-most-common cause of food poisoning after *Campylobacter* spp. It has been found in unpasteurised milk, eggs and raw egg products, meat and poultry. *Salmonella* spp., can be inside of the completely normal-appearing eggs, and if the eggs are eaten raw or undercooked, the bacterium can cause illness [4].

Listeria monocytogens can cause illness in pregnant women, babies and people with reduced immunity and febrile gastroenteritis in healthy people. The estimated annual incidence of listeriosis is quite low and poultry farms are not frequently examined for *L. monocytogens*, but relatively high prevalence of contaminated raw chicken products has been reported. Thus, contaminated farms can be the source for contamination of the slaughter and processing environment [5]. *E. coli* is one of the most common bacteria which cause

diarrhea especially in children. The elderly, infants, and those with impaired immune systems may have a more severe illness. In these patients, the infection may spread from the intestines to the blood stream, and then to other body sites and can cause death unless the person is treated promptly with antibiotics.

Many investigations around the world reported the outbreak, contamination of egg by the *Salmonellas* spp., *C. jejuni*, *L. monocytogens* and *E. coli* [2,6,7]. A couple of reports have confirmed and introduced the egg as dangerous food stuff. So, for the first time we investigated the contamination of the retail eggs by these bacteria in Shahrekord, Iran.

Material and methods

One hundred normal eggs were randomly purchased in the spring and summer of 2008 from small and big supermarkets of Shahrekord, and sent to the food microbiology lab for bacteriological tests. To produce statistically reliable results, the minimum number of eggs was selected based on the number of samples from which the standard error starts converging to an asymptotic value. In the culture of the egg contents, surface of each of the eggs was first disinfected with 70% ethanol. The eggs were broken and the content thoroughly mixed for approximately 1min using a blender.

Each sample was serially diluted with sterile peptone water and presumptive, confirmatory and complementary tests were done for counting and identifying coli form and, *E. coli*. All tests were duplicated and the means recorded. We used selective agar to culture *Salmonella* spp. (Selenite cystein, Xylose Lysine Deoxycholate, Salmonella Shigella agar SS), *C. jejuni* (Skirrow agar, microaerophilic condition) and *L. monocytogens* (Frazer broth, BA-PALCAM; Blood agar with lithium

chloride, polymyxin B sulfate, acriflavine-HCl, and ceftazidime). After 24-48h incubation, standard microbiological techniques including cellular morphology and staining and biochemical (TSI, SIM, IMViC, nalidixic acid resistance, cephalothin sensitivity) tests performed to identify the isolated organism; *E. coli*, *Proteus* spp. and *Klebsiella* spp. [6,8]. All of the culture media were provided by Merck, Germany.

Results and discussion

The result showed that there was no contamination by *Salmonella* spp, *L. monocytogenes*, and *C. jejuni* in the whole content of all 100 eggs purchased in the spring and summer of 2008 from small and big supermarkets of Shahrekord, Iran. In the present study 19 samples (19%) were contaminated by *E. coli*, 4(4%) samples by *Proteus* and one sample (1%) by *Klebsiella* spp. The overall prevalence rate of bacterial contamination of egg was 24 %. Average colony count of coli form bacteria was 20cfu/g and *E. coli* was 12/6cfu/g.

Regarding the increasing consumption of egg and its products, it is necessary to investigate egg contamination. Several factors implicated in egg contamination. The egg shell contamination resulted from deposition of faecal material on the shell, ovarium or oviduct and gut flora, debris material, egg crates, packing and storage, cloths and hands of poultry workers, dust, the environment, weather conditions, transporting and marketing [9]. Among the common contaminant organisms pathogenic to human beings are *Salmonella* spp., *L. monocytogenes*, *C. jejuni* and *E. coli* [7,10]. The isolated bacteria could cause severe health problems like, diarrhea, nausea and abdominal pain since they are pathogenic. The results of this study showed that there was no contamination by *Salmonella* spp.,

L. monocytogenes, and *C. jejuni* in the whole content of all 100 eggs.

Several reports showed that egg was not internally infected with *C. jejuni*. Shane *et al.* [11] reported that hens with faecal shedding *C. jejuni* in farms did not produce infected eggs. Jones investigated the shell and egg content and showed that one shell sample (0.5% of total samples) was *Campylobacter* positive. Two shell samples (1.1% of total samples) were *Salmonella* positive. Twenty-one percent of samples were positive for *Listeria* (33 shells and 5 contents) and no *Salmonella* was found. Other report showed that the *Salmonella* contamination in Spain was 1%, Poland 5%, England 0-7%, and India 1.8% [12].

Fortunately in our study *Salmonella* was not isolated and this suggested that all the study eggs were *Salmonella* free. This may be attributed to the fact that poultry farmers practice strict medication and care. Nineteen (19%) samples were contaminated by *E. coli*, four samples to *Proteus* and one sample to *Klebsiella*. Average colony count of coli form bacteria was 20cfu/g and *E. coli* was 12cfu/g. The mean total of coli form count and mean log were higher than the acceptable limits of 10.00 as set by the International Commission on the Microbiological Specification for Food (ICMSF), showing a hazardous implication on the health of egg consumers [13,14].

Jones *et al.* [15] reported the average *Enterobacteriaceae* less than 0.1 log cfu/ml for the egg contents, with 36.7% of the samples being positive. Cortes *et al.* [7] showed that 45% of eggs were contaminated with *E. coli*. *E. coli* are known to contaminate the surface of egg while mechanical process can spread the bacteria through eggs and meat. Contamination with the pathogen while in the field, occur through improperly decomposed manure, contaminated water

and poor hygienic practices of the farm workers [15,16].

Conclusion

Contamination by *Salmonella* spp. is a major concern in the poultry industry. Fortunately, in our study, the absence of *Salmonella* spp., *L. monocytogenes*, and *C. jejuni* showed the increased notice to hygiene, and hazard analysis critical control point (HACCP) management system have reduced the potential for contamination of these bacteria. However, egg was a source of *E. coli*, *Proteus* and *Klebsiella* contamination. It is important to remember that control is required at all levels in the food chain and by separating cooked and raw. We can help prevent bacterial contamination of table egg from causing egg borne diseases.

Conflict of interest statement: All authors declare that they have no conflict of interest.

Sources of funding: Research Council of the Azad University of Shahrekord, Iran (Grant no. 13787).

References

- 1) Kaneko KI, Hayashidani H, Ohtomo Y, *et al.* Bacterial contamination of ready-to-eat foods and fresh products in retail shops and food factories. *J Food Prot.* 1999; 62: 644-9. PMID: 10382654
- 2) Cox Stern NJ, Wilson JL, Musgrove MT, Buhr RJ, Hiatt KL. Isolation of *Campylobacter* Spp. from semen samples of commercial broiler breeder roosters. *Avian Dis.* 2002; 46: 717-20. PMID: 12243539
- 3) Newell DG, Fearnley C. Sources of *Campylobacter* colonization in broiler chickens. *Appl Envir Microbiol.* 2003; 69: 4343-51. PMID: 12902214
- 4) Davies RH, Breslin M. Investigation of *Salmonella* contamination and disinfection in farm egg-packing plants. *J Appl Microbiol.* 2003; 94: 191-6. PMID: 12534810
- 5) Esteban JI, Oporto B, Aduriz G, Juste RA, Hurtado A. A survey of food-borne pathogens in free-range poultry farms. *Int J Food Microbiol.* 2008; 123: 177-82. PMID: 18234386
- 6) Gorman BS, Adley CC. A study of cross-contamination of food-borne pathogens in the domestic kitchen in the Republic of Ireland. *Int J Food Microbiol.* 2002; 76: 143-50. PMID: 12038571
- 7) Cortés CR, Isaias GT, Cuello CL, Flores JMV, Anderson RC, Campos CE. Bacterial isolation rate from fertile eggs, hatching eggs, and neonatal broilers with yolk sac infection. *Rev Latinoam Microbiol.* 2004; 46: 12-6. PMID: 17061521
- 8) Roberts D, Greenwood M. Practical food microbiology. 3rd ed, UK, Blackwell Publishing Ltd, 2003; 131-91.
- 9) De Reu K, Grijspeerdt K, Messens W, *et al.* Eggshell factors influencing egg shell penetration and whole egg contamination by different bacteria, including *Salmonella Enteritidis*. *Int J Food Microbiol.* 2006; 112: 253-60. PMID: 16822571
- 10) De Reu, Grijspeerdt K, Heyndrickx M, *et al.* Bacterial eggshell contamination in conventional cages, furnished cages and aviary housing systems for laying hens. *Br Poul Sci.* 2005; 46: 149-55. PMID: 15957434
- 11) Shane SM, Gifford DH, Yogasundram K. *Campylobacter jejuni* contamination of eggs. *Vet Res Commun.* 1986; 10: 487-92. PMID: 3798738
- 12) Messens W, Grijspeerdt K, Herman L. Eggshell penetration of hen's eggs by *Salmonella enterica* serovar *Enteritidis* upon various storage conditions. *Br Poult Sci.* 2006; 47: 554-60. PMID: 17050098
- 13) ICMSF. Microorganisms in foods http://www.icmsf.edu/pdf/icmsf2pdf Second edition 175-6.
- 14) Frazier WC, Westhoff DC. Food microbiology. 4th, New York, McGraw Hill, 1988; Table B -1.
- 15) Jones DR, Curtis PA, Anderson KE, Jones FT. Microbial contamination in inoculated shell eggs: II. Effects of layer strain and egg

- storage. *Poult Sci.* 2004; 83: 95-100. PMID: 14761090
- 16) Rahimi E, Momtaz H, Ameri M, Ghasemian-Safaei H, Ali-Kasemi M. Prevalence and antimicrobial resistance of *Campylobacter* species isolated from chicken carcasses during processing in Iran. *Poult Sci.* 2010; 89: 1015-20. PMID: 20371855