

### **Original article**

# A study on risk factors of chronic hepatitis B carriers

Mahmoud Jahangirnezhad, MD<sup>1</sup> Eskandar Hajiani, MD<sup>2</sup> Manochehr Makvandi, PhD<sup>3</sup> Fariba Jalali, BSc<sup>2</sup>

<sup>1</sup>Department of Periodontics, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran <sup>2</sup>Department of Internal Medicine and Gastroenterology, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran <sup>3</sup>Department of Virology, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

#### Address for correspondence:

Dr. Mahmoud Jahangirnezhad, Department of Periodontics, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran Tel: 0098611 3205169 Fax : 0098611 3364366 E-mail:drmahjahan@yahoo.com

#### How to cite this article:

Jahangirnezhad M, Hajiani E, Makvandi M, Jalali F. A study on risk factors of chronic hepatitis B carriers. Jundishapur J Microbiol. 2011; 4(4): 267-272.

Received: August 2010 Accepted: January 2011

#### Abstract

**Introduction and objective:** Hepatitis B is a disease of public health importance in Iran. The aim of this study was to evaluate some possible risk factors for the spread of hepatitis B infection from the hepatitis B virus (HBV) carriers.

**Materials and methods:** A cross-sectional study was conducted among chronic HBV individuals who referred to gastrointestinal department, Imam Khomeini hospital, Ahvaz Jundishapur University of medical sciences from October 2009 to June 2010. All subjects were evaluated using a face-to-face questionnaire about demographic aspects. The analysis included data on past medical history, physical examination and periodic evaluation clinically and serologically. The control group consisted of the patients referred to the gastrointestinal clinics with negative HBV serologic markers of HBV infection. The risk factors among infected subjects (HBV-positive) were compared to those of subjects never exposed (HBV-negative) to HBV.

**Results:** A total of 560 subjects were studied for HBV, of which 272 were HBV-positive and 288 HBV negative cases comprised the control group. Mean age of the patients was  $35\pm9$  years. HBV-positive subjects were more likely to be of female gender (61.39% versus 41.31%, P<0.0001). Endoscopy 54.77%, major surgery 44.48%, and tattooing history 8.45% were found to be independent risk factors of being chronically infected with hepatitis B virus.

**Conclusion:** Our data indicate that a history of endoscopy, major surgery and tattooing are important risk factors for spreading of HBV infection.

**Significance and impact of the study:** Improvements in certain lifestyle patterns and customs in this area may be essential to prevent transmission of the infection.

Keywords: Hepatitis B virus; Risk factors; Seroprevalence



# Introduction

Hepatitis B virus (HBV) infection is a health problem worldwide. with approximately 400 million chronically infected people. HBV is an important agent of acute and chronic hepatitis and the endemicity of a given population is determined by detecting hepatitis B surface antigen (HBsAg) in serum and is classified as low, intermediate, and high endemicity according to the percentage of infection (2%, 2-7.9%, and >10%, respectively) [1,2]. HBV has a circular and partially double stranded DNA genome of 3.2Kb containing four overlapping open reading frames.

HBV strains isolated worldwide have been classified into six genomic groups deduced from genome comparisons and designated genotypes A to F [3,4,]. Chronic, persistent HBV infection develops at the end-stage of liver disease, such as liver cirrhosis and hepatocellular carcinoma [5,6]. Although safe and effective HBV vaccine is available but new cases of HBV infection are still reported throughout the world [7]. Recently occult HBV infections were also reported on which the serological HBsAg is negative in the patient sera but the HBV DNA could be detected by PCR [8].

One of the important priorities for elimination and control of HBV is to know the factors involved in HBV transmission especially in the endemic regions. Iran is the intermediate for the HBV infection [9]. The prevalence of HBV infection among the blood donors have been reported in Iran [10]. It is evident that 70-84% of cirrhotic patients and 72% of individuals with hepatocellular carcinoma in Iran have evidence of exposure to HBV [11]. The prevalence of HBV among the injecting drug user in Ahvaz was 44.3% [12].

The prevalence of HBV infection among the hemodialysis patients in Ahvaz was reported to be 5.1% [13]. Thus important measures should be implemented to the risk factors causing spread of HBV infection. The aim of this study was to evaluate some possible risk factors for the spread of hepatitis B infection by HBV carriers.

# Materials and methods

A cross-sectional study was conducted 272 HBV carriers including among 167(61.39%) female and 105(38.6%) male who referred to gastrointestinal department, Imam Khomeini hospital, Ahvaz Jundishapur University of medical sciences from October 2009 to June 2010. The evidence of HBV infection among the HBV carriers was tested for HBsAg by ELISA test (Behring, Germany).

All the HBV carriers having the following inclusion criteria: documented case of HBsAg positive for more than six months, HBV carriers without clinical acute symptoms, and the HBV carriers with normal liver function test of alanine transaminase and aspartate transaminase. The excluding criteria were individual with acute HBV symptoms, individual with abnormal normal liver function test of alanine transaminase and aspartate transaminase, and individual refusing to answer the questions. While 288 subjects 119(41.31%) including female and 169(58.68%) male as a control group were found negative for HBsAg.

All the HBV patients and control group were registered at the department of gastroenterology of the Imam Khomeini. The HBV positive carriers; age was between 14 to 70 years old with the mean age of 35±9 years while the age of the control group was between 19-65 years with the mean age of  $39\pm8$  years. The independent risk factors including tattooing, history endoscopy. major surgery. phlebotomy (Hejamat), intravenous drug users, dentist visiting, were determined by

Jundishapur Journal of Microbiology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran, Tel: +98611 3738283; Fax: +98611 3362537; URL: http://jjm.ajums.ac.ir; Email: editorial office: jjm@ajums.ac.ir JJM. (2011); 4(4): 267-272.



using a questionnaire. The questioners were general practitioners, blinded to the HBsAg test results of the subjects before and during the interview. The results were analyzed

Chi-squared analysis performed to determine whether significance of risk factor and/or also presented to indicate the chance of risk factor on HBsAg positive in contrast to HBsAg negative. The P value below 0.05 was considered to be significant. to the control group 119(41.31%) was found to be significant (P<0.0001). The independent risk factors among the HBV carriers included; tattooing history, 23(8.45%), endoscopy 149(54.77%), major surgery 121(44.48%) and high risk job 14 (5.14%). Therefore, among the control group the risk factors included tattooing history 1(0.34%), endoscopy 4(1.4%) and major surgery 2(0.7%). Regarding the above data, most of independent risk factors were found to be significant (table 1).

#### Results

The results of this study revealed that the ratio of female HBV carriers 167(61.39%)

**Table 1:** The odds ratio for each independent variable among the HBV carriers and control group (the p value in the majority of the variable found significant)

Risk factor	HBsAg positive	HBsAg negative	Odds ration	Р
	(n=272)	(n=288)	(95% CI)	value
Sex female	167(61.39%)	119(41.31%)	2.26(1.6-3.177)	0.0001
Tattooing history,	23(8.45%)	1(0.34%)	26.5 (3.55-197.7)	0.0001
Endoscopy	149(54.77%)	4(1.38%)	86(31.16-237.4)	0.0001
Major surgery	121(44.48%)	2(0.69%)	114.6(27.944-469.8)	0.0001
Phlebotomy	1(0.36%)	0	2.12(.07-63.6)	0.65
IDU©	3(1.1%)	1(0.34%)	3.2(.3-30.95)	0.28
High risk job®	14(5.14%)	0	31.25(1.85-527.46)	0.0001
Dentist visit	205(75.36%)	200(69.44%)	1.35(0.92-1.95)	0.12

©IDU, Intravenous Drug Use, ®High risk job included nurse and barber

#### Discussion

Persistent or chronic HBV infection is one of the most common persistent viral infections in human. The results of this study revealed that the ratio of female HBV carriers167 (61.39%) to male carriers119 (41%). was found to be significant (P<0.0001).

The horizontal transmission of HBV infection occurred mainly by exposure to blood transfusion [14], sexual depravity [15], tattooing[16], endoscopy [17], surgical procedure [18], needle sticks [19], hemodialysis [20], dental procedure [21], intravenous drug use (IDU) [22], history of sexually transmitted disease (STD) [23]. The prevalence of carriers HBsAg carriers

in Iran varied from 1.07% in Fars province [24] to over 6% in the Balouchestan province [25]. Although sexual transmission of the HBV is the important key and has been reported [13,26], in Iranian culture due to social values which is strongly noticeable and might reduce the rout of sexual transmission. Thus the role of STD in Iran is not predictable.

Sex especially among the married couple may play a significant role in the transmission of HBV infection [27]. The transmission of HBV infection from the infected mother to newborn infant has been reported [28]. In our study about 61.39% of HBV positive were married female which may have crucial role in spreading of HBV

Jundishapur Journal of Microbiology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran, Tel: +98611 3738283; Fax: +98611 3362537; URL: http://jjm.ajums.ac.ir; Email: editorial office: jjm@ajums.ac.ir JJM. (2011); 4(4): 267-272.



infection. The health care workers (HCW) are at a higher risk of HBV infection than general population because they are more frequently exposed to infected patients [29,30]. In the present study 5.14% of positive HBV carriers were nurses, whose jobs were highly risky.

There are some reports on infected physicians, dentists and oral surgeons which have been implicated in transmission of HBV infection due to multiple contacts with patients [31,32]. Tattooing is another risk factor for the HBV infection. It has been found that the tattooing could be a significant factor to be considered in relation to the transmission of hepatitis B and tattooing [33]. In our data 8.45% of the patients had history of tattooing. Vaccination against hepatitis B is an important factor for the prevention of HBV infection [34].

It has been revealed that the are more than 30 independent risk factors including tooth brushes, baby bottles, toys, eating utensils, razors, blood feeding insects, heamodialysis, breast feeding, acupuncture` which are implicated in the transmission of HBV infection [35] but these risk factors were not considered in the present study.

## Conclusion

Our data indicate that a history of endoscopy and major surgery, tattooing are important risk factors for HBV infection in our area and that more careful screening of blood for anti-HBV must be introduced in our clinical practice. Improvements in certain lifestyle patterns and customs in this area may be essential to prevent transmission of the infection.

## Acknowledgements

Authors would like to thank B Eftekhar, A. Jahangirnezhad, and M. Jahangiri.

**Conflict of interest statement:** All authors declare that they have no conflict of interest.

**Sources of funding:** Vice Chancellor for Research affairs of Ahvaz Jundishapur University of Medical Sciences (Grant no. U-88091).

### References

- Hou J, Liu Z, Gu F. Epidemiology and prevention of hepatitis B virus infection. *Int J Med Sci.* 2005; 2: 50-7. PMID: 15968340
- Gish RG, Gadano AC. Chronic hepatitis B: current epidemiology in the Americas and implications for management. *J Viral Hepat*. 2006; 13: 787-98. PMID: 17109678
- Magnius LO, Norder H. Subtypes, genotypes and molecular epidemiology of the hepatitis B virus as reflected by sequence variability of the S-gene. *Intervirology*. 1995; 38: 24-34. PMID: 8666521
- Okamoto H, Tsuda F, Sakugawa H, *et al.* Typing hepatitis B virus by homology in nucleotide sequence: comparison of surface antigen subtypes. *J Gen Virol.* 1988; 69: 2575-83. PMID: 3171552
- Bruix J, Llovet JM. Hepatitis B virus and hepatocellular carcinoma. *J Hepatol*. 2003; 39(Suppl. 1): S59-S63. PMID: 14708679
- Zoulim F. Detection of hepatitis B virus resistance to antivirals. *J Clin Virol* 2001; 21: 243-53. PMID: 11397661
- Lavanchy D. Hepatitis B virus epidemiology, disease burden, treatment, and current and emerging prevention and control measures. *J Viral Hepat.* 2004; 11: 97-107. PMID: 14996343
- Adachi S, Shibuya A, Miura Y, Takeuchi A, Nakazawa T, Saigenji K. Impact of occult hepatitis B virus infection and prior hepatitis B virus infection on development of hepatocellular carcinoma in patients with liver cirrhosis due to hepatitis C virus. *Scand J Gastroenterol.* 2008; 43: 849-5. PMID: 18584524
- 9) Alavian SM, Mostajabi P, Malekzadeh R. Evaluation of hepatitis B transmission risk

Jundishapur Journal of Microbiology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran, Tel: +98611 3738283; Fax: +98611 3362537; URL: http://jjm.ajums.ac.ir; Email: editorial office: jjm@ajums.ac.ir JJM. (2011); 4(4): 267-272.



factors in Tehran blood donors. *Govaresh*. 2005; 39: 26-36.

- 10) Lankarani KB, Nabipoor I. Reassessment of the role of hepatitis B and C viruses in postnecrotic cirrhosis and chronic hepatitis in southern Iran. *Irn J Med Sci* 1999; 24: 117-21.
- 11) Taghavi S, Sedighi S, Mehrabani D, Khademolhosseini F. Hepatitis D in chronic active hepatitis B: prevalence, liver enzymes and histopathology-an epidemiological study in Shiraz, southern Iran, 2003-2004. *Hepat Mon.* 2008; 4: 248-51.
- 12) Alavi SM, Etemadi A. HIV/HBV, HIV/HCV and HIV/HTLV-1 co infection among injecting drug user patients hospitalized at the infectious disease ward of a training hospital in Iran. *Pak J Med Sci*. 2007; 23: 510-13.
- 13) Assarehzadegan MA, Shakerinejad G, Noozkohnejad R, Amini A, Rahim Rezaee SA. Prevalence of hepatitis C and B infection and HC V genotypes among hemodialysis patients in Khuzestan province, southwest Iran. Saudi J Kidney Dis Transpl. 2009; 20: 681-4. PMID: 19587521
- 14) Buseri FI, Muhibi MA, Jeremiah ZA, Buseri FI, Muhibi MA, Jeremiah ZA. Seroepidemiology of transfusion-transmissible infectious diseases among blood donors in Osogbo, south-west Nigeria. *Blood Transfus*. 2009; 7: 293-9. PMID: 20011640
- 15) Rich JD, Anderson BJ, Schwartzapfel B, Stein MD. Sexual risk for hepatitis B virus among hepatitis C virus-negative heroin and cocaine users. *Epidemiol Infect.* 2006; 143: 478-84. PMID: 16194294
- 16) Samuel MC, Doherty PM, Bulterys M, Jenison SA. Association between heroin use, needle sharing and tattoos received in prison with hepatitis B and C positivity among street-recruited injecting drug users in New Mexico, USA. *Epidemiol Infect.* 2001; 127: 475-84. PMID: 11811881
- 17) Morris IM, Cattle DS, Smits BJ. Endoscopy and transmission of hepatitis B. *Lancet*. 1975; 2: 1152. PMID: 53640
- 18) Spijkerman IJ, van Doorn LJ, Janssen MH *et al.* Transmission of hepatitis B virus from

a surgeon to his patients during high-risk and low-risk surgical procedures during 4 years. *Infect Control Hosp Epidemiol*. 2002; 23: 306-12. PMID: 12083233

- 19) Gerberding JL. Management of occupational exposures to blood-borne viruses. *N Engl J Med.* 1995; 332: 444-51. PMID: 7824017
- 20) Siagris D, Christofidou M, Triga K, et al. Occult hepatitis B virus infection in hemodialysis patients with chronic HCV infection. J Nephrol. 2006; 19: 327-33. PMID: 16874693
- 21) Nagao Y, Matsuoka H, Kawaguchi T, Ide T, Sata M. HBV and HCV infection in Japanese dental care workers. *Int J Mol Med.* 2008; 21: 791-9. PMID: 18506374
- 22) Lum PJ, Hahn JA, Shafer KP, *et al.* Hepatitis B virus infection and immunization status in a new generation of injection drug users in San Francisco. *J Viral Hepat.* 2008; 15: 229-36. PMID: 18233994
- 23) Trepka MJ, Weisbord JS, Zhang G, Brewer T. Hepatitis B virus infection risk factors and immunity among sexually transmitted disease clinic clients. *Sex Transm Dis.* 2003; 30: 914-8. PMID: 14646641
- 24) Ghavanini AA, Sabri MR. Hepatitis B surface antigen and anti-hepatitis C antibodies among blood donors in the Islamic republic of Iran. *East Mediterr Health J.* 2000; 6: 1114-6. PMID: 12197336.
- 25) Merat S, Malekzadeh R, Rezvan H, Khatibian M. Hepatitis B in Iran. *Arch Iran Med.* 2000; 3: 192-201.
- 26) Shethwala ND, Mulla SA, Kosambiya JK, Desai VK. Sexually transmitted infections and reproductive tract infections in female sex workers. *Indian J Pathol Microbiol.* 2009; 52: 198-9. PMID: 19332911
- 27) Wang Cs, Chang TT, Yao WJ, Chou P. Comparison of hepatitis B virus and hepatitis C virus prevalence and risk factors in a community-based study. *Am J Trop Med Hyp.* 2002; 66: 389-93. PMID: 12164293
- 28) Tan HH, Lui HF, Chow WC. Chronic hepatitis B virus (HBV) infection in

Jundishapur Journal of Microbiology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran, Tel: +98611 3738283; Fax: +98611 3362537; URL: http://jjm.ajums.ac.ir; Email: editorial office: jjm@ajums.ac.ir **JJM. (2011); 4(4): 267-272.** 



pregnancy. *Hepatol Int.* 2008; 2: 370-5. PMID: 19669267

- 29) Gunson RN, Shouval D, Roggendorf M, et al. Hepatitis B virus (HBV) and hepatitis C virus (HCV) infections in health care workers (HCWs): guidelines for prevention of transmission of HBV and HCV from HCW to patients. J Clin Virol. 2003; 27: 213-30. PMID:12878084
- 30) Daha TJ, Bilkert-Mooiman AJM, Ballemans C, et al. Hepatitis B virus infected health care workers in the Netherlands, 2000-2008. Eur J Clin Microbiol Infect Dis. 2009; 28: 1041-4. PMID: 19350292
- 31) Thurn J, Willenbring K, Crossley K. Needlestick injuries and needle disposal in Minnesota physicians' offices. *Am J Med.* 1989; 86: 575-9. PMID: 2712064
- 32) Ammon A, Reichart PA, Pauli G, Petersen LR. Hepatitis B and C among Berlin dental

personnel: incidence, risk factors, and effectiveness of barrier prevention measures. *Epidemiol Infect*. 2000; 125: 407-13. PMID: 11117965

- 33) Limentani AE, Elliott LM, Noah ND, Lamborn, JK. An outbreak of hepatitis B from tattooing. *Lancet*. 1979; 2: 86-8.
  PMID: 87977
- 34) Batista SM, Andreasi MS, Borges AM, et al. Seropositivity for hepatitis B virus, vaccination coverage, and vaccine response in dentists from Campo Grande, Mato Grosso do Sul, Brazil. Mem Inst Oswaldo Cruz. 2006; 101: 263-7. PMID: 16862319
- 35) Seeger C, Mason W, Zoulim F. Hepadnaviruses. In: Knipe DM, Howley PM, (eds), *Fields virology*, Philadelphia, Lippincott Williams & Wilkins. 2007; 2977-3029.