Jundishapur J Microbiol. 2013;6(1): 42-46. DOI: 10.5812/jjm.4756



Studies on Drug Sensitivity and Bacterial Prevalence of UTI in Tribal Population of Paschim Medinipur, West Bengal, India

Suman Kumar Maji¹, Chiranjit Maity¹, Suman Kumar Halder¹, Tanmay Paul¹, Pratip Kumar Kundu², Keshab Chandra Mondal^{1*}

¹ Department of Microbiology, Vidyasagar University, Midnapore, West Bengal, India

² Department of Microbiology, Culcutta School of Tropical Medicine, Kolkata, West Bengal, India

ARTICLE INFO ABSTRACT Background: There is paucity of information in regards to prevalence of bacteria in uri-Article type: nary tract infection (UTI) in tribal population. Original Article **Objectives:** This study was conducted to observe dominant organisms causing UTI in tribal community of Medinipur, West Bengal, India and evaluate their sensitivity to com-Article history: Received: 07 Mar 2012 mon antibiotics. Patients and Methods: A total number of 4,416 urine samples of clinically suspected Revised: 15 Apr 2012 UTI attending outpatient department and hospitalized tribal patients of West Medini-Accepted: 29 Apr 2012 pur zone were collected. Urine samples were inoculated on Blood agar, MacConkey's agar, and cystine lactose electrolyte deficient (CLED) agar and incubated at 37°C for 24 Keywords: hr and extended up to 48 hr in cases of negative growth. Antibiotic susceptibility testing Urinary Tract Infection Bacteria against the most common causative bacteria was also performed. **Results:** A total number of 1,190 samples revealed positive bacterial growth. Seven types Antibiotics Sensitivity of dominant organisms were isolated as causative agents like Escherichia coli (63.44%), Klebsiella sp. (14.62%), Pseudomonas aeruginosa (4.53%), Proteus sp. (4.62%), other Gram negative bacteria (5.79%), Staphylococcus aureus and coagulase negative Streptococci (5.21%), and other Enterococcus sp. (1.76%). E.coli was the most common pathogen of UTI in tribal population with female susceptibility predominantly more than male patients. Among16 antibiotics, amikacin (14.30%) and meropenem (12.0%) were shown to be the most sensitive to and Tobramycin (0.66%) the least effective on E.coli. Conclusions: Females are more susceptible to UTI than males and the most effective drug (sensitive against isolated E. coli) is Amikacin. Therefore, culture and antimicrobial drug sensitivity testing are essential procedures for proper management of UTI in tribal population. Published by Kowsar Corp, 2013. cc 3.0.

▶ Implication for health policy/practice/research/medical education:

The result will be very much helpful for medical practitioners and clinicians of the region to create a common database for UTI causing microorganisms and their antibiogram profile.

Please cite this paper as:

Maji SK, Maity C, Halder SK, Paul T, Kundu PK, Mondal KC. Studies on Drug Sensitivity and Bacterial Prevalence of UTI in Tribal Population of Paschim Medinipur, West Bengal, India. *Jundishapur J Microbiol*. 2013;**6**(1):42-6. DOI: 10.5812/jjm.4756

DOI:10.5812/jjm.4756

^{*} Corresponding author: Keshab Chandra Mondal, Department of Microbiology, Vidyasagar University, Midnapore 721-102, West Bengal, India. Tel: +91-9434196483, Fax: +91-3222275329, E-mail: mondalkc@gmail.com

^{© 2013} Ahvaz Jundishapur University of Medical Sciences; Published by Kowsar Corp.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Background

Bacterial urinary tract infection (UTI) is the most common kind of infection affecting the urinary tract and causing inflammation of bladder and kidneys. Urine is a favorable medium for growth of bacteria due to its enriched chemical composition (1-3). UTI is the third most common cause of admission to hospitals in India. It has been estimated that about 6 million patients per year are visited worldwide for UTI out of which around 30,000 are treated in the wards. (4). Particularly, those sub-populations who are at increased risk of UTI include infants, pregnant women, and elderly of both sexes, as well as those with spinal cord injuries, indwelling catheters, diabetes, multiple sclerosis, immunodeficiency, and underlying urological abnormalities.

Majority of women are recurrently infected within one year (5). Compared to males, young sexually active females and teenage students girls are mostly infected by UTI (10.57% higher). In this group, most prominent pathogens are Escherichia coli (32.8%), Klebsiella pneumoniae (22.4%), and Staphylococcus aureus (15.1%) (6). E. coli cause 75 - 90% of uncomplicated UTIs (7) whereas S. saprophyticus frequently causes UTI in younger women at estimated rate of 5 - 15%(8). Golechha and Solanki (9) performed a bacteriological study on preoperative urine samples of 100 urolithiasis cases in India. E.coli was encountered as the commonest pathogen recovered from pre-operative urine and stone cultures (32.25% and 21.73%, respectively) followed by Pseudomonas aeruginosa (22.58% and 17.39%, respectively). Enterococcus and other Gram negative rods other than E.coli have also been implicated in some cases (10).

UTI treatment incurs a considerable cost, both directly and indirectly, on the health care system. In the United States of America, approximately 5% of all patients acquired infections after admission to hospital with an estimated cost of \$1.6 million annually (11-13). The predominant urinary tract pathogens are *E.coli* (93%); other minor groups include *Enterococcus sp.*, hemolytic *Streptococci sp.*, *S. aureus*, and *Pseudomonas sp.*

Antimicrobial susceptibility profile in respect to causative microbes may significantly reduce morbidity and mortality, cost of treatment, and duration of hospitalization if duly provided to medical practitioners and clinicians in a rapid and timely fashion (14).

2. Objectives

The West Medinipur is a habitat of diverse group of populations including primitive tribal like Lodha, Kheria / Sabar, Munda, Santal, Kohl, Oraon, Mahali, and Bhumij. Because of poor health care system, most of clinicians prescribe antibiotics to treat UTI blindly, resulting in failure of treatment in many cases due to occurrence of bacterial drug resistance.. The major objectives of this study include isolation and identification of predominant UTI causing bacteria in different groups of tribal people and observation of the current sensitivity trend of most common commercial drugs against predominant pathogens. This may also highlight the prevalence of pathogens in tribal population of different age and sex.

3. Patients and Methods

3.1. Study Design

Urine samples of clinically suspected outpatients with UTI as well as hospitalized patients were collected. A detailed history was taken and complete clinical examination was carried out for each case of UTI. Some samples were collected from clean catch midstream: in neonatal cases the samples were collected through suprapubic approach and in children (less than 3 years) by sterile urine bags (any colony count of bacterial growth was significant). It was noticed to discontinue all antibiotics 72 hr prior to urine collection for culture and sensitivity. Urine samples were delivered to the laboratory within 1hr and processed within 24 hr from collection. Each and every sample was observed under microscope and examined for the presence of puss cells, RBCs, epithelial cells, casts, and crystals. A total number of 4,416 early morning midstream urine samples of patients from different areas were collected from June 2006 to October 2007.

3.2. Isolation and Identification

A standard loop technique was used to place 0.01 mL of urine for inoculation on Blood agar, MacConkey's agar, and cystine lactose electrolyte deficient (CLED) agar, and incubated at 37°C for 24 hr and extended up to 48 hr in cases of negative growth. At any multiple growths, the culture was repeated before acceptance of outcomes. All positive samples were rechecked by collecting second urine samples to rule out contaminations. The number of colonies was counted to quantify organisms. Diagnosis of urinary tract infection for a single pathogen was defined based on significant colony count of $\geq 10^5$ CFU/ mL for Gram negative and $\geq 10^4$ CFU/mL for Gram positive bacteria. The organisms were identified through monitoring general biochemical tests such as catalase, oxidase, Triple Sugar Iron agar (TSI), citrate utilization (Simmon's citrates medium), urease (Christensen's Urea Agar), indole, motility, H₂S production (Sulphide Indole Motility Medium), esculin hydrolysis (Bile esculin agar), and sugar fermentation tests. All culture media were provided by Himedia Laboratories Pvt. Ltd., India.

3.3. Antimicrobial Susceptibility of Most Common Bacteria of UTI

Antibiotic susceptibility testing against the most common causative bacteria was performed according to Kirby Bauer's method (12). The antibiotic discs (Himedia,

www.SID ir

India) (each 6.3 mm diameter) used were Amoxicillin with clavulanic acid (30mcg), Ceftazidime(30mcg), Cefoperazone (75mcg), Cefotaxime (30mcg), Cifixime (5mcg), Ceftriaxone (30mcg), Tobramycin (10mcg), Gentamycin (10mcg), Netilmycin (30mcg), Amikacin (30mcg), Cotrimoxazole (25mcg), Gatifloxacin (5mcg), Norfloxacin (10mcg), Levofloxacin (5cg), Ofloxacin (5mcg), and Meropenem (10mcg). The zone of inhibition of each antibiotics against the causative bacteria were compared using standard CLSI protocol (15).

4. Results

To find out predominant causative organism of UTI in our district, 4,416 urine samples (both community and hospital acquired) were cultured from different sexes and ages of tribal population.

In 1,596 out of 4,416 samples, bacterial growth was observed as a cause of significant and insignificant urinary tract infection. Among them 1,190 samples showed significant bacteriuria, 383 samples showed insignificant bacteriuria, and 23 samples showed colony count between 10⁴–10⁵ CFU/mL of mixed growth which was considered as contaminated samples (16-18). Urine samples with negative growth were marked in 2,820 cases.

Generally, we had identified seven bacteria sp. mainly *E.coli, Klebsiella sp., P. aeruginosa, Proteus sp., S. aureus* and coagulase negative streptococci (CONS), *Enterococcus sp.,* and others Gram negative spp which consists of *Alkaligenes feacalis, Acinetobacter bawmani, Providentia sp.,* and some *Enterobacter spp.*

In our study it was observed that *E. coli* was the most common causative organism for UTI in both community and hospital acquired infection (63.44%). *Klebsiella sp.* was the second most common isolated organisms (14.62%) in all ages of this population. The infection rate of *P. aeruginosa* and *Proteus sp* (both *P. vulgaris* and *P. mirabilis*) were 4.53% and 4.62%, respectively in respect to total number of positive growth samples.

Gram positive bacteria including *Staphylococcus sp.* and *Enterococcus sp.* were responsible for UTI in both genders. In our West Medinipur zone from tribal community, the rate of positive samples for *S. aureus &* CONS, and *Enterococcus sp* were 5.21% and 1.76%, respectively. Infection rate by other unidentified species was 5.79% (*Figure 1*).



In respect of sex variation, isolated bacteria were sex-dependent. The percentages of *Pseudomonas sp.* and *Proteus sp.* were slightly higher in male patients, whereas females were infected more by *E.coli* and *Klebsiella sp.* (*Figure 2*).



Figure 2. Occurrence of Different Isolated Bacteria (in Number and Percentage) Among Males and Females

Antibiotic susceptibility tests were performed for common isolated pathogen *E.coli*. 755 out of 1,190 positive growth, showed positive growth of *E. coli*. All of these isolates were processed individually to pass sensitivity test against 16 different antibiotics. *Figure* 3 revealed antibiotic susceptibility pattern of different isolated *E.coli*, in which Amikacin (14.30%) was the most common effective drug followed by Meropenem (12.45%). One aminoglycoside, Tobramycin, was the least effective drug on *E.coli* in UTI.



Figure 1. Percentage (%) of the Common Bacteria Isolated From Urine Samples (1,190)



5. Discussion

Asymptomatic bacteriuria is the major cause of UTI because under favorable conditions colonized bacteria in urinary tract may ascend towards the bladder and cause cystitis which is usually associated with the classic symptoms of UTI (i.e., pain, frequency, and urgency). If UTI remains untreated, it can proceed via ureters, to the kidneys that may cause pyelonephritis which may lead to irreversible kidney damage, renal failure, and death (16, 17).

The tribal people have different culture, custom, and life style and most of them are dependent upon natural medicine. Due to poor economic condition, they are not able to avail the facility of modern medicinal system. This is probably the first study in this group of people to understand prevalent organism for UTI and related antibiotic sensitivity. In this study, we found that percentage of urinary tract infection with significant bacteriuria was 74.56% among 1,596 positive growth samples with suspected fever with or without UTI symptoms, randomly. A study was conducted among 1,000 pregnant women with asymptomatic UTI from National Medical College; Calcutta that revealed the prevalence of bacteriuria was 10.2% (16). In this study a comparable different pattern of bacterial prevalence was found among UTI infected tribal people such as E.coli (63.44%), Klebsiella sp. (14.62%), P. aeruginosa (4.53%), Proteus sp. (4.62%), S. aureus and CONS (5.21%), Enterococcus sp. (1.76%) and other Gram negative spp. including Alkaligenes feacalis, A. bawmani, Providentia *sp.*, and some *Enterobacter* (1.76%).

E.coli was the commonest bacteria (63.44%) in UTI patients, but with a different rate obtained from other populations in U.S.A. study (75.5% - 87.0%) (19, 20), where the number are 68.69% and 83.0% as seen by Rayan *et al.* in general population in India (21). On the contrary, in the report of Nabeela *et al.*, *Klebsiella sp.* and *Proteus sp.* were responsible for 16% and 11% of all urinary tract infections, respectively (22). This variation may be attributed to different life style, poor healthcare system, lack of education, and inadequate availability of water, and also may be due to geographical variations.

However, it should be pointed out that other investigators have observed *E. coli* as the common bacterium in significant bacteriuria. In our study same findings were obtained in tribal people of West Medinipur zone and enriched the fact that females are more susceptible to UTI than males and that the most effective drug (effective on isolated *E.coli*) is Amikacin (from amino-glycosides group) for this tribal people. Therefore, culture and antimicrobial drug sensitivity testing are prepared for surveillance purposes to guide clinicians on proper management and prevent empirical treatment of tribal population with both asymptomatic & symptomatic bacteriuria.

Our study will be a common and convenient database for medical practitioners for treatment of UTI patients in tribal

population of Paschim Medinipur, West Bengal, India.

Acknowledgements

The authors are thankful to the Director, Midnapore Scan Centre Private Ltd. and Department of Microbiology, Vidyasagar University, Midnapur for providing laboratory facilities to carry out the research.

Financial Disclosure

There is no financial conflict among the authors or with the Institute for the carrying out the particular study.

Funding/Support

We are thankful to Defense Institute of Physiology & Allied Sciences (DIPAS) as well as Council of Scientific & Industrial Research (CSIR), India, for their financial supports to carry out this work.

Authors' Contribution

None declared.

References

- Asscher AW, Sussman M, Waters WE, Davis RH, Chick S. Urine as a medium for bacterial growth. *Lancet*. 1966;288(7472):1037-41.
- Asscher AW, Sussman M, Weisser R. Bacterial growth in human urine. In, "Urinary Tract Infections". Edited by Francis O'Grady and Brumfitt W. Oxford University Press, London, New York & Toronto. 1968:3-14.
- Chernew I, Braude AI. Depression of phagocytosis by solutes in concentrations found in the kidney and urine. J Clin Invest. 1962;41:1945-53.
- Bano K, Khan J, Rifat, Begum H, Munir S, Akbar NU, et al. Patterns of antibiotic sensitivity of bacterial pathogens among urinary tract infections (UTI) patients in a Pakistani population. Afr J Microbiol Res. 2012;6(2):414-20.
- Koljalg S, Truusalu K, Vainumae I, Stsepetova J, Sepp E, Mikelsaar M. Persistence of Escherichia coli clones and phenotypic and genotypic antibiotic resistance in recurrent urinary tract infections in childhood. J Clin Microbiol. 2009;47(1):99-105.
- Kumar CS, Jairam A, Chetan S, Sudesh P, Kapur I. Asymptomatic bacteriuria in school going children. *Indian J Med Microbiol*. 2002;20(1):29-32.
- Ejrnaes K, Sandvang D, Lundgren B, Ferry S, Holm S, Monsen T, et al. Pulsed-Field Gel Electrophoresis Typing of Escherichia coli Strains from Samples Collected before and after Pivmecillinam or Placebo Treatment of Uncomplicated Community-Acquired Urinary Tract Infection in Women. J Clin Microbiol. 2006;44(5):1776-81.
- 8. Widerstrom M, Wistrom J, Ferry S, Karlsson C, Monsen T. Molecular epidemiology of Staphylococcus saprophyticus isolated from women with uncomplicated community-acquired urinary tract infection. *J Clin Microbiol*. 2007;**45**(5):1561-4.
- Solanki A, Golechha S. Bacteriology and chemical composition of renal calculi accompanying urinary tract infection. 2001.
- Darbro BW, Petroelje BK, Doern GV. Lactobacillus delbrueckii as the Cause of Urinary Tract Infection. J Clin Microbiol. 2009;47(1):275-7.
- Cox CE. Nosocomial urinary tract infections. Urology. 1988;32(3):210-5.
- Hua HT, Bollet C, Tercian S, Drancourt M, Raoult D. Aeromonas popoffii urinary tract infection. J Clin Microbiol. 2004;42(11):5427-8.



- Meares EM, Jr. Current patterns in nosocomial urinary tract infections. Urology. 1991;37(3 Suppl):9-12.
- Ataee RA, Tavana AM, Hosseini SMJ, Moridi K, Zadegan MG. A Method for Antibiotic Susceptibility Testing: Applicable and Accurate. Jundishapur J Microbiol. 2011;5(1):341-5.
- Clinical, Institute LS. Performance standards for antimicrobial susceptibility testing, twenty first informational supplement M100-S21. CLSI Wayne, PA; 2011.
- Ghosh AK, Dey NC, Guha AC. Significance of bacteriuria in pregnancy. J Indian Med Assoc. 1975;64(3):63-8.
- Ngwai YB, Iliyasu H, Young E, Owuna G. Bacteriuria and Antimicrobial Susceptibility of Escherichia coli Isolated From urine of Asymptomatic University Students in Keffi, Nigeria. Jundishapur J Microbiol. 2012;5(1):323-7.
- 18. Nolte F, Metchock B, Murray P, Baron E, Pfaller M, Tenover F, et al.

Manual of clinical microbiology. Manual of clinical microbiology. 1995.

- Ghedira Besbes L, Messaoudi A, Ben Meriem C, Guediche MN. [Profile of antimicrobial resistance of agents causing urinary tract infections in children]. *Tunis Med.* 2004;82(3):299-305.
- 20. Mangiarotti P, Pizzini C, Fanos V. Antibiotic prophylaxis in children with relapsing urinary tract infections: review. *J Chemother*. 2000;**12**(2):115-23.
- 21. Rayan AJ, Bai KT, Unnisa A, Lalitha K. Drug sensitivity of urinary pathogenic bacteria. *J Indian Med Assoc*. 1978;**70**(11):245-7.
- 22. Noor N, Ajaz M, Rasool SA, Pirzada ZA. Urinary tract infections associated with multidrug resistant enteric bacilli: characterization and genetical studies. *Pak J Pharm Sci.* 2004;**17**(2):115-23.