Molecular Characterization of Vibrio cholerae Isolated From Clinical Samples in Kurdistan Province, Iran

Rashid Ramazanzadeh ^{1,2}; Samaneh Rouhi ³; Pegah Shakib ³; Babak Shahbazi ^{2,*}; Farzam Bidarpour⁴; Mohammad Karimi⁴

¹Cellular and Molecular Research Center, Kurdistan University of Medical Sciences, Sanandaj, IR Iran

⁴Department of Microbiology, Faculty of Medicine, Kurdistan University of Medical Sciences, Sanandaj, IR Iran ³Student Research Committee, Kurdistan University of Medical Sciences, Sanandaj, IR Iran ⁴Deputy of Public Health Affairs, Kurdistan University of Medical Sciences, Sanandaj, IR Iran

*Corresponding author: Babak Shahbazi, Department of Microbiology, Faculty of Medicine, Kurdistan University of Medical Sciences, Sanandaj, IR Iran. Tel: +98-9353929575, E-mail: microbiol_sci2013@yahoo.com

Received: February 9, 2014; Revised: April 12, 2014; Accepted: April 23, 2014

Background: Vibrio cholerae causes diarrhoeal disease that afflicts thousands of people annually. V. cholerae is classified on the basis of somatic antigens into seroyroups and there are at least 200 known serogroup. Two serogroups, OI and O139 have been associated with epidemic diseases. Virulence genes of these bacteria are OmpW, ctxA and tcpA.

Objectives: Due to the importance of V. cholerae infection and developing molecular diagnostics of this organism in medical and microbiology sciences, this study aimed to describe molecular characterization of V. cholerae isolated from clinical samples using a molecular method.

Materials and Methods: In this study, 48 samples were provided during summer 2013 (late August and early September) by reference laboratory. Samples were assessed using biochemical tests initially. The primer of OmpW. ctxA and tcpA genes was used in Polymerase Chain Reaction (PCR) protocols. Enterobacterial Repetitive Intergenic Consensus (ERIC) PCR and Repetitive Extragenic Palindromic (REP)-PCR methods were used to subtype V. cholerae.

Results: In this study, from a total of 48 clinical stool samples 39 (81.2 %) were positive for V. cholerae in biochemical tests and bacteria culture tests. The PCR results showed that of 39 positive isolates 35 (89.7%), 34 (87.1%) and 37 (94.8%) were positive for *ctxA*, *tcpA* and *OmpW* gene, respectively. Also, in the REP-PCR method with ERIC primer strains were divided into 10 groups. In the REP-PCR method with REP primer, strains were divided into 13 groups.

Conclusions: Polymerase chain reaction has specificity and accuracy for identification of the organism and is able to differentiate biotypes. Enterobacterial repetitive intergenic consensus sequence is one of the informative and discriminative methods for the analysis of V. cholerae diversity. The REP-PCR is a less informative and discriminative method compared to other methods for the analysis of V. cholerae diversity.

Keywords: Molecular Characterization; Vibrio cholerae; Clinical Samples; Virulence Genes

1. Background

Cholera is an important infectious diarrheal disease and sometimes causes death in humans. This disease is both epidemic and endemic in different regions of the world, especially in the countries that lack proper sanitary management. Pandemics that are caused by this bacterium have severely affected many countries in multiple continents over a span of many years (1). Vibrio cholerae is a natural inhabitant of aquatic environments and the toxigenic strains are causative agents of Cholera disease. These bacteria are rod-like, Gram-negative, facultative anaerobe and motile (2, 3).

Pathogenicity of Vibrio spp. is depended on the expression of some virulence factors such as Cholerae Toxin (CT) that is facilitated by the Toxin-Coregulated Pilus (TCP). These two genes are encoded by the *ctx* and *tcp* genes, respectively (4). Another virulence factor is the Outer-Membrane Protein W (*OmpW*) that is involved in stimulating the immune response via induction of protective immunity. It also plays an important role in bacterial pathogenesis by increasing the adaptability of pathogenic strains and is encoded by OmpW gene (5).

Different studies showed that toxigenic V. cholerae that caused epidemic and pandemic potential forms of cholerae diarrhea are O1 or O139 sero-groups. V. cholerae O1 and O139 serogroups are positive for ctx, tcp and OmpW genes (6).

Traditionally, identification of Vibrio spp. is consisted as isolation on the selective blood agar medium, followed by biochemical and serological testing (7). However, Polymerase Chain Reaction (PCR) is a rapid and versatile in vitro method for identification of Vibrio spp. and it amplifies defined target DNA sequences that are present

Copyright © 2015, Ahvaz Jundishapur University of Medical Sciences. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

within a source of DNA. Also, this method is designed to permit selective amplification of a specific targeted DNA sequence within a heterogeneous collection of DNA sequences (8). Enterobacterial Repetitive Intergenic Consensus sequence (ERIC)-PCR is a method for interspecies profiling of several bacteria such as Bacillus anthracis and B. cereus, Enterobacter sakazakii, Lactobacillus, Listeria monocytogenes, Salmonella enteritidis and it provides discriminative DNA patterns for these bacteria (9). Also, Repetitive Extragenic Palindromic (REP)-PCR is a technique that has been used to study epidemiological relationships among V. cholerae isolates and DNA fingerprints production. It is used to discriminate among bacterial species or strains of the same species. This technique includes the application of oligonucleotide primers based on families of short and highly conserved extragenic repetitive sequences (9, 10).

Different studies showed molecular characterization of *V. cholerae* in different samples; Waturangi et al. (9) in Indonesia, with examining 75 isolates of *V. cholerae* from ice samples and using the REP- and ERIC-PCR methods showed that fingerprinting profiles of *V. cholerae* isolated from ice samples were very diverse. In another study, Alishahi et al. (11) in Iran, by using multiplex PCR on 72 *V. cholerae* samples of stool showed that all of them carried *toxR* and *ompU* genes, but *ctxA* gene was detected in only 61 isolates. Also, Naha et al. (12) in India, in 180 diarrheal stool samples detected 247 *V. cholerae* O1 and by using PCR detected *rstR*, *rtxC* and *tcpA* genes (12).

2. Objectives

V. cholerae O1 and O139 are causative agents of cholerae epidemics and endemics. Therefore, molecular detection and identification of *V. cholerae* strains are important. The REP and ERIC-PCR are fast, simple and valuable methods for molecular typing, genomic fingerprinting and proving of relationships within and among *V. cholerae* strains, which can be used in epidemiological studies of this microorganism infections (13, 14). Given the importance of the items listed above, this study aimed to describe molecular characterization of isolated *V. cholerae* from clinical samples by REP and ERIC-PCR methods in Kurdistan Province, Iran.

3. Materials and Methods

3.1. Samples

Forty-eight clinical specimens that suspected for cholerae were considered (under the health center, Kurdistan Province, Iran). Stool samples were collected from these clinical specimens in different cities of Kurdistan Province, Iran, summer 2013. Samples were collected using sterile rectal swabs.

3.2. Bacterial Cultures

Collected samples were transferred to Cary-Blair trans-

port medium (Merck, Germany). Alkaline peptone water was used for enrichment (Merck, Germany) and then bacteria were isolated on Thiosulphate Citrate Bile Salts Sucrose (TCBS) agar plates (Merck, Germany).

3.3. Biochemical and Agglutination Characterization

All bacterial isolates were screened by the biochemical test including oxidase reaction, motility status on Sulfur-Indole-Motility (SIM) agar (Merck, Germany), Triple Sugar Iron (TSI) agar (Merck, Germany) for sucrose, mannose and arabinose fermentation tests, Kligler Iron Agar (KIA) (Merck, Germany) for glucose fermentation, lactose fermentation, sulfur reduction and Methyl Red Voges-Proskauer test (MRVP) (Merck, Germany). Agglutination test in the presence of polyvalent O1 and O139 antisera was used for confirmation of *V. cholerae* strains (Mast Diagnostics Ltd., Bootle, Mersey side, UK). Positive samples (O1 and O139) were placed in a Luria-Bertani (LB) broth (Merck, Germany) with 30% glycerol at -80°C for other tests (11).

3.4. DNA Extraction

For DNA extraction, screened isolates were cultured in a LB broth for 24 hours at 37°C; DNA genome of each sample was purified using a DNA extraction kit according to manufacturer's instructions. Afterwards, 500 µl Thioglycollate broth (Merck, Germany) and 2×10^9 (10 - 20 mg) bacterial cells were added to 2ml tubes, then they were vortexed. In other step, tubes were centrifuged at 4.000 rpm for 5 minutes. Then 100 μ l of prelysis buffer and 10 μ l Ribotinase were added to the tubes and they were incubated at 55°C for 30 minutes. Then 100 µl of each samples were added to a sterile 1.5 or 2 polypropylene tubes. After this stage, 400 µl of lysis buffer was added to tubes and they was vortexed at max speed for 20 seconds, then 300 µl precipitation solutions was added to them and they vortexed at max speed for 5 seconds. Solution was transferred to a spin column with a collection tube. Tubes were centrifuged at $12.100 \times g$, 13.000rpm for 1 minute. Spin column was placed in a new collection tube and 400 μ l wash buffer 1 was added to tubes. Then they were centrifuged at $12.100 \times g$, 13.000 rpm for 1 minute. After centrifuge, spin column was placed in a new collection tube and wash buffer 2 was added to it and it was centrifuged at 12.100 × g, 13.000 rpm for 1 minute (wash buffer 2 stage was repeated). Column was carefully transferred to a new 1.5 ml tube. Afterwards, 50 µl of 65°C preheated elution buffer was poured in the center of column and then it was incubated for 3 - 5 minutes at 65°C. This solution was centrifuged at 12.100 × g, 13.000 rpm for 1 minute (According to the manufacturer's instructions, (CinnaGene, Iran).

3.5. Polymerase Chain Reaction Assay

Uniplex PCR assays were performed for *tcpA*, *ctxA* and *OmpW* genes in a reaction volume of 26 μ l; 26 μ l of the reaction mixture contained the following reagents:

Mastermix: 12 µl, MgCl₂: 25 µl, Distilled Water (DWs): 8.75 μl, primer (pr) Reverse (R): 1 μl, primer Forward (F): 1 μ l and 3 μ l DNA. The cycling profile for *ctxA* and *OmpW* was as follows: initial denaturation was at 94°C for 5 minutes, and 35 cycles of denaturation at 94°C for 1 minute, annealing at 59°C for 1 minute, extension at 72°C for 1 minute and final extension at 72°C for 16 minutes. The cycling profile for *tcpA* was as follows: initial denaturation at 94°C for 5 minutes. 35 cycles of denaturation at 94°C for 1 minute, annealing at 61°C for 1 minute, extension at 72°C for 1 minute and final extension at 72°C for 16 minutes. The PCR products were electrophoresed at100 v for approximately 1 hour on a 1.5% agarose gel in 0.5- TAE. The PCR primer set that was used: pr ctxA-R: (5' - CAAGCACCCCAAAATGAACT-3'), pr ctxA-F: (5' - TTGTTAG-GCACGATGATGGA-3'), pr tcpA-R: (5' - GACTAAGGCTGCG-CAAAATC-3'), pr *tcpA*-F: (5' - CCCCTACGCTTGTAACCAAA-3'), pr ompW-R: (5' - GCATCTGCACCTGCTTTGTA-3') and pr *OmpW*-F: (5' – ACTTGACGACTCATGGGGAC-3') (Cinna Clon, Iran) (11).

3.6. Enterobacterial Repetitive Intergenic Consensus Polymerase Chain Reaction Assay

The amplification protocols that were employed for ERIC-PCR was as follows: 1 cycle of 95°C for 5 minutes, 35 cycles of 92°C for 45 seconds, 52°C for 1 minute and 70°C for 10 minutes with a final extension step of 70°C for 10 minutes. The PCR products were electrophoresed at 100V for approximately 1 hour on a 1.5% agarose gel (Merck, Germany) in 0.5-TAE (Tris-acetate- EDTA) (Merck, Germany). The PCR primer set that was used as follows: pr R: (5'-ATGTAAGCTCCTGGGGATTCAC-3') and pr F: (5' - AAGTA-AGTGACTGGGGTGAGCC-3') (Cinna Clon, Iran) (9).

3.7. Repetitive Extragenic Palindromic Polymerase Chain Reaction Assay

The amplification protocols that were employed for REP-PCR was as follows: 1 cycle of 94° C for 5 minutes, 35 cycles of 94° C for 1 minute, 61° C for 1 minute and 72° C for 1 minute with a final extension step of 72° C for 16 minutes. Then PCR products were electrophoresed at100 v for approximately 1 hour on a 1.5% agarose gel in 0.5-TAE. The PCR primer set used: pr R: (5' – IIIICGICATCIGGC-3') and pr F: (5' - ICGICTTATCIGGCCTA-3') (Cinna Clon, Iran) (9).

3.8. Composed Matrix of 0 and 1

In the using of molecular markers, schemas are bands that are produced with a primer. After electrophoresis, the presence or absence of bands were determined with the numbers 0 and 1 for the isolates, respectively. For data analysis, a 0 and 1 similarity matrix was drawn. Detection of bands was performed by manual editing. After the similarity matrix formation, data were analyzed using the software NTSYS-pc ver 2. 02 and cluster analysis was performed (14).

4.1. Biochemical Tests and Bacterial Culture Results

Biochemical tests and bacterial culture results showed that of 48 clinical specimen samples that suspected for cholera, 39 (81.2%) samples were positive for *V. cholerae*.

4.2. Polymerase Chain Reaction Assay Results

The PCR assay results showed that of 39 *V. cholerae* positive samples, *ctxA* was present in 35 (89.7%) (Figure 1), *tcpA* was present in 34 (87.1%) (Figure 2) and *OmpW* was present in 37 (94.8%) samples (Figure 3). These results showed specificity for all tested *V. cholerae* strains. The nucleotide sequence of the *tcpA*, *ctxA* and *OmpW* genes had revealed an amplicon of 287-252 and 132 Base Pair (bp).

4.3. Repetitive Extragenic Palindromic Polymerase Chain Reaction Results With Enterobacterial Repetitive Intergenic Consensus Primer

The results achieved by REP-PCR with the ERIC primer showed that strains were divided into 10 groups. Similarity coefficient between these groups was 1% to 51%. Most strains were placed in group 4, which had 30% similarity coefficient (Figures 4 and 6).



Figure 1. Gel Electrophoresis of the Polymerase Chain Reaction Products From CtxA Gene



Figure 2. Gel Electrophoresis of the Polymerase Chain Reaction Products From *Tcpa* Gene

www.SID.ir



Figure 3. Gel Electrophoresis of the Polymerase Chain Reaction Products From *OmpW* Gene



Figure 4. The Frequency of Different Genotypes Obtained From Repetitive Extragenic Palindromic Polymerase Chain Reaction Results With the Enterobacterial Repetitive Intergenic Consensus Primer in This Study

4.4. Repetitive Extragenic Palindromic Polymerase Chain Reaction Results With the Repetitive Extragenic Palindromic Primer

The results that obtained from the REP-PCR method with the REP primer showed that strains were divided into 13 groups and similarity coefficient between these groups was 1% to 50%. Maximum strains were placed in group 1 and similarity coefficient was 22% (Figures 5 and 6).



Figure 5. The Frequency of Different Genotypes Obtained From Repetitive Extragenic Palindromic Polymerase Chain Reaction Results With Repetitive Extragenic Palindromic Primer in This Study



Figure 6. Dendrogram Analysis of V. cholerae Gene

5. Discussion

V. cholerae causes one of the most potent diarrheal diseases in the world. Toxigenic *Vibrio* spp. expresses virulence factors of different genes such as *ctxA*, stn, *OmpW* and *tcpA* (15, 16).

Goel and Jiang (17) in India showed that out of 114 suspected clinical cases for *V. cholerae* in different locations from 2004 to 2007, 100% were positive for *V. cholerae* O1 strains. According to biochemical tests and bacterial cultures in our study, out of 48 clinical specimens that suspected for cholera, 39 (81.2%) were positive. These results indicate that a large number of samples were unfortunately positive for this bacteria. However, in our study 9 (18.7%) samples were not positive for *V. cholerae*. Different factors such as poor economics, inadequate sanitation, poor hygienic practices, poor nutritional status and low immunity may be effective on *V. cholerae* rate that was isolated from patients in different regions (15).

Sharma et al. (16) in India using the PCR method reported that out of 115 environmental isolates of Vibrio, *ctxA* gene was isolated from 14 (13%) and *OmpW* gene existed in 100% of the samples. The PCR results in our research showed that from 39 *V. cholerae* isolates, 35 (89.7%), 34 (87.1%) and 37 (94.8%) were positive for *ctxA*, *tcpA* and *OmpW* genes, respectively. These toxin genes are mobile among isolates and they spread in environment. It is possible that a different mechanism of gene transfer such as horizontal gene transfer between serotypes and genetic changes cause *V. cholerae* transmission in different geographic areas (16, 18).

Maleki et al. (19) by taking advantage of PCR, isolated *ctxA*, zot, ace and *tcpA* genes of 39 *V. cholerae* strains from the summer epidemic in Iran. These genes were present in 89.7%, 84.6%, 100% and 100% of the isolates, respectively. Maleki's study results (19) are very close to our obtained results. Sampling in our study was conducted in summer (late August and early September). Virulence genotype and phenotype of these pathogens that are associated with infection may differ according to climatic changes (20). Some studies showed seasonal patterns of *V. cholerae* transmission. Larger number of these transmissions occurred in summer (from June to September) compared to winter (21). Consumption of untreated water and uncooked seafood in summer is another epidemiologic evidence of *V. cholerae* transmission (22).

Pourshafie et al. (23) in Iran studying 50 *V. cholerae* O1 serotype Inaba isolates that were collected during several cholerae outbreaks during the summer by PCR method showed that 100%, 98% and 98% carried the *ctx, zot* and *ace* genes, respectively. Also, in the Pourshafie's study strains were genotyped using Randomly Amplified Polymorphic DNA (RAPD), Pulsed-Field Gel Electrophoresis (PFGE) and ribotyping techniques. In our study a PCR-based technique was used for detection of toxigenic genes of *V. cholerae* O1 and O139 from different cities of Kurdistan Province in Iran. *V. cholerae* is a well-defined species on the

basis of biochemical tests and commercial biochemical identification systems, but these traditional methods are time- consuming, laborious or not always accurate. Also, these methods are not able to specify bacterial strains. So, many researchers use PCR techniques that are fast, have specificity and accuracy for identification of the organism (16, 24). Since strain typing and determination of relationships among *V. cholerae* strains are important processes for diagnosis, treatment and epidemiological investigations of this bacterium, molecular methods for genotyping and gene analysis are applied. Polymerase chain reaction is able to differentiate biotypes and also detect virulence factors (16, 18, 25).

Goel et al. (26) in India using ERIC- PCR, revealed similar DNA patterns during the outbreak in Chennai City. Moreover, the ERIC sequence is one of the most informative and discriminative methods for the analysis of *V. cholerae* diversity (9). Ten strain groups were obtained using ERIC-PCR in our results and it classified the tested strains into 4 groups with 30% similarity coefficient. These groups were also highly homogenous.

Taneja et al. (27) by PFGE, REP-PCR and ribotyping studied genetic characteristics of *V. cholerae* isolates during sporadic and outbreak cases. In Taneja's study, the REP-PCR divided all clinical isolates into four major profiles. Also, the REP-PCR method with REP primer in our study showed that strains were divided into 13 groups with 1% to 50% similarity coefficient. Group 1 had maximum strains with 22% similarity coefficient, also groups were highly homogenous. Different studies showed that REP-PCR is a less informative and discriminative method compared to other applied methods for the analysis of *V. cholerae* diversity. However, it is a good technique to differentiate clinical toxigenic isolates from nontoxigenic environmental isolates (9, 27).

In our results, extracted DNA of V. cholerae using the REP-PCR method showed different fingerprints that had a variety of sizes between 150 to 5400 bp. Also, the results of our study showed that V. cholerae that isolated from patients created different fingerprint patterns in REP-PCR, so, these results showed different origins for the infection and the fact that it was spreading. In cluster analysis of REP-PCR results, isolates were classified into 13 categories. In ERIC-PCR analysis, isolates were sorted in 10 categories. These results showed marker similarity between the isolates that were obtained from ERIC-PCR was more than REP-PCR. In the dendrogram of REP-PCR results, 6 samples had 22% similarity. In ERIC-PCR, most strains belonged to group 4, which had 30% similarity. The comparison of two primers used in this test revealed that results of REP primer had 1% - 50% similarity between selected isolates. Therefore, the results that belonged to REP-PCR, ERIC-PCR and cluster groups were similar. In conclusion, considering the prevalence of V. cholerae outbreaks in different countries, detection of reservoir of V. cholerae and

appropriate methods to prevent these outbreaks should be considered.

Acknowledgements

Cellular and molecular research center and microbiology department of Kurdistan University of Medical Sciences, Sanandaj, Iran, supported this study. The authors declare that there is no conflict of interest.

Authors' Contributions

Rashid Ramazanzadeh and Babak Shahbazi developed the original idea and protocol and also they analyzed data. Samaneh Rouhi and Pegah Shakib abstracted and wrote the manuscript. Farzam Bidarpour and Mohammad Karimi contributed to protocol development and preparation of the manuscript.

Financial Disclosure

There is no financial interest to disclose in this survey.

Funding/Support

This study was funded and supported by cellular and molecular research center and microbiology department, Kurdistan University of Medical Sciences, Sanandaj, Iran.

References

- Fiore AE, Michalski JM, Russell RG, Sears CL, Kaper JB. Cloning, characterization, and chromosomal mapping of a phospholipase (lecithinase) produced by Vibrio cholerae. *Infect Immun.* 1997;65(8):3112-7.
- Tracz DM, Backhouse PG, Olson AB, McCrea JK, Walsh JA, Ng LK, et al. Rapid detection of Vibrio species using liquid microsphere arrays and real-time PCR targeting the ftsZ locus. J Med Microbiol. 2007;56(Pt 1):56–65.
- Faruque SM, Asadulghani N, Rahman MM, Waldor MK, Sack DA. Sunlight-induced propagation of the lysogenic phage encoding cholera toxin. *Infect Immun.* 2000;68(8):4795–801.
- Goss TJ, Morgan SJ, French EL, Krukonis ES. ToxR recognizes a direct repeat element in the toxT, ompU, ompT, and ctxA promoters of Vibrio cholerae to regulate transcription. *Infect Immun.* 2013;81(3):884–95.
- Alizadeh J, Ranjbar R, Kamali M, Farhadi N, Davari A, Sadeghifard N. Cloning of Vibrio cholerae outer membrane protein W in Pichia pastoris. *Iran J Microbiol*. 2013;5(3):252–8.
- Talkington D, Bopp C, Tarr C, Parsons MB, Dahourou G, Freeman M, et al. Characterization of toxigenic Vibrio cholerae from Haiti, 2010-2011. *Emerg Infect Dis*. 2011;**17**(11):2122–9.
- Harwood VJ, Gandhi JP, Wright AC. Methods for isolation and confirmation of Vibrio vulnificus from oysters and environmental sources: a review. J Microbiol Methods. 2004;59(3):301–16.
- 8. Hu X, Zhang Y, Zhou X, Xu B, Yang M, Wang M, et al. Simultaneously typing nine serotypes of enteroviruses associated with hand, foot, and mouth disease by a GeXP analyzer-based multiplex reverse transcription-PCR assay. *J Clin Microbiol.* 2012;**50**(2):288–93.
- Waturangi DE, Joanito I, Yogi Y, Thomas S. Use of REP- and ERIC-PCR to reveal genetic heterogeneity of Vibrio cholerae from edible ice in Jakarta, Indonesia. *Gut Pathog.* 2012;4(1):2.
- 10. Caburlotto G, Lleo MM, Gennari M, Balboa S, Romalde JL. The use

of multiple typing methods allows a more accurate molecular characterization of Vibrio parahaemolyticus strains isolated from the Italian Adriatic Sea. *FEMS Microbiol Ecol.* 2011;77(3):611–22.

- Alishahi A, Imani Fooladi AA, Fallah Mehrabadi J, Mahmoodzadeh Hosseini H. Facile and Rapid Detection of Vibrio cholerae by Multiplex PCR Based on ompU, ctxA, and toxR Genes. *Jundishapur J Microbiol.* 2013;6(10):e7933.
- Naha A, Chowdhury G, Ghosh-Banerjee J, Senoh M, Takahashi T, Ley B, et al. Molecular characterization of high-level-choleratoxin-producing El Tor variant Vibrio cholerae strains in the Zanzibar Archipelago of Tanzania. J Clin Microbiol. 2013;51(3):1040-5.
- Mancuso M, Avendano-Herrera R, Zaccone R, Toranzo AE, Magarinos B. Evaluation of different DNA-based fingerprinting methods for typing Photobacterium damselae ssp. piscicida. *Biol Res.* 2007;40(1):85–92.
- Madhusudana Rao B, Surendran PK. Genetic heterogeneity of non-O1 and non-O139 Vibrio cholerae isolates from shrimp aquaculture system: a comparison of RS-, REP- and ERIC-PCR fingerprinting approaches. *Lett Appl Microbiol.* 2010;51(1):65-74.
- Centers for Disease Control and Prevention. Global Disease Detection (GDD) Manual Rapid Diagnostic Tests for Epidemic Diseases. 2011. Watery Diarrhea Differential Diagnosis: Outbreaks of Acute Watery Diarrhea.
- Sharma A, Chaturvedi AN. Prevalence of virulence genes (ctxA, stn, OmpW and tcpA) among non-O1 Vibrio cholerae isolated from fresh water environment. Int J Hyg Environ Health. 2006;209(6):521-6.
- Goel AK, Jiang SC. Genetic determinants of virulence, antibiogram and altered biotype among the Vibrio cholerae O1 isolates from different cholera outbreaks in India. *Infect Genet Evol.* 2010;10(6):815–9.
- Gubala AJ, Proll DF. Molecular-beacon multiplex real-time PCR assay for detection of Vibrio cholerae. *Appl Environ Microbiol.* 2006;72(9):6424–8.
- 19. Maleki A, Pakzad I, Hosseini S, Ranjbar R, Purshafie M, Mohebi R, et al. Frequency Analysis of Genes ctxA, tcpA, Ace and Zot in Vibrio Cholerae Isolated of Epidemy of Summer 2005 in Iran By PCR. *Sci J Ilam Univ Med Sci.* 2011;**18**(4):1–6.
- 20. Schirmeister F, Dieckmann R, Bechlars S, Bier N, Faruque SM, Strauch E. Genetic and phenotypic analysis of Vibrio cholerae non-01, non-0139 isolated from German and Austrian patients. *Eur J Clin Microbiol Infect Dis.* 2014;**33**(5):767–78.
- Scientific Committee on Enteric Infections and Foodborne Diseases.. Epidemiology, Prevention and Control of Cholera in Hong Kong. 2011. Available from: http://www.chp.gov.hk/files/pdf/ epidemiology_prevention_and_control_of_cholera_in_hong_ kong_r.pdf.
- 22. Florida Department of Health.. *Cholera (Toxigenic Vibrio cholerae* 01, 0139). *Guide to Surveillance and Investigation*. 2011. Available from: http://www.floridahealth.gov/diseases-and-conditions/ disease-reporting-and-management/disease-reporting-and-surveillance/_documents/gsi-vibrio-cholerae.pdf.
- Pourshafie M, Bakhshi B, Ranjbar R, Sedaghat M, Sadeghifard N, Yazdi J, et al. Dissemination of a single Vibrio cholerae clone in cholera outbreaks during 2005 in Iran. J Med Microbiol. 2007;56(12):1615–9.
- 24. Srisuk C, Chaivisuthangkura P, Rukpratanporn S, Longyant S, Sridulyakul P, Sithigorngul P. Rapid and sensitive detection of Vibrio cholerae by loop-mediated isothermal amplification targeted to the gene of outer membrane protein ompW. *Lett Appl Microbiol*. 2010;50(1):36–42.
- 25. Yildirim I. Molecular methods for bacterial genotyping and analyzed gene regions. *J Microbiol Infect Dis.* 2011;**1**(1):42–6.
- Goel AK, Jain M, Kumar P, Jiang SC. Molecular characterization of Vibrio cholerae outbreak strains with altered El Tor biotype from southern India. World J Microbiol Biotechnol. 2010;26(2):281–7.
- 27. Taneja N, Sangar G, Chowdhury G, Ramamurthy T, Mishra A, Singh M, et al. Molecular epidemiology of Vibrio cholerae causing outbreaks & sporadic cholera in northern India. *Indian J Med Res.* 2012;**136**(4):656–63.