

Antibiotic Resistance and Biofilm Formation of *Enterococcus faecalis* in Patient and Environmental Samples

Malihe Talebi,¹ Nastaran Asghari Moghadam,² Zeynab Mamooii,³ Mohsen Enayati,³ Mahnaz Saifi,⁴ and Mohammad Reza Pourshafie^{2,*}

¹Department of Microbiology, School of Medicine, Iran University of Medical Sciences, Tehran, IR Iran

²Department of Microbiology, Pasteur Institute of Iran, Tehran, IR Iran

³Microbiology Department, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, IR Iran

⁴Department of Bacteriology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, IR Iran

*Corresponding author: Mohammad Reza Pourshafie, Department of Microbiology, Pasteur Institute of Iran, Tehran, IR Iran. Tel/Fax: +98-2166405535, E-mail: pour@pasteur.ac.ir

Received: October 27, 2014; Revised: February 17, 2015; Accepted: March 28, 2015

Background: Enterococci are opportunistic pathogens and are a major factor in nosocomial infections. They may contain *ebp* operon, which upon expression makes them highly prone to biofilm formation on biotic and abiotic surfaces.

Objectives: The aim of the current study was to detect the polymorphism of *ebp* genes in *Enterococcus faecalis*.

Materials and Methods: Samples were isolated from patients (n = 58) and hospital environments (n = 32) of two hospitals in Tehran, Iran. All enterococcal species were identified by species-specific polymerase chain reaction (PCR); the antibiotic resistance pattern against nine antibiotics was determined. The *ebp A*, *ebp B*, *ebp C* and *srt C* genes were detected by PCR and the biofilm formation by the isolates was evaluated using the microtiter plate method. The genetic diversity of *ebp* genes was analyzed by restriction fragment length polymorphism (RFLP).

Results: The results indicated that, 86% of patient and 29% of environmental isolates carried *ebp* genes. The ability of the isolates to strongly attach was 62% and 71% for patient and environmental samples, respectively. The RFLP of the *ebp* showed no genetic variations amongst the isolates.

Conclusions: The results of the antibiotic resistance and other data suggest that there is a possible common clone of *E. faecalis*, which could rapidly disseminate in patients and the environment.

Keywords: Drug Resistance; Microbial; *ebp* operon; Biofilms; *Enterococcus faecalis*

1. Background

Enterococci are important nosocomial pathogens (1). They cause various infections including urinary tract infections, sepsis, endocarditis, wound infections and bacteremia (2). Amongst enterococci, *Enterococcus faecalis* is the most common species among clinical enterococcal infections. On the other hand, *E. faecium* causes 10 - 20 percent of infections yet in comparison to other enterococci species, they show higher antibiotic resistance (1, 3).

Enterococci, not only carry intrinsic resistance against some antibiotic agents, they can also acquire resistance via mutation, transposable elements or conjugative plasmids (1). In addition, the ability of enterococci to form biofilm has also been of growing concern in the recent years. The use of indwelling catheters can intensify biofilm production of enterococci, which would lead to increase in urinary infections in catheterized patients (4). Antibiotic resistance and biofilm formation in bacteria have been under intensive investigation in urinary tract infections and endocarditis, during the recent years (1), in particular the *pili* and its gene cluster

(*ebp*). This attention, of course, is worthwhile because biofilms can be the source of 80% of chronic infections in humans and their presence can increase antibiotic concentration by up to 1,000 folds (5, 6).

2. Objectives

In this study, we assessed the antibiotic resistance and biofilm formation ability of *E. faecalis* strains isolated from environmental and clinical samples in Tehran, Iran. In addition, the polymorphisms of *ebp* genes were determined.

3. Materials and Methods

3.1. Sampling, Isolation and Identification

From October 2009 to April 2010, a total of 90 clinical and hospital environmental isolates of *E. faecalis* were collected from two main hospitals in Tehran, Iran (Shariati Hospital (SH) and Shahid Rajaie Hospital (SRH)). Overall, 58 of the isolates were from patient samples

and 32 from environmental sites. The clinical isolates were collected from blood, wound, suction secretion, sterile body fluid and urine of catheterized patients. Environmental samples were collected from patients' bathroom, beds and tables and staff's bathrooms and tables, as well as ventilators and oxygen pumps in the patients' rooms.

The collected samples from hospitals were cultured on blood agar and incubated at 37°C for 24 hours. Next, the isolates were cultured on m-Enterococcus agar (Becton Dickinson Co., MD, USA). *Enterococcus faecalis* strains were detected based on colony morphology, gram staining and polymerase chain reaction (PCR). To amplify genus and species specific gene (*sodA*) of *E. faecalis*, PCR was performed as mentioned before (7).

3.2. Antimicrobial Susceptibility

Antimicrobial susceptibility test was performed by the disk diffusion method, according to the Clinical and Laboratory Standards Institute (CLSI) guidelines, for nine antimicrobial agents including: gentamycin (G) (10 µg), ciprofloxacin (CIP) (5µg), ampicillin (AMP) (10 µg), co-trimoxazole (TS) (25 µg), erythromycin (E) (15 µg), tetracycline (T) (30 µg), linezolid (LZD) (30 µg), teicoplanin (TEC) (30 µg) and vancomycin (V) (30 µg) (Mast Diagnostics Ltd, UK).

3.3. Biofilm Formation Assay by the Quantitative Method

Microtiter plate test was used to quantitatively assay biofilm formation as mentioned previously. The cut-off calculation was used to interpret the test results by the formula presented below: Cut-off = (mean (OD) ± 3SD (OD)) + OD (Blank). Strains with OD of ≤ 0.45, 0.45 - 0.55 and ≥ 0.55 were considered as non-adherent, weakly adherent and strongly adherent, respectively (4, 6, 8, 9).

3.4. Polymerase Chain Reaction Test to Detect Endocarditis and Biofilm-Associated Pili Genes (*ebpA*, *ebpB*, *ebpC*) and Sortase C (*srtC*)

Amplification of *ebpA*, *ebpB*, *ebpC* and *srtC* genes was done as described by previous studies (10, 11).

3.5. Restriction Fragment Length Polymorphism of Amplified Genes of *ebp* Operon

For amplified *ebpA*, *ebpB* and *ebpC*, gene products were digested by HaeIII, EcoRI and HindIII restriction enzymes, respectively. The results showed that *ebpA* gene digestion by HaeIII resulted in three fragments with a size of 223 bp, 480 bp and 361 bp. EcoRI digestion of *ebpB* produced 397 bp and 129 bp fragments. The 963 bp *ebpC* gene was divided to two fragments with a length of 760 bp and 199 bp by Hind III.

4. Results

4.1. Bacterial Isolates and Identification of *Enterococcus faecalis* Strains

Out of a total of 80 patient isolates of enterococci, 58 were *E. faecalis*. From the two hospitals, 49 and 9 isolates were from SH and SRH, respectively. Overall, hospital environmental samples were obtained from 250 different locations. The distribution of isolates obtained from patients and hospital environments are shown in Table 1.

4.2. Antibiotic Resistance of Isolates

Among the clinical isolates, 86% of isolates were resistant to tetracycline (86%). The resistance observed against teicoplanin (3%), vancomycin (9%), linezolid (9%) and ampicillin (9%) was low. Among environmental isolates only 25% were sensitive to tetracycline with no resistance against vancomycin, teicoplanin, linezolid and ampicillin. Overall, 20 different antibiotic resistance patterns were obtained for 58 patient isolates and nine for environmental isolates (Figure 1).

Table 1. Isolation sites of *Enterococcus faecalis* strains^a

Source of Isolates	Values
Clinical Isolates	
Urinary culture	16 (27)
Blood culture	14 (24)
Wound culture	11 (19)
Catheter	8 (14)
Suction secretion	5 (9)
Body sterile fluid	4 (7)
Total	58 (100)
Environmental Isolates	
Staffs' bathroom	7 (21.9)
Patients' bathroom	5 (15.6)
Staffs' table	5 (15.6)
Patients' table	3 (9.4)
Patients' bed	5 (15.6)
Ventilator	4 (12.5)
Oxygen pump	3 (9.4)
Total	32 (100)

^a The values are presented as No. (%)

4.3. Identification of Virulence and Biofilm Formation Genes

Polymerase chain reaction product sizes for *ebpA* (1064 bp), *ebpB* (530 bp) and *ebpC* (964 bp) were found in 50 (86%), 55 (95%) and 56 (97%) of the patient isolates and 29 (91%), 30 (94%) and 29 (91%) of environmental isolates, respectively. The PCR product of *srtC* gene was a 500 bp fragment. The presence of this gene was confirmed in all patient and 30 (94%) of the environmental isolates. Distribution of the mentioned genes among *E. faecalis* strains in this study is shown in Table 2.

4.4. Results of Biofilm Formation Assay by the Quantitative Method

Quantitative assay of biofilm formation was conducted three times for each strain. Amongst the 58 studied patient isolates 36, 6 and 16 isolates were strongly adherent, weakly adherent and non-adherent, respectively. Amongst 32 environmental isolates, 23, 4 and 5 were strongly adherent, weakly adherent and non-adherent.

4.5. Results of Antibiotic Sensitivity Test and Biofilm Formation Assay

The antibiotic resistance and the ability of biofilm formation of all 90 isolates of *E. faecalis* were examined. The samples were divided to two groups; biofilm positive isolates with strongly and weakly adherent and biofilm negative isolates with non-adherent strains. The results are shown in Table 3.




Figure 1. Antibiotic Resistance Pattern Amongst Patient and Environmental Isolates

Table 2. Distribution of Genes Among *Enterococcus faecalis* Strains^a

The number of genes	Clinical Isolates	Environmental Isolates
4 genes	47 (81)	25 (78)
3 genes	9 (16)	6 (19)
2 genes	2 (3)	0 (0)
1 gene	0 (0)	0 (0)
None of the genes	0 (0)	1 (3)
Total	58 (100)	32 (100)

^a The values are presented as No. (%).

Table 3. Antibiotic Resistance Amongst Biofilm Positive and Biofilm Negative Strains^a


Antibiotics	Clinical Isolates		Environmental Isolates	
	Biofilm Positive	Biofilm Negative	Biofilm Positive	Biofilm Negative
Vancomycin	5	19	0	0
Teicoplanin	0	13	0	0
Linezolid	10	6	0	0
Tetracycline	95	62	81	40
Erythromycin	67	50	48	20
Cotrimoxazole	76	50	26	20
Ampicillin	0	31	0	0
Ciprofloxacin	63	63	26	0
Gentamicin	44	44	30	0

^a The values are presented as percent.

4.6. Results of Restriction Fragment Length Polymorphism Analysis of *ebp* A, B and C Gene Fragments

Out of 90 isolates, 72 had all four genes. Twenty-three including 15 patient and 8 environmental isolates were selected and polymorphism in their *ebp* gene by RFLP was examined (Figure 2). The results showed no difference in the obtained fragment patterns in all analyzed isolates.

Figure 2. Fragments Derived From Enzymatic Restriction of *ebp* Operon by Hae III, EcoRI and Hind III



5. Discussion

In this study, strains of *E. faecalis* isolated from patient and environmental specimens were analyzed. The patient isolates showed higher antibiotic resistance in comparison to environmental isolates, which could be due to higher transferability of antibiotic resistant genes amongst the patient isolates than environmental isolates. More than 60% of the patient isolates were resistant to tetracycline, erythromycin, co-trimoxazole, ciprofloxacin and gentamycin. On the other hand, the environmental isolates showed resistance only against tetracycline. Furthermore, 5% of the patient isolates were resistant to vancomycin (VRE), whereas none showed this resistance amongst environmental isolates. Other investigators have found various *E. faecalis* resistance rates to vancomycin, up to 11% in Europe (7), or 17% in isolates from hospital staff hands in Nigeria (12). In this study only 7% (5) of the patient isolates were sensitive to all tested antibiotic agents. Out of all *E. faecalis* examined, 43 were resistant to three or more antibiotics. Isolation of strains with identical antibiotic resistant patterns indicates widespread dissemination of resistance genes among strains as well as distribution of a common clone in different hospital wards.

In accordance with other reports (13, 14), we also found that biofilm formation was higher in isolates with antibiotic resistance to vancomycin, teicoplanin and ampicillin (as shown in Table 3). This may suggest a genetic linkage between the biofilm and these antibiotic genes. This conclusion was, however, not inclusive for all antibiotics. Quantitative assessment of biofilm formation exhibited that 72% and 85% of all patient and environmental isolates could produce biofilms, respectively. This finding was similar to reports by other investigators. In Japan 62.5% (4), Spain 62% (6) and in USA 79% (9) of the *E. faecalis* clinical samples were found to form biofilms.

Other studies showed a connection between genes and biofilm existence (15). The presence of *ebp* gene has been shown to be variable. In the present study, 86% of the isolates carried all four genes, which is lower than previous reports (16, 17). This difference could be due to the fact that there is variable genetic make-up amongst the isolates within the *ebp* gene. Restriction Fragment Length Polymorphism analysis of our amplified product, however, revealed no such difference which suggests complete gene sequencing is required. The lack of variants in RFLP analysis may indicate the dissemination of a single *E. faecalis* clone within the hospital setting, which may have gone under slight single nucleotide mutations.

In order to identify the effects of *ebp* genes in biofilm formation by *E. faecalis*, the ability of biofilm formation of isolates having four *ebp* and *srtC* genes were compared with isolates in which the presence of one or more genes had not been confirmed. Eighty-one percent of patient isolates having four genes could produce biofilms. On the other hand, 61% of strains not possessing one or two of the genes did not form biofilms. For environmental isolates the two possibilities mentioned above were 48% and 14%, respectively. It can be concluded that *ebp* genes play an essential role in biofilm formation, since the absence of one or two genes have a negative role in production of biofilms. The study of Nallapareddy on *E. faecalis* OG1RF revealed that mutation in *ebp A-srt* considerably reduced the ability of biofilm formation.

In conclusion the high antibiotic resistance seen in this study may indicate the intrinsic resistance of enterococci; circulation of transposable elements carrying resistant genes in clinical isolates, indiscriminate and uncontrolled usage of antibiotics and the presence of biofilm are reasons for higher antibiotic resistance among the patient isolates when compared to environmental isolates.

Authors' Contributions

Study concept and design: Malihe Talebi and Zeynab Mamooii. Analysis and interpretation of data: Mohsen Enayati and Nastaran Asghari Moghadam. Drafting of the manuscript: Nastaran Asghari Moghadam. Critical revision of the manuscript for important intellectual content: Mahnaz Saifi and Mohammad Reza Pourshafie. Statistical analysis: Zeynab Mamooii.

References

1. Fisher K, Phillips C. The ecology, epidemiology and virulence of *Enterococcus*. *Microbiology*. 2009;**155**(Pt 6):1749-57.
2. Murray BE. The life and times of the *Enterococcus*. *Clin Microbiol Rev*. 1990;**3**(1):46-65.
3. Giraffa G. *Enterococci* from foods. *FEMS Microbiol Rev*. 2002;**26**(2):163-71.
4. Seno Y, Kariyama R, Mitsuhashi R, Monden K, Kumon H. Clinical implications of biofilm formation by *Enterococcus faecalis* in the urinary tract. *Acta Med Okayama*. 2005;**59**(3):79-87.
5. Mohamed JA, Huang DB. Biofilm formation by enterococci. *J Med Microbiol*. 2007;**56**(Pt 12):1581-8.
6. Tendolkar PM, Baghdayan AS, Gilmore MS, Shankar N. Enterococcal surface protein, Esp, enhances biofilm formation by *Enterococcus faecalis*. *Infect Immun*. 2004;**72**(10):6032-9.
7. Kuhn I, Iversen A, Finn M, Greko C, Burman LG, Blanch AR, et al. Occurrence and relatedness of vancomycin-resistant enterococci in animals, humans, and the environment in different European regions. *Appl Environ Microbiol*. 2005;**71**(9):5383-90.
8. Mohamed JA, Huang W, Nallapareddy SR, Teng F, Murray BE. Influence of origin of isolates, especially endocarditis isolates, and various genes on biofilm formation by *Enterococcus faecalis*. *Infect Immun*. 2004;**72**(6):3658-63.
9. Toledo-Arana A, Valle J, Solano C, Arrizubieta MJ, Cucarella C, Lamata M, et al. The enterococcal surface protein, Esp, is involved in *Enterococcus faecalis* biofilm formation. *Appl Environ Microbiol*. 2001;**67**(10):4538-45.
10. Nallapareddy SR, Singh KV, Sillanpaa J, Garsin DA, Hook M, Erlandsen SL, et al. Endocarditis and biofilm-associated pili of *Enterococcus faecalis*. *J Clin Invest*. 2006;**116**(10):2799-807.
11. Kemp KD, Singh KV, Nallapareddy SR, Murray BE. Relative contributions of *Enterococcus faecalis* OG1RF sortase-encoding genes, *srtA* and *bps* (*srtC*), to biofilm formation and a murine model of urinary tract infection. *Infect Immun*. 2007;**75**(11):5399-404.
12. David OM, Oluduro AO, Olawale AK, Osuntuyinbo RT, Olowe OA, Famurewa O. Incidence of multiple antibiotic resistance and plasmid carriage among *Enterococcus faecalis* isolated from the hands of health care workers in selected hospitals in Ekiti, Ondo and Osun states, Nigeria. *Int J Acad Res*. 2010;**22**:43-7.
13. Sandoe JA, Wysesome J, West AP, Heritage J, Wilcox MH. Measurement of ampicillin, vancomycin, linezolid and gentamicin activity against enterococcal biofilms. *J Antimicrob Chemother*. 2006;**57**(4):767-70.
14. White HJ. Correlation of single multiple species biofilms and antibiotic resistance.; Missouri Junior Academy of Science.; University of Central Missouri. 2007.
15. Moniri R, Ghasemi A, Moosavi SGA, Dastehgoli K, Rezaei M. Virulence Gene's Relationship With Biofilm Formation and Detection of *aac* (6')/*aph* (2'') in *Enterococcus faecalis* Isolated From Patients With Urinary Tract Infection. *Jundishapur J Microbiol*. 2013;**6**(5):e6244.
16. Cobo Molinos A, Abriouel H, Omar NB, Lopez RL, Galvez A. Detection of *ebp* (endocarditis- and biofilm-associated pilus) genes in enterococcal isolates from clinical and non-clinical origin. *Int J Food Microbiol*. 2008;**126**(1-2):123-6.
17. Sillanpaa J, Xu Y, Nallapareddy SR, Murray BE, Hook M. A family of putative MSCRAMMs from *Enterococcus faecalis*. *Microbiology*. 2004;**150**(Pt 7):2069-78.