

Epidemiologic and Drug Resistance Pattern of *Vibrio cholerae* O1 Biotype El Tor, Serotype Ogawa, in the 2011 Cholera Outbreak, in Alborz Province, Iran

Hojatolah Barati,¹ Ghobad Moradi,² Mohammad Aziz Rasouli,³ and Parvin Mohammadi^{4,*}

¹Health Deputy, Alborz University of Medical Sciences, Karaj, IR Iran

²Social Determinants of Health Research Center, Kurdistan University of Medical Sciences, Sanandaj, IR Iran

³Department of Epidemiology and Biostatistics, School of Medicine, Kurdistan University of Medical Sciences, Sanandaj, IR Iran

⁴Department of Nursing and Midwifery, Sanandaj Branch, Islamic Azad University, Sanandaj, IR Iran

*Corresponding author: Parvin Mohammadi, Department of Nursing and Midwifery, Sanandaj Branch, Islamic Azad University, Sanandaj, IR Iran. Tel: +98-33235826, Fax: +98-33235826, E-mail: moham_pa@yahoo.ca

Received 2014 September 8; Revised 2014 December 29; Accepted 2015 January 14.

Abstract

Background: Although the national guidelines recommend special antibiotics, based on the antibiogram of National Reference Laboratory, it seems that, because of uncontrolled usage of antibiotics in the society and due to the changes in the serotypes causing the disease, it is essential to monitor the status of drug resistance, permanently, and to revise the current prescriptions guidelines.

Objectives: This study aimed to assess the epidemiological aspects and drug resistance pattern of *Vibrio cholerae* O1, biotype El Tor, serotype Ogawa, in cholera outbreak, in Alborz province in Iran, during 2011.

Materials and Methods: This is a cross-sectional study, which reviews a cholera epidemic that occurred in Iran. A total of 9844 specimens were taken from suspected cases, among diarrheal patients, via rectal swabs. The specimens were placed in Cary-Blair transport medium and sent to laboratory. Samples were enriched, in alkaline peptone water, and isolated on thiosulphate-citrate-bile salt-sucrose agar. From the 244 confirmed cases, 239 cases underwent antibiogram test, via disk diffusion method and based on national committee for clinical laboratory standards (NCCLS) instructions. The standard *Escherichia coli* ATCC 25922 was used for antibiogram quality control and, eventually, all results were interpreted and reported using NCCLS standard table.

Results: In total, until October 22, 2011, which was announced as the end of outbreak, 9844 samples were taken from diarrheal patients. Regarding the type of *V. cholerae*, 244 El Tor biotype positive cases were reported. The case fatality rate was 1.3%. The mean age of patients was 37.8 years and the highest incidence rate occurred in the age group 21 - 30 years. After conducting antibiotic susceptibility test in the 244 *V. cholerae*, biotype El Tor, serotype Ogawa, it was found that ciprofloxacin had the highest level of antibiotic susceptibility (99.6%) and the highest level of antibiotic resistance was observed in co-trimoxazole (95.4%).

Conclusions: The results of our study show that the resistances to doxycycline and tetracycline, which are mentioned in multiple resources, as the most common antibiotic drugs for treating cholera, are increasing.

Keywords: Antibiotic Resistance, Bacterial, Outbreaks, Iran, Ciprofloxacin, Co-Trimoxazole, *Vibrio cholerae* O1

1. Background

Cholera is a major public health concern, in countries where access to safe water and adequate sanitation cannot be guaranteed, for all residents. *Vibrio cholerae* serogroups O1 and O139 are the causative agents of cholera (1). Reports from the epidemic forms of cholera diarrhea, particularly in developing countries, show that toxigenic *V. cholerae* with O1 or O139 serogroup is the main causative microorganism. Cholera toxin, encoded by *ctx* gene, is a potent factor, which induces intense dehydration and pathogenesis (2). There are two biotypes of *V. cholerae* O1, classical and El Tor, which are believed to have evolved from separate lineages among the two biotypes, strains belonging to the El Tor biotype have better adaptability to survive in the environment and in the human host (3). Diagnosis of *V. cholerae*, based on multiplex polymerase chain reaction (PCR) can be useful.

Multiplex PCR provides the possibility of evaluating several genes, through one PCR reaction, in a short time interval, by designing different target-specific primers. This method is simple, fast and cost-saving and has the ability to test a large number of samples, at the same time. Moreover, technicians can interpret the results easily (2). In 2012, the world health organization estimated that every year, there are 3 - 5 million cholera cases and more than 100,000 people die of the disease, with the majority (99%) of the cases and deaths occurring in Sub-Saharan Africa and Southern Asia (4). However, *V. cholerae* serogroup O1 still is the most common cause of cholera epidemics. Cholera is an endemic disease in Iran and is common in summer and early month of autumn. The causative serogroup, in our country, is O1 and we did not have any report for isolation of O139 serogroup in recent outbreaks (5).

Alborz is one of Iran's provinces, which has a population of 2,412,513 people, based on 2011 census. During the last recent decades, several cases of cholera epidemics have occurred in Karaj, in 3 - 5-year intervals (6). In each outbreak, it has always been a big concern to choose the type of antibiotic. Although the national guidelines recommend special antibiotics, based on the antibiogram of national reference laboratory (7), it seems that, because of uncontrolled usage of antibiotics in the society and due to the changes in the serotypes causing the disease, it is essential to monitor the status of drug resistance permanently and to revise the current prescriptions guidelines (8).

2. Objectives

In this study, we aimed to determine the antibiotic susceptibility pattern of *V. cholerae*. Therefore, we reviewed the epidemiologic characteristics of the disease and conducted the antibiotic susceptibility test for the *V. cholerae* samples, collected from El Tor infected patients, in 2011.

3. Materials and Methods

This was a cross-sectional study, which reviewed a cholera epidemic that occurred in Alborz province, in 2011. From the beginning of the cholera epidemic (June 14, 2011) until the end of the disease (October 22, 2011), a total of 9844 specimens were taken from suspected cases, among diarrheal patients, via rectal swabs. The specimens were placed in Cary-Blair transport medium and sent to laboratory. Samples were enriched in alkaline peptone water and isolated on thiosulphate-citrate-bile salt-sucrose agar. Strains were serotyped using polyva-

lent O1 and monospecific Inaba and Ogawa antisera (9). All antisera type determination serotyping test were performed with 1TOR 1R polyvalent antiserum 1T and monovalent Ogawa and Inaba (Baharafshan institute of research and development, Tehran, Iran) were purchased, which was approved by the national reference laboratory. From the 244 confirmed cases, 239 cases underwent antibiogram test, via disk diffusion method (Kirby-Bauer) and based on National committee for clinical laboratory standards (NCCLS) instructions. The standard *Escherichia coli* ATCC 25922 was used for antibiogram quality control and, eventually, all results were interpreted and reported, using NCCLS standard table (10).

Using the chi-square and Fisher exact test, the relationship between antibiotic resistance and other variable was tested.

4. Results

The first case of cholera occurred in a 58-year-old man, in June 14, 2011. In total, until October 22, 2011, which was announced as the end of outbreak, 9844 samples were taken from diarrheal patients. Regarding the type of *V. cholerae*, 244 El Tor biotype positive cases were reported. All cases were serotype Ogawa. Figure 1 shows the trend of the incidence of El Tor, in the province, in terms of days. From all patients in Alborz province, three patients died. Therefore, the case fatality rate reached 1.3%. The mean age of patients was 37.8 years and the highest incidence rate occurred in the age group 21 - 30 years. Based on antibiogram tests, ciprofloxacin had the highest level of antibiotic susceptibility (99.6%) and co-trimoxazole showed the highest level of antibiotic resistance (95.4%).




Figure 1. Distribution of Patients Based on the Time of the Emergence of Symptoms in Patients With Acute Watery Diarrhea Caused by *V. cholerae* in 2011 in Alborz Province

Table 1. *V. cholerae* Antibiogram Result Based on the Type of Antibiotics, in Alborz Province, 2011^a

Result of Antibiotic Susceptibility	Ciprofloxacin	Azithromycin	Tetracycline	Doxycycline	Erythromycin	Furazolidone	Ampicillin	Co-Trimoxazole
Susceptible	238 (99.6)	227 (95)	194 (81.2)	209 (84.7)	6 (2.5)	37 (15.5)	33 (13.8)	7 (2.9)
Resistant	0 (0)	2 (0.8)	36 (15.1)	10 (4.2)	13 (5.4)	202 (84.5)	110 (46)	228 (95.4)
Intermediate	1 (0.4)	2 (4.2)	9 (3.8)	20 (8.4)	220 (92.1)	0 (0)	96 (40.2)	4 (1.7)

^aData are presented as No. (%).

Drugs like doxycycline and tetracycline, which are among the drugs recommended in treatment guidelines, had antibiotic sensitivities of 84.7% and 81.2%, respectively. More details on drug resistance of different antibiotics are shown in Table 1. Using Chi-square and Fisher exact tests, the relationship between antibiotic resistance to doxycycline and sex was tested. This was not significant ($P = 0.622$). Using the Chi-square test, the relationship between antibiotic resistance to tetracycline and sex was tested. This was also not significant ($P = 0.073$). Using Fisher exact test, the relationship between antibiotic resistance to tetracycline and doxycycline was tested. This was significant ($P = 0.000$).

5. Discussion

Most of outbreaks, which occurred in Iran from 1977 to 2005, were caused by Ogawa serotype. In recent years, a change has been made in the number of the dominant types of outbreaks in Iran and there has been a shift between Ogawa and Inaba, as the main types of outbreaks in the country (4). Based on the main results of this study, the highest level of susceptibility was observed in the case of ciprofloxacin (99.6%) and then azithromycin (95%). Nevertheless, it is noteworthy that the levels of resistance to tetracycline (15.1%) and doxycycline (4.2%), which are mentioned as the main drugs, in most of sources, are raised (11). Hence, the large number of tetracycline resistant cases (14.7%) is remarkable. Various studies in Iran and elsewhere have reported resistance of *V. cholerae* to chloramphenicol, erythromycin, kanamycin, tetracycline and sulfamethoxazole/trimethoprim (12). There was a significant relationship between the antibiotic resistance to tetracycline and doxycycline ($P = 0.000$), i.e. if a person is resistant to one of these two antibiotics, there is a chance of more than 90% for him/her to be resistant to the other drug, as well.

Comparing the results of this study, with other studies, the following results have been achieved. The Barati and colleagues study shows that the resistance to co-trimoxazole has had an increase and there has not been a notable change in ciprofloxacin susceptibility (6). In the study of Abera, the resistance to co-trimoxazole was 100%. A high level of resistance to chloramphenicol (94%) and ampicillin (89%) was observed. In this study, all cases were suscep-

tible to doxycycline. The antibiotic susceptibility to ciprofloxacin and erythromycin in that study is almost similar to the results of our study; however, the resistance to ampicillin is greater while for tetracycline is lower (13). Data generated in the present study revealed that tetracycline, one of the main antimicrobial agents used for cholera treatment, was the most effective antibiotic. Our previous study, during the 2005 outbreak, also showed that all isolated stains of *V. cholerae* were susceptible to tetracycline. In Rahbar et al. study, all strains of *V. cholerae* that were susceptible to tetracycline, were also susceptible to doxycycline. Doxycycline is used as the first line drug for the treatment of cholera in Iran (5).

In Seas et al. study, data from these well-conducted trials show clearly that the quinolones are equally effective, as the standard therapy (multiple dose tetracycline regimen or single dose doxycycline regimen), in all clinical parameters (i.e., duration, total volume of stools). However, earlier eradication of bacterium from the stool was attained (14). In Talkington et al. study, all cases were susceptible to doxycycline and azithromycin. The susceptibility to azithromycin was similar to our study; however, in our study, susceptibility to doxycycline was lower than the result of Talkington study (15). In Pal et al. study, all the strains were similarly susceptible to ampicillin, chloramphenicol, gentamicin, ciprofloxacin and tetracycline. Ogawa biotype was resistant to furazolidone, nalidixic acid and neomycin. However, the results of the study of Pal et al. are considerably different from the results of our study, in terms of susceptibility to ampicillin and tetracycline and, considering the susceptibility to ciprofloxacin and resistance to co-trimoxazole, they were similar (16). In Rahbar et al. study, the researchers studied the *V. cholerae* epidemic, which occurred in 2005, and showed that all isolated strains were resistant to co-trimoxazole, nalidixic acid and furazolidone. On the contrary, all strains were sensitive to tetracycline, doxycycline, ciprofloxacin, erythromycin and ampicillin (11).

The results of our study show that the resistance to the doxycycline and tetracycline, which are mentioned by multiple authors as the most common antibiotic drugs for treating cholera, is increasing. However, the drug resistance to ciprofloxacin is very low. The results of this study can be used as an evidence to formulate new guidelines for the treatment of cholera.

Acknowledgments

The authors acknowledge the assistance of Masoud Moradi in the study. We would like to thank the assistance of Faezeh Mazoji in laboratory. We would like to thank the Eisa Momeni in gathering data.

Footnotes

Authors' Contribution: Conception of the research idea, study design: Parvin Mohammadi; data collection, part of laboratory work and acquisition of data and study supervision: Hojatollah Barati; analysis and interpretation of the data and critical revision: Ghobad Moradi; the draft of the manuscript: Mohammad Aziz Rasouli. All authors have read and approved the final version of the manuscript.

Financial Disclosure: The authors declare that the research was conducted in the absence of any commercial or financial support that could be considered as a potential conflict of interest.

Funding/Support: Financial support was only from the Alborz university of medical sciences, Karaj, IR Iran.

References

- Sithivong N, Morita-Ishihara T, Vongdouangchanh A, Phouthavane T, Chomlasak K, Sisavath L, et al. Molecular subtyping in cholera outbreak, Laos, 2010. *Emerg Infect Dis*. 2011;17(11):2060-2. doi: 10.3201/eid1711.110280. [PubMed: 22099098]
- Alishahi A, Imani Fooladi AA, Fallah Mehrabadi J, Mahmoodzadeh Hosseini H. Facile and Rapid Detection of *Vibrio cholerae* by Multiplex PCR Based on ompU, ctxA, and toxR Genes. *Jundishapur J Microbiol*. 2013;6(10):7933. doi: 10.5812/jjm.7933.
- Kumar P, Jain M, Goel AK, Bhadauria S, Sharma SK, Kamboj DV, et al. A large cholera outbreak due to a new cholera toxin variant of the *Vibrio cholerae* O1 El Tor biotype in Orissa, Eastern India. *J Med Microbiol*. 2009;58(Pt 2):234-8. doi: 10.1099/jmm.0.002089-0. [PubMed: 19141742]
- Msyamboza KP, Kagoli M, M'Bang'ombe M, Chipeta S, Masuku HD. Cholera outbreaks in Malawi in 1998-2012: social and cultural challenges in prevention and control. *J Infect Dev Ctries*. 2014;8(6):720-6. doi: 10.3855/jidc.3506. [PubMed: 24916870]
- Rahbar M, Zahraei M, Omidvarnia A, Afshani MT, Glami M, Sabourian R, et al. Survey of epidemiology and bacteriology features of cholera in Iran. *Asia Pac J Trop Med*. 2010;3(1):45-7. doi: 10.1016/S1995-7645(10)60030-2.
- Barati HA, Golmohammadi A, Momeni I, Mpradi G. A cholera outbreak investigation in Karaj district in 2008. *Iran J Epidemiol*. 2010;6(3):28-34.
- Koneman EW. *Color atlas and textbook of diagnostic microbiology*. Lippincott-Raven Publishers; 1997.
- Ataee RA, Mehrabi-Tavana A. An analysis on Recent Cholera Epidemic in IR IRAN. *J Mil Med*. 2005;7(2):177-85.
- World Health Organization, Centers for Disease Control and Prevention. *Laboratory methods for the diagnosis of epidemic dysentery and cholera/Centers for Disease Control and Prevention*. Atlanta: Centers for Disease Control and Prevention; 1999. Available from: http://apps.who.int/iris/bitstream/10665/66885/1/WHO_CDS_CSR_EDC_99.8.pdf.
- Mohammadi P, Barati H, Moradi G. Study of vibrio cholera anti-biogram results in Karaj cholera outbreak in 2008.; Second seminar on Enteric Pathogens Sanandaj, Iran.; Kurdistan University of Medical Sciences. 2010.
- Rahbar M, Sabourian R, Saremi M, Abbasi M, Masoumi Asl H, Soroush M. Epidemiologic and drug Resistant Pattern of *Vibrio Cholerae* O1 Biotype El Tor, Serotype Inaba during the summer of 2005 outbreak in Iran. *J Ardabil Univ Med Sci*. 2007;7(1):41-5.
- Sedaghat M, Rahimi F, Talebi M, Pourshafie M. Serotyping, Antibiotic Susceptibility Pattern and Detection of hlyA Gene Among Cholera Patients in Iran. *Jundishapur J Microbiol*. 2012;6(1):20-3. doi: 10.5812/jjm.4709.
- Abera B, Bezabih B, Dessie A. Antimicrobial susceptibility of *V. cholerae* in north west, Ethiopia. *Ethiop Med J*. 2010;48(1):23-8. [PubMed: 20607994]
- Seas C, Gotuzzo E. Cholera: Overview of epidemiologic, therapeutic, and preventive issues learned from recent epidemics. *International J Infect Dis*. 1996;1(1):37-46. doi: 10.1016/S1201-9712(96)90076-4.
- Talkington D, Bopp C, Tarr C, Parsons MB, Dahourou G, Freeman M, et al. Characterization of toxigenic *Vibrio cholerae* from Haiti, 2010-2011. *Emerg Infect Dis*. 2011;17(11):2122-9. doi: 10.3201/eid1711.110805. [PubMed: 22099116]
- Pal BB, Khuntia HK, Samal SK, Das SS, Chhotray GP. Emergence of *Vibrio cholerae* O1 biotype El Tor serotype Inaba causing outbreaks of cholera in Orissa, India. *Jpn J Infect Dis*. 2006;59(4):266-9. [PubMed: 16936349]