

# Antibiotic Resistance Pattern and Distribution of *pslA* Gene Among Biofilm Producing *Pseudomonas aeruginosa* Isolated From Waste Water of a Burn Center

Shiva Emami,<sup>1</sup> Iraj Nikokar,<sup>1\*</sup> Yusuf Ghasemi,<sup>1</sup> Monireh Ebrahimpour,<sup>1</sup> Hadi Sedigh Ebrahim-Saraie,<sup>1</sup> Afshin Araghian,<sup>1</sup> Sobhan Faezi,<sup>1</sup> Mojtaba Farahbakhsh,<sup>1</sup> and Abdolhalim Rajabi<sup>2</sup>

<sup>1</sup>Department of Biotechnology, Laboratory of Microbiology and Immunology of Infectious Diseases, Para Medicine Faculty, Guilan University of Medical Sciences, Rasht, IR Iran

<sup>2</sup>Department of Epidemiology, Faculty of Public Health, Iran University of Medical Sciences, Tehran, IR Iran

\*Corresponding author: Iraj Nikokar, Department of Biotechnology, Laboratory of Microbiology and Immunology of Infectious Diseases, Para Medicine Faculty, Guilan University of Medical Sciences, Rasht, IR Iran. Tel: +98-1425237070, Fax: +98-1425237171, E-mail: Nikokariraj@yahoo.com; Nikokariraj@gums.ac.ir

Received 2014 September 15; Revised 2015 January 16; Accepted 2015 February 5.

## Abstract

**Background:** *Pseudomonas aeruginosa* is considered as a major cause of hospital-acquired infections due to its high antibacterial resistance. Biofilm formation is a well-known pathogenic mechanism in *P. aeruginosa* infections, since sessile bacteria are protected in an extracellular matrix of exopolysaccharide. The expression of polysaccharide synthesis locus (*pslA* gene) can be important for biofilm formation by *P. aeruginosa*.

**Objectives:** The purpose of this research was to evaluate the antibiotic resistance pattern and distribution of the *pslA* gene among biofilm-producing *P. aeruginosa* isolates obtained from waste water of Burn Centre in Guilan, Iran.

**Materials and Methods:** Fifty isolates of *P. aeruginosa* were obtained from waste water of a burn center. The *P. aeruginosa* isolates were identified using standard bacteriological procedures. Drug susceptibility test was performed by disk diffusion method for all the isolates against nine antimicrobial agents. Biofilm formation was measured by microtiter plate assay. Polymerase chain reaction (PCR) was used to identify the presence of the *pslA* gene among the isolates.

**Results:** Biofilm formation was observed in 70% of the *P. aeruginosa* isolates. The potential formation of biofilm was significantly associated with resistance to gentamicin, imipenem, tobramycin and piperacillin. In addition, the *pslA* gene only existed in biofilm-producing isolates with a frequency of 42.9% (n=15).

**Conclusions:** The findings of the present study well demonstrated that the *P. aeruginosa* biofilm-producing isolates were more resistant to the tested antibiotics. Furthermore, because of wide distribution, it seems that the *pslA* gene is associated with biofilm formation.

**Keywords:** Biofilms, Antimicrobial Drug Resistance, Burn Units, *Pseudomonas aeruginosa*

## 1. Background

Biofilm formation is considered as a main problem in infection control (1, 2). Due to recent investigations, there is strong evidence that *Pseudomonas aeruginosa* strains form multicellular seeds within sites of infection, e.g. in the lungs of patients with cystic fibrosis or on the surfaces of infected catheters or burn infections (3). *Pseudomonas aeruginosa* is a prototype organism for studying biofilm formation. The persistence of long-lasting infections caused by biofilm-forming *P. aeruginosa* isolates has created serious problems in burn hospitals and related infections are difficult to treat, even in individuals with normal immune responses (1).

Biofilm formation demonstrates a protective mode of growth which allows the bacterium to survive in different environments. (4, 5). In some studies, it has been shown that the *P. aeruginosa* strains that produce bio-

film, tolerate ceftazidime, ciprofloxacin, and tobramycin antibiotics at concentrations more than those necessary to kill planktonic bacteria (6, 7). This antibiotic resistance may be due to a variety of bacterial populations which are protected from treatment in biofilm structure (8, 9). In vitro susceptibility tests with bacterial biofilm models have shown that after treatment with antibiotics, bacteria in biofilm survive at concentrations hundreds or even a thousand times greater than the minimum inhibitory concentration of the planktonic bacteria (10).

The biofilm matrix of bacteria is composed of diverse biomolecules including polysaccharide, proteins, and even DNA (11, 12). Different types of polysaccharides can be found within the matrix: alginate, polysaccharide encoding locus (Pel), and polysaccharide synthesis locus (Psl) (13). Psl that is encoded by the *pslA* gene is a neutral-

charge exopolysaccharide, comprised of D-mannose, D-glucose, and L-rhamnose, arranged in pentasaccharide repeats and provides structural support during biofilm formation, playing a role in both cell-to-cell and cell-to-substrate attachment (14). Some previous investigations showed an important role of the *pslA* gene in primary stages of biofilm formation among *P. aeruginosa* isolates (15). The expression of the *pslA* gene restores the biofilm forming phenotype among mutant strains; so, it indicates that the *pslA* gene is required for biofilm formation by *P. aeruginosa* (16). Finally, since reporting antibiotic resistance is a common phenomenon among biofilm-producing bacteria, it was interesting to determine the probable association of this event and the presence of the *pslA* gene among multidrug-resistant isolates of *P. aeruginosa*.

## 2. Objectives

The aim of the present study was to assess biofilm formation and antibiotic resistance and identify their probable correlation with the presence of the *pslA* gene among *P. aeruginosa* isolates obtained from waste water in a new burn centre in Guilan, Iran.

## 3. Materials and Methods

### 3.1. Study Area and Bacterial Strains

This cross-sectional study was conducted from October 2012 to April 2013 on 50 isolates of *P. aeruginosa* which were collected from waste water samples from a new burn center (Velayat Burn Center, Rasht) in Guilan, Iran. The biofilm-positive and negative strains of *Staphylococcus epidermidis* (RP62A and RP62NA, respectively) were used as controls for the biofilm formation test.

### 3.2. Antibacterial Susceptibility Tests

Antibacterial susceptibility was determined by disc diffusion, as recommended by the Clinical and Laboratory Standards Institute (CLSI), using the following antibiotic discs (MAST, UK): amikacin (AK), gentamicin (GM), tobramycin (TN) ceftazidime (CAZ), cefazolin (CZ), ciprofloxacin (CIP), carbenicillin (CB), imipenem (IPM), and piperacillin (PRL) (17).

### 3.3. Biofilm Formation Assay

Biofilm formation was determined in vitro using microtiter plate assay. In brief, overnight cultures (24 hours, 37°C) were adjusted to an OD<sub>600</sub> of 0.8 and were diluted 100 folds in tryptic soy broth. Aliquots (200 µL) of each isolate suspension were then inoculated into four wells of a 96-well flat-bottomed polystyrene plate and incubated overnight at 37°C. The content of each well was washed two times with 250 µL of sterile physiological saline; the plates were severely shaken to remove all non-adherent bacteria. Then, the plate was dried with heat which helps

with the fixation of the attached biofilm. Then, each well was stained with 200 µL of safranin 0.1% v/v for 15 minutes. After washing and drying the plate, the dye bound to the adherent cells was dissolved with 200 µL of ethanol 95% per well. The optical density of the biofilms was measured at 492 nm using an ELISA reader (Stat Fax 2100, Awareness Tech Inc., USA) (18). The cut-off optical density (OD<sub>c</sub>) for the microtiter plate was defined as three standard deviations plus the mean OD of the negative control. The isolates were classified as follows (19): biofilm-negative (OD < 0.625) and biofilm-positive (OD > 0.625).

### 3.4. DNA Extraction and Molecular Assay

The genomes of all the isolates were extracted by boiling method. Previously described primers were used to check the presence of the *pslA* gene: F, 5'-CACTGGACGTCTACTCCGACGATAT-3'; R, 5'-GTTCTTGATCTGTGCAGGGTGC-3', which amplify an 1119 base-pair (bp) amplicon (20). The used primers were manufactured by Bioneer, Korea. *P. aeruginosa* PAO1 was used as positive control for the *pslA* gene. PCR assay was performed in 25 µL using the specific primers with the following time and thermal program (initial denaturation: 95°C five minutes, followed by 30 cycles of denaturation at 94°C for 30 seconds, annealing at 55°C for 30 seconds, extension at 72°C for one minute, and a final extension of 10 minutes at 72°C) in a thermocycler device (Eppendorf Mastercycler Gradient, Germany).

### 3.5. Statistical Analysis

Statistical analysis was performed using SPSS™ software, version 19.0. Chi squared and t test were used to analyze the results whenever needed. P < 0.05 was considered as the significance level.

## 4. Results

Of the 50 isolates, 35 (70%) were biofilm producers and 15 (30%) were non-biofilm producers. All the isolates were resistant to cefazolin and carbenicillin followed by 82% resistance to carbenicillin, 40% to ceftazidime, 30% to gentamicin, 28% to tobramycin, 22% to piperacillin and imipenem, 20% to ciprofloxacin, and 10% to amikacin. The comparison between the two groups of strains, biofilm and non-biofilm producers, was shown in Table 1. Biofilm producers have been more resistance to most antibiotics than non-producer groups. On the other hand, resistance to gentamicin, imipenem, tobramycin, and piperacillin was significantly higher among the biofilm-producing isolates than the non-producing ones (P < 0.05). The PCR results showed that the *pslA* gene was present in 15 isolates of biofilm producers.

Interestingly, none of the biofilm-negative isolates contained the *pslA* gene. The differences in the presence of the *pslA* gene between the two groups were statically significant (P < 0.001). The results of PCR reaction for the *pslA* gene are presented in Figure 1.


**Table 1.** The Resistance of Two Groups of Isolates to Tested Antibiotics<sup>a,b</sup>

Antibiotics	Biofilm Producer (n = 35)	Biofilm Non-producer (n = 15)	P Value
Gentamicin	13 (37.1)	2 (13.3)	0.046 <sup>c</sup>
Imipenem	10 (28.6)	1 (6.7)	0.043 <sup>c</sup>
Ceftazidime	16 (45.7)	4 (26.7)	0.10
Amikacin	5 (14.3)	0	0.61
Ciprofloxacin	7 (20)	3 (20)	1.0
Piperacillin	11 (31.4)	0	0.007 <sup>c</sup>
Tobramycin	13 (37.1)	1 (6.7)	0.014 <sup>c</sup>
<i>pslA</i> gene presentation	15 (42.9)	0	0.001

<sup>a</sup>The two groups of biofilm producers were compared with the biofilm negative group.

<sup>b</sup>Data are presented as No. (%).

<sup>c</sup>P < 0.05.

**Figure 1.** A Representative Polymerase Chain Reaction Gel Image for the *pslA* Gene

Lane 1, 50 bp DNA marker; lane 2, negative control; lane 3, positive control; lane 4, 6, 8, and 9, positive samples; lane 5 and 7, negative samples.

## 5. Discussion

In recent years, despite offering a variety of antibiotics with anti-*Pseudomonas* activity, this organism with the acquisition of drug resistance genes still causes severe infections in burns (11). Bacterial biofilm communities are introduced as one of the important ways for the acquisition of resistance genes (21). We found that the majority of our *P. aeruginosa* isolates were capable to produce biofilm (70%) and subsequently showed high level of antibiotic resistance. Previously, closest to our findings, Gotaslo et al. in a hospital survey from Tabriz, Iran, reported that 79% of their clinical *P. aeruginosa* strains were biofilm producers (22). Jabalameli et al. from capital of Iran, Tehran, documented biofilm formation in more than 96% of the *P. aeruginosa* isolates collected from burn patients (23). Biofilm-related infections of *P. aeruginosa* are of particular clinical importance in skin burns and lead to chronic wounds with long healing time (24, 25).

One of the most important characteristics of bacterial biofilm is tolerance to antibiotics and the host immune system components. Therefore, the possibility of infection recurrence is an important clinical consequence of biofilm-related infections (15, 26). Multidrug resistance correlates with the ability to form biofilm on abiotic and biological surfaces (27). In our investigation, drug-resistant isolates existed in both biofilm-positive and negative groups, but most of them were significantly associated with the biofilm group. In support of our findings, Drenkard et al. found that the antibiotic-resistant variants of *P. aeruginosa* had high ability to form biofilm both in vivo and in vitro (28). In some similar studies it was documented that isolates recovered from waste waters had notable antibiotic resistance (29, 30). The transferring of such high-resistance isolates can occur in environments such as groundwater or somehow in healthcare centers, where it becomes a potential risk for the human health (29).

We found that the *pslA* gene only existed in biofilm-producing isolates; it seems that this gene is one of the most critical factors for biofilm formation in *P. aeruginosa*. To best of our knowledge, the present study was the first report from Iran showing the association of the *pslA* gene with the ability of biofilm formation in *P. aeruginosa* isolates recovered from waste water. To evaluate the essential role of *pslA* in biofilm formation, Overhage et al. generated a nonpolar isogenic *pslA* knockout mutant of *P. aeruginosa*. They found that this *pslA* knockout mutant was impaired in attachment and biofilm formation and the mutant showed about 30% less attachment to tissue culture plates than the respective wild type (16). In another study by Ghafoor et al. they found that *pslA* mutant was still able to form biofilm, but this biofilm was flat and much more compact than the biofilm formed by all other studied mutants, and both live and dead cells were present in this biofilm (31). These results showed that the *pslA* gene was an important factor to form biofilm. However, since this gene was not found in all of our biofilm-producing isolates, it seems there might be other genes or

factors that played role in biofilm formation.

As a preliminary study to determinate the role of the *pslA* gene in biofilm formation of *P. aeruginosa*, our work had a number of limitations. Apart from the limited sample size, generalizing the results of waste water isolates as clinical isolates can be doubtful. This requires a separate sampling of clinical isolates from hospital and investigates the molecular relationship between the isolates in future studies. Beside the limitations, the present study showed that the *P. aeruginosa* biofilm producing isolates were more resistance to antibiotics, especially to tobramycin, piperacillin, gentamicin, and imipenem. In addition, it seems that the *pslA* gene had association with biofilm formation, since it was widely distributed among the biofilm-producing isolates. However, since this gene was not found in all the biofilm producers, perhaps there were other genes or factors that played role in forming biofilm. Therefore, we should consider other genetic and phenotypic factors as well, which afford for future studies.

## Footnotes

**Authors' Contribution:** Study concept and design: Shiva Emami and Iraj Nikokar; sampling: Monireh Ebrahimpour; phenotypic detection: Shiva Emami, Afshin Araghian and Mojtaba Farahbakhsh; molecular detection: Yusuf Ghasemi; drafting of the manuscript: Shiva Emami, Iraj Nikokar and Sobhan Faezi; critical revision: Iraj Nikokar, Hadi Sedigh Ebrahim-Saraie and Sobhan Faezi; statistical analysis: Abdolhalim Rajabi; study supervision: Iraj Nikokar.

**Funding/Support:** This study was supported by Guilan University of Medical Sciences.

## References

1. Hoiby N, Fomsgaard A, Jensen ET, Johansen HK, Kronborg G, Pedersen SS, et al. In: The Immune Response to Bacterial Biofilms. Lappin-Scott HM, Costerton JW, editors. Cambridge: Cambridge University Press; 1995.
2. Imani Fooladi AA, Aghelimsour A, Nourani MR. Evaluation of the Pathogenesis of *Pseudomonas aeruginosa*'s Flagellum Before and After Flagellar Gene Knockdown by Small Interfering RNAs (siRNA). *Jundishapur J Microbiol*. 2013. doi:10.5812/jjm.5401
3. Deretic V. In: *Pseudomonas aeruginosa* Infections. Nataro JP, Blaser MJ, Cunningham-Rundles S, editors. Washington, DC: American Society for Microbiology Press; 2000.
4. Nivens DE, Ohman DE, Williams J, Franklin MJ. Role of alginate and its O acetylation in formation of *Pseudomonas aeruginosa* microcolonies and biofilms. *J Bacteriol*. 2001;183(3):1047-57. doi: 10.1128/JB.183.3.1047-1057.2001. [PubMed: 11208804]
5. Sarkisova S, Patrauchan MA, Berglund D, Nivens DE, Franklin MJ. Calcium-induced virulence factors associated with the extracellular matrix of mucoid *Pseudomonas aeruginosa* biofilms. *J Bacteriol*. 2005;187(13):4327-37. doi: 10.1128/JB.187.13.4327-4337.2005. [PubMed: 15968041]
6. Anwar H, Costerton JW. Enhanced activity of combination of tobramycin and piperacillin for eradication of sessile biofilm cells of *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother*. 1990;34(9):1666-71. [PubMed: 2126686]
7. Moriarty TF, Elborn JS, Tunney MM. Effect of pH on the antimicrobial susceptibility of planktonic and biofilm-grown clinical *Pseudomonas aeruginosa* isolates. *Br J Biomed Sci*. 2007;64(3):101-4. [PubMed: 17910277]
8. Balaban NQ, Merrin J, Chait R, Kowalik L, Leibler S. Bacterial persistence as a phenotypic switch. *Science*. 2004;305(5690):1622-5. doi: 10.1126/science.1099390. [PubMed: 15308767]
9. Gefen O, Balaban NQ. The importance of being persistent: heterogeneity of bacterial populations under antibiotic stress. *FEMS Microbiol Rev*. 2009;33(4):704-17. doi: 10.1111/j.1574-6976.2008.00156.x. [PubMed: 19207742]
10. Ceri H, Olson ME, Stremick C, Read RR, Morck D, Buret A. The Calgary Biofilm Device: new technology for rapid determination of antibiotic susceptibilities of bacterial biofilms. *J Clin Microbiol*. 1999;37(6):1771-6. [PubMed: 10325322]
11. Stewart PS, Costerton JW. Antibiotic resistance of bacteria in biofilms. *Lancet*. 2001;358(9276):135-8. [PubMed: 11463434]
12. Costerton JW, Lewandowski Z, Caldwell DE, Korber DR, Lappin-Scott HM. Microbial biofilms. *Annu Rev Microbiol*. 1995;49:711-45. doi: 10.1146/annurev.mi.49.100195.003431. [PubMed: 8561477]
13. Franklin MJ, Nivens DE, Weadge JT, Howell PL. Biosynthesis of the *Pseudomonas aeruginosa* Extracellular Polysaccharides, Alginate, Pel, and Psl. *Front Microbiol*. 2011;2:167. doi: 10.3389/fmicb.2011.00167. [PubMed: 21991261]
14. Billings N, Millan M, Caldara M, Rusconi R, Tarasova Y, Stocker R, et al. The extracellular matrix Component Psl provides fast-acting antibiotic defense in *Pseudomonas aeruginosa* biofilms. *PLoS Pathog*. 2013;9(8):e1003526. doi: 10.1371/journal.ppat.1003526. [PubMed: 23950711]
15. Ma L, Jackson KD, Landry RM, Parsek MR, Wozniak DJ. Analysis of *Pseudomonas aeruginosa* conditional psl variants reveals roles for the psl polysaccharide in adhesion and maintaining biofilm structure postattachment. *J Bacteriol*. 2006;188(23):8213-21. doi: 10.1128/JB.01202-06. [PubMed: 16980452]
16. Overhage J, Schemionek M, Webb JS, Rehm BH. Expression of the psl operon in *Pseudomonas aeruginosa* PAO1 biofilms: PslA performs an essential function in biofilm formation. *Appl Environ Microbiol*. 2005;71(8):4407-13. doi: 10.1128/AEM.71.8.4407-4413.2005. [PubMed: 16085831]
17. Noyal MJ, Menezes GA, Harish BN, Sujatha S, Parija SC. Simple screening tests for detection of carbapenemases in clinical isolates of nonfermentative Gram-negative bacteria. *Indian J Med Res*. 2009;129(6):707-12. [PubMed: 19692754]
18. Stepanovic S, Vukovic D, Dakic I, Savic B, Svabic-Vlahovic M. A modified microtiter-plate test for quantification of staphylococcal biofilm formation. *J Microbiol Methods*. 2000;40(2):175-9. [PubMed: 10699673]
19. Dheepa M, Vinitha LR, Appalaraju B. Comparison of biofilm production and multiple drug resistance in clinical isolates of *Acinetobacter baumannii* from a tertiary care hospital in South India. *Int J Pharm Biomed Sci*. 2011;2(4):103-7.
20. Hou W, Sun X, Wang Z, Zhang Y. Biofilm-forming capacity of *Staphylococcus epidermidis*, *Staphylococcus aureus*, and *Pseudomonas aeruginosa* from ocular infections. *Invest Ophthalmol Vis Sci*. 2012;53(9):5624-31. doi: 10.1167/iovs.11-9114. [PubMed: 22736609]
21. Beceiro A, Tomas M, Bou G. Antimicrobial resistance and virulence: a successful or deleterious association in the bacterial world? *Clin Microbiol Rev*. 2013;26(2):185-230. doi: 10.1128/CMR.00059-12. [PubMed: 23554414]
22. Gottaslo R, Salahi B. Effects of Oxygen on In-vitro Biofilm Formation and Antimicrobial Resistance of *Pseudomonas aeruginosa*. *Pharmaceutical Sci*. 2013;19(3):96-9.
23. Jabalameli F, Mirsalehian A, Khoramian B, Aligholi M, Khoram-rooz SS, Asadollahi P, et al. Evaluation of biofilm production and characterization of genes encoding type III secretion system among *Pseudomonas aeruginosa* isolated from burn patients. *Burns*. 2012;38(8):1192-7. doi: 10.1016/j.burns.2012.07.030. [PubMed: 22995427]
24. Nikokar I, Tishayar A, Flakiyan Z, Alijani K, Rehana-Banisaeed S, Hossinpour M, et al. Antibiotic resistance and frequency of class I integrons among *Pseudomonas aeruginosa*, isolated from burn patients in Guilan, Iran. *Iran J Microbiol*. 2013;5(1):36-41. [PubMed: 23466812]
25. Davis SC, Ricotti C, Cazzaniga A, Welsh E, Eaglstein WH, Mertz PM. Microscopic and physiologic evidence for biofilm-associated wound colonization in vivo. *Wound Repair Regen*. 2008;16(1):23-9.

- doi:10.1111/j.1524-475X.2007.00303.x. [PubMed:18211576]
26. Whiteley M, Bangera MG, Bumgarner RE, Parsek MR, Teitzel GM, Lory S, et al. Gene expression in *Pseudomonas aeruginosa* biofilms. *Nature*. 2001;**413**(6858):860-4. doi: 10.1038/35101627. [PubMed:11677611]
  27. Gaddy JA, Actis LA. Regulation of *Acinetobacter baumannii* biofilm formation. *Future Microbiol*. 2009;**4**(3):273-8. doi: 10.2217/fmb.09.5. [PubMed:19327114]
  28. Drenkard E, Ausubel FM. *Pseudomonas* biofilm formation and antibiotic resistance are linked to phenotypic variation. *Nature*. 2002;**416**(6882):740-3. doi: 10.1038/416740a. [PubMed:11961556]
  29. Bolaji AS, Akande IO, Iromini FA, Adewoye SO, Opasola OA. Antibiotic resistance pattern of bacteria spp isolated from hospital waste water in Ede South Western, Nigeria. *Euro J Exp Bio*. 2011;**1**(4):66-71.
  30. Basu S, Dastidar SG, Mukhopadhyay S, Gangopadhyay A. Utilization of Bulk Drugs by a Highly Antibiotic Resistant Microorganism Isolated from Waste Water of a Bulk Drug Industry. *J Pharm Sci Tech*. 2012;**2**(1):41-6.
  31. Ghafoor A, Hay ID, Rehm BH. Role of exopolysaccharides in *Pseudomonas aeruginosa* biofilm formation and architecture. *Appl Environ Microbiol*. 2011;**77**(15):5238-46. doi: 10.1128/AEM.00637-11. [PubMed:21666010]

Archive of SID