Published online 2015 November 14.

**Research Article** 

# Comparative Proteomics Analysis of Two Strains of *Neisseria meningitidis* Serogroup B and *Neisseria lactamica*

Raheleh Sheikhi, <sup>1</sup>Mansour Amin, <sup>2,3</sup>Maryam Hamidinia, <sup>4</sup>Mohammad Ali Assarehzadegan, <sup>5</sup> Soodabeh Rostami, <sup>6</sup> and Zahra Mojtahedi <sup>4,\*</sup>

<sup>1</sup>Department of Microbiology, Guilan University of Medical Sciences, Rasht, IR Iran

2Health Research Institute, Infectious and Tropical Diseases Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, IR Iran

<sup>3</sup>Department of Microbiology, Ahvaz, Iludishapur University of Medical Sciences, Ahvaz, IR Iran

<sup>4</sup>Shiraz Institute for Cancer Research, Shiraz University of Medical Sciences, Shiraz, IR Iran <sup>5</sup>Department of Immunology, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, IR Iran

6 Infectious Diseases and Tropical Medicine Research Center, Isfahan University of Medical Sciences, Isfahan, IR Iran

\*Corresponding author: Zahra Mojtahedi, Shiraz Institute for Cancer Research, Shiraz University of Medical Sciences, P. O. Box: 7134845794, Shiraz, IR Iran. Tel/Fax: +98-7112303687, E-mail: mojtahedizahra@hotmail.com

Received 2014 November 9; Revised 2015 March 6; Accepted 2015 March 30.

#### Abstract

**Background:** Antigenic similarities between *Neisseria lactamica* as a commensal species and *N. meningitidis* serogroup B (NmB) as an important cause of meningitis infection have been considered for the development of an effective vaccine based on their common proteins to prevent life-threatening bacterial meningitis.

**Objectives:** The main aims of this study were to determine whole proteome profiles of *N. lactamica* strains and to compare them with whole proteome profile of a reference strain of NmB for identification of some of common proteins between the two species.

**Materials and Methods:** We compared the whole proteomic profiles of *N. lactamica* strains and a reference strain of NmB. Lysates from bacterial strains were resolved by two-dimensional gel electrophoresis (2-DE), followed by Coomassie Brilliant blue staining. Some of the protein spots were excised from the gel and subjected to matrix-assisted laser desorption/ionization-tandem time-of-flight mass spectrometry (MALDI-TOF/TOF MS) analysis.

**Results:** The analysis of Coomassie-stained gels using ImageMaster 2D Platinum software identified approximately 800 reproducible protein spots in the range of pl 4.5 - 9.5 and  $M_{\rm r}$  of 8 - 100 kDa for each 2-DE gel of the studied bacterial strains. By comparing proteome maps of 2-DE gels, more than 200 common protein spots were recognized between the two species. Forty-eight common protein spots between the studied bacterial strains were identified by MALDI-TOF/TOF-MS. The results indicated that among the protein spots identified by MOLDI-TOF/TOF mass spectrometry, the groups of proteins included cell surface, energy metabolism, amino acid transport and metabolism, coenzyme metabolism, defense, multifunctional cellular processes, DNA, RNA and protein synthesis, ribosomal structure, regulatory functions, replication, transcription, translation, unknown and hypothetical proteins with unknown function. We found that *N. lactamica* strains have a proteome profile somewhat similar to each other and slightly different with NmB.

**Conclusions:** These results show the usefulness of proteome analysis in successful identification of the common proteins between *N. lactamica* strains and NmB. This proteomics analysis is the starting point in the path of knowledge development about whole proteome profiles of *N. lactamica* strains.

Keywords: Proteomics, Mass Spectrometry, Two-Dimensional Gel Electrophoresis, Neisseria meningitidis Serogroup B, Neisseria lactamica

## 1. Background

*Neisseria meningitidis* serogroup B (NmB) is an encapsulated Gram-negative organism and the cause of lifethreatening bacterial meningitis worldwide, especially among infants and young adults (1-3). Unlike the other serogroups of *N. meningitidis* (A, C, Y and W135), the B capsular polysaccharide serogroup is similar to sialic acid of the foetal neural cell adhesion molecule (NCAM) and is poorly immunogenic; thus, there is no effective vaccine based on capsular polysaccharides against NmB (4-6). Therefore, to advance effective vaccines against NmB, outer membrane vesicle (OMV) proteins have been considered. Outer membrane vesicle-based vaccines have been used locally based on strain-specific OMV of NmB but the usage of these vaccines has been decreased by a variety of NmB strains, short period of immunogenicity, and lack of immune responses in young children (2, 5, 6).

Copyright @ 2015, Ahvaz Jundishapur University of Medical Sciences. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

Among nonpathogenic Neisseria species, N. lactamica shows close antigenic similarities with N. meningitidis. N. lactamica is a harmless commensal species and is more frequently colonized in the upper respiratory tract of infants and young children when compared to adults; its colonization rates decrease with increasing age (7-9). Previous studies have shown that development of crossprotective immunity against N. meningitidis will be seen in people who are carriers of *N. lactamica*. Therefore, it has been proposed that colonization with N. lactamica during early childhood will protect the host against colonization with *N. meningitidis* by natural immunity, which is thought to be due to shared common antigens (5, 7, 8, 10-12). Proteomics, as a key technology, is being widely used in the study of proteins using two-dimensional gel electrophoresis (2-DE), followed by mass spectrometric (MS) protein identification (13). With an aim to carry out *N. meningitidis* proteomics, Mignogna et al. published the proteome map of the NmB serogroup B strain MC58 (14).

Investigation of the proteomic profiles of the N. lactamica strains is rarely performed. Some proteomic investigations have been devoted to analyzing and comparing common outer-membrane vesicle (OMVs) proteins (12, 15) and OMVs vaccine antigens (16) between NmB and N. lactamica. Previous studies have reported that cytoplasmic proteins, in addition to outer membrane proteins, may be involved in immune responses, (17, 18). However, the whole proteome profiles of N. lactamica isolates are not well understood. Also, no comparison has been done on the cytoplasmatic and periplasmic proteome of *N. lactamica* and NmB, so far. Therefore, this proteomic approach is the first whole proteome analysis of *N. lactamica* strains. This study is also the first comparative whole proteome analysis of the N. lactamica and NmB strains for the comparison of their common cytoplasmic and membrane proteins using proteomics approaches.

## 2. Objectives

The main aims of this study were to determine whole proteome profiles of *N. lactamica* strains and to compare them with whole proteome profile of a reference strain of NmB, for identification of common cytoplasmic and membranous proteins between the proteins of the two species by two-dimensional electrophoresis (isoelectric focusing (IEF)/SDS-PAGE) (2DE) and subsequent matrix-assisted laser desorption/ionization-tandem time-of-flight mass spectrometry (MALDI-TOF/TOF-MS).

## 3. Materials and Methods

#### 3.1. Bacterial Strains and Growth Conditions

Four *N. lactamica* strains were used in this study. Three *N. lactamica*, Nl1, Nl2 and Nl3 were isolated from the

oropharynx of healthy young children in Ahvaz, southwest of Iran (ethics number: ETH609). In our study, N. lactamica isolates, as gram-negative and oxidase-positive diplococci, were identified by growth on modified thaver-martin (MTM agar) medium (Oxoid, Basingstoke, UK) (19), O-Nitrophenyl-B-D-Galactopyranoside disc (ONPG) test and polymerase chain reaction (PCR) for the presence of the *pdhC* gene (9). Identification of *N*. lactamica strains may be confirmed by detecting betagalactosidase in an enzyme substrate test (16). Therefore, B-galactosidase activity was tested with ONPG (Rosco Diagnostica, Denmark), as the substrate. O-Nitrophenyl-B-D-Galactopyranoside disc test was performed according to the instructions of the manufacturer. The ONPG-positive isolates were yellow after incubation at 37°C for up to one hour (20). Neisseria lactamica ATCC 23970 and NmB ATCC 13090 were used as reference strains and obtained from the American type culture center, London, United Kingdom. All the bacteria were grown to confluence on Gonococcal (GC) agar (Oxoid, Basingstoke, UK) plus supplements at 37°C in a humidified atmosphere containing 5% CO<sub>2</sub> (14, 21).

## 3.2. Whole-Cell Extract Preparation

Whole-cell protein extraction method was performed according to Bernardini (21). Briefly, the bacteria were harvested from three plates, resuspended in 2 mL of PBS plus an EDTA-free protease inhibitor cocktail (Roche, Mannheim, Germany), centrifuged at 8,000 × g (Eppendorf, Germany), for ten minutes at 4°C, and washed twice with the same buffer. The cell pellets were resuspended in 1.0 mL of reswelling lysis buffer: 7 M urea, 2 M thiourea, 2% (w/v) 3-((3-Cholamidopropyl)-dimethylammonio)-1-proane sulphonate CHAPS, 2% (w/v) amidosulfobetaine-14 (ASB-14) (Sigma-Aldrich, USA), 2% (v/v) 3 - 10 Non-Linear (NL) Immobilized pH Gradient (IPG) buffer, 2 mM Tributylphosphine solution (TBP) (Sigma-Aldrich, USA) and 65 mM Dithiothreitol (DTT). Lysates were added with 1000 U of benzonase solution (Sigma-Aldrich, USA) and 2 mM MgSO<sub>4</sub> (Merck, Germany) and incubated at room temperature for 30 minutes while shaking. The cell extracts were then centrifuged at 12,000  $\times$  g (Eppendorf, Germany) for two hours at 15°C and the supernatant was collected. The soluble protein extracts were stored at -80°C in single use aliquots. The total protein concentration was determined by the Bradford assay (22). Urea, thiourea, CHAPS, Immobilised pH Gradient (IPG) buffer pH 3 - 10 non-linear (NL), and dithiothreitol (DTT) were obtained from GE Healthcare Biosciences, USA.

#### 3.3. Two-Dimensional Gel Electrophoresis (2-DE)

First dimension Isoelectric Focusing (IEF) was performed using the PROTEAN IEF Cell (Bio-Rad, USA). Approximately, 500 µg of protein extracts from each sample was brought to a final volume of 440  $\mu$ l with reswelling buffer and a small amount of bromophenol blue (Merck, Germany) (21, 23). Next, proteins were adsorbed by active rehydration onto an Immobiline Dry-Strip (18 cm, pH 3 - 10 NL gradient; GE Healthcare, USA) by applying 50 V at 20°C for 16 - 18 hours. All IEFs were carried out at 20°C and focused at 10,000 V for 20 hours. Linear voltage was increased from 250 V to 5000 V during the first two hours, from 5000 V to 10,000 V for 4.5 hours until 70,000 V/hour was obtained at a maximum of 10.000 V. Before the second dimension, strips were equilibrated for 15 minutes in equilibration buffer (6 M urea, 30% v/v glycerol) (Merck, Germany), 2% w/v Sodium Dodecyl Sulfate (SDS) (GE Healthcare, USA), 0.05 M Tris-HCl (GE Healthcare, USA) pH 6.8) containing 65 mM DTT. This step was followed by a 15-minute incubation in equilibration buffer plus 2.5% (w/v) iodoacetamide (GE Healthcare. USA). The second dimension was run on a 12% acrylamide gradient gel ( $18 \times 20$  cm  $\times 1.5$  mm) with a constant current of 20 mA, using a Mini-PROTEAN 3 Cell (Bio-Rad, USA), until the dye reached the bottom of the gel. The gels were stained with colloidal Coomassie brilliant blue G-250 (Bio-Rad, USA), as previously described (24). The 2-DE was carried out in triplicates for each bacterial strain.

#### 3.4. Image Analysis

The gels were scanned using a computing densitometer (GS-800 Imaging Densitometer, Bio-Rad, USA) at 400 dpi resolution and analyzed using the ImageMaster 2D Platinum software, version 7 (Melanie 7) (Swiss Institute of Bioinformatics, Switzerland). Spot quantification was determined according to the percentage volume (relative volume). Approximately, 800 reproducible protein spots were detected by comparing three replicas of the gels for each *Neisserial* strain. The reference image was set as a gel with the highest number of detected spots. Using match analysis, we matched the common spots automatically. The detected spots and matches were edited manually if necessary.

## 3.5. Matrix-Assisted Laser Desorption/Ionization-Tandem Time-of-Flight Mass Spectrometry Analysis

Protein spots, stained with colloidal Coomassie brilliant blue G-250, were cut from the gel using a pipette tip in a free keratin environment. The punched spots were transferred to microcentrifuge tubes (Greiner, Germany). Then, the spots were sent to the bioscience technology facility, department of biology, University of York, Wentworth Way, York, YO10 5DD, UK, for trypsin ingel digestion and MALDI-TOF/TOF-MS analysis. Furthermore, MALDI-TOF/TOF-MS was performed using a Bruker ultraflex III MALDI-TOF/TOF (Bruker Daltonics, Bremen, Germany).

#### 3.6. Database Search

The monoisotopic peptide masses created by MALDI-TOF/TOF MS were searched against the NCBInr protein database in the matrix science data base (MSDB) 20131226 and 20140122 using the MASCOT program (www.matrixscience.com) in the taxonomy group of Bacteria. A maximum of one trypsin missed cleavage per peptide, a peptide mass tolerance of  $\pm$  100 ppm and mass tolerance of  $\pm$  0.5 Da were allowed on the fragment. Other search parameters were carbamidomethylation of cysteine residues and oxidation of methionine residues, which were set as fixed and variable modifications, respectively. Peak intensity for each spot showed more than 60 peaks for peptide mass fingerprinting data. The matched peptides mainly demonstrated the strongest peaks. A maximum of 15 precursors were selected for fragmentation from each MS spectrum. In this study, MASCOT scores greater than 55 for all identified protein were indicated to be significant (P < 0.05). The theoretical  $M_r$  (molecular mass) and pI values (pH isoelectric) for each identified protein were calculated individually to ensure the quality of data, using the ExPASY-Compute pI-M, tool (http://web.expasy. org/compute\_pi/).

## 4. Results

### 4.1. Two-Dimensional Gel Electrophoresis Analysis

Bacterial strains were extracted in reswelling lysis buffer. Five hundred micrograms of any bacterial wholecell lysates was separated by the use of 2-DE with Immobilized pH Gradient (IPG) strips (nonlinear pH 3 - 10) in the first dimension (isoelectric focusing). Using the ImageMaster 2D Platinum software, version 7 (Melanie 7), approximately 800 reproducible distinct protein spots stained with colloidal Coomassie brilliant blue G-250 from the two dimensional gels were visualized for each bacterial strain. Reproducibility was evaluated by comparing three replicas of the gels for each bacterial strain. For each gel, approximately 75% of the gel spots were matched according to the reference gel. Two-dimensional gel electrophoresis map of the most representative gel of the whole-cell extract of NmB and *N. lactamica* strains is shown in Figure 1. As shown in this Figure, by comparing 2-DE gels between NmB and N. lactamica strains, we found some of the variations in the proteome profile of the studied bacterial strains. The difference of the proteome profile between the studied N. lactamica strains was lower than NmB and N. lactamica strains. Therefore, after image analysis, we selected some of the common protein spots between the studied Neisseria strains from the 2-DE gel related to the reference strain of *N*. *lactamica* and some of the protein spots from the 2-DE gel related to the NmB for MOLDI-TOF/TOF mass spectrometry analysis.

#### Sheikhi R et al.

Figure 1. Representative Two-Dimensional Gel Electrophoresis Proteome Map of a Total Protein Extract Stained With Colloidal Coomassie Brilliant Blue G-250



Neisseria meningitidis serogroup B ATCC 13090 (A) N. lactamica ATCC 23970 (B) N. lactamica NL1 (C) N. lactamica NL2 (D) and N. lactamica NL3 (E).

## 4.2. Identification of proteins by Matrix-Assisted Laser Desorption/Ionization-Tandem Time-of-Flight Mass Spectrometry

After image analysis, some of the protein spots were excised from the two 2-DE gels related to NmB and reference strain of *N. lactamica* and identified using MOLDI-TOF/TOF mass spectrometry. As shown in Figure 2, a total of 48 common protein spots were identified between NmB and *N. lactamica* strains, corresponding to 37 identified spot proteins in the *N. lactamica* strains and 11 identified spot proteins in NmB. Two of the protein spots (No. 7 and 38) did not significantly match using MALDI-TOF/TOF MS analysis. A list of MALDI-TOF/TOF MS-identified common protein spots between NmB and *N. lactamica* strains using the MASCOT program and the ImageMaster 2D Platinum software is reported in Table 1. As mentioned

in Table 1, identified proteins ranged from theoretical pI 4.84 (spot 41) to 9.2 (spot 26) and from theoretical  $M_r$  55 kDa (spot 23) to 15 kDa (spot 2). Other unidentified proteins were located between experimental pI 4.5-9.5 and experimental  $M_r$  of 9 - 100 kDa based on 2-DE gel image analysis. As shown in Table 1, in most identified proteins, a good agreement between theoretical and experimental pI and M<sub>r</sub> was observed. The result of the peptide mass fingerprinting for spot number 15 is shown in Figure 3. As shown in Figure 3, the database search result using MAS-COT and also protein sequence indicate the identity of the protein spot as phosphoglyceromutase. The standard used to accept protein spots identifications included the expanse of sequence coverage, the number of peptides matched, and the protein score. For most of the identified proteins, scores greater than 55 were considered significant (P < 0.05).



Figure 2. Numbered Protein Spots Indicate Proteins Identified By Matrix-Assisted Laser Desorption/Ionization-Tandem Time-of-Flight Mass Spectrometry Analysis

 Table 1. Matrix-Assisted Laser Desorption/Ionization-Tandem Time-of-Flight Mass Spectrometry Identification of Common Protein

 Spots Between Neisseria meningitidis serogroup B and Neisseria lactamica Strains

Spot	Protein Name	Cell <sup>a</sup>	M <sub>r</sub> (theor/	p/(theor/	Score	Matchedpep-	Accession No.	Sequence-	Function
I			exp)	exp)		tides		coverage, %	
1	50S ribosomal protein L10	С	17/11	8.8/7.7	272	2	gi-489807274	27	Translation
2	50S ribosomal protein L9	С	15/10	6.6/7	384	3	gi-313668457	36	Translation
3	Pilus assembly protein PilW	U	34/34	8.5/6.7	70	1	gi-313668647	4	Cell envelope (surface structures)
4	Single-stranded DNA- binding protein	С	19/17	5.7/6.1	400	5	gi-489806670	37	Synthesis and modification of DNA
5	Superoxide dismutase	PS	21/20	6.1/6.5	155	4	gi-489803560	12	Cell envelope (periplasmic proteins)
6	50S ribosomal protein L25	С	21/20	7/7.4	431	4	gi-489851556	31	Translation
7	Not identified		/20	/7.1					
8	50S ribosomal protein L25	С	21/20	7 7	187	2	gi-489851556	13	Translation
9	Acetyl-CoA carboxylase carboxyl transferase subunit	C	35/23	6.4/7.2	575	6	gi-313668288	29	Fatty acid biosynthesis
10	3-ketoacyl-ACP reductase	U(ps)	25/21	6.6/7.1	591	6	gi-313669143	34	Fatty acid biosynthesis
11	Uridylate kinase	С	25/21	6.7/7.4	458	5	gi-488172473	30	Purines, pyrimidines, nucleosides and nucleotides biosynthesis
12	FerredoxinNADP reductase	С	29/21	5.7/5.4	239	3	gi-15676930		
12	Energy metabolism (respiration)								
13	Enoyl-ACP reductase	CM	27/21	5.6/5.7	312	3	gi-518101971	14	Fatty acid biosynthesis
14	CRISPR-associated protein Csh2	С	32/22	5.8/6.1	590	7	gi-489804777	32	Synthesis and modification of macromolecules (DNA and RNA)
15	Phosphoglyceromutase	С	25/21	6.1/6.5	610	7	gi-489805517	37	Energy metabolism (glycolysis)
16	Short-chain dehydrogenase	С	26/21	6/6.5	164	2	gi-489803682	11	Energy metabolism (respiration)
17	Thioredoxin reductase	U(ps)	33/22	5.1/3.7	412	4	gi-518101583	21	Purines, pyrimidines, nucleosides and nucleotides metabolism
18	Oxidoreductase	С	20/20	5.6/5.3	205	2	gi-489806338	15	Miscellaneous

Jundishapur J Microbiol. 2015;8(11):e25228

<sup>(</sup>A) 2-DE gel related to the NmB (B) 2-DE gel related to the reference strain of N. lactamica. Proteins corresponding to spot numbers are given in the Table 1.

Sheil	khi	R	et	al.
-------	-----	---	----	-----

19	Phosphoribosyl-amino- imidazole-succinocarbox- amide synthase	С	32/22	5.4/4.6	507	5	gi-489804157	25	Purines, pyrimidines, nucleosides and nucleotides biosynthesis
20	Hypothetical protein	С	33/22	5.5/4.8	169	2	gi-313668735	10	hypothetical protein
21	Methylenetetrahydrofo- late dehydrogenase/cyclo- hydrolase	U(ps)	30/23	5.4/4.7	447	5	gi-489804966	26	Biosynthesis of cofactor, prosthetic groups and carriers (folic acid)
22	NAD synthetase	С	30/23	5.6/4.3	301	4	gi-489803893	19	Synthesis and modification of macromolecules
23	Nutilization substance protein A	С	55/27	4.5/2.6	531	6	gi-313667982	15	Synthesis and modification of macromolecules (RNA and DNA)
24	Opa protein	OM	26/23	9.2/9.3	184	2	gi-8346997	12	Cell envelope (surface structures)
25	Competence lipoprotein	OM	30/25	8.6/7.6	250	3	gi-313668817	13	Transport/binding protein
26	ABC transporter	СМ	27/25	5.7/6.3	79	1	gi-518102710	5	Transport/binding proteins (cations)
27	Inosine-5'-monophosphate dehydrogenase	С	52/53	6.7/7.4	384	4	gi-313668309	12	Purines, pyrimidines, nucleosides and nucleotides biosynthesis
28	Inosine-5'-monophosphate dehydrogenase	С	52/53	6.7/7.2	513	6	gi-313668309	19	Purines, pyrimidines, nucleosides and nucleotides biosynthesis
29	Inosine-5'-monophosphate dehydrogenase	С	52/53	6.7/7	75	1	gi-313668309	3	Purines, pyrimidines, nucleosides and nucleotides biosynthesis
30	Peptidoglycan-binding protein LysM	U	45/47	8.3/6.9	314	3	gi-489810301	10	Cell envelope
31	Transcription termination factor rho	U(c)	38/49	6.8/6.9	98	1	gi-3421248	6	Synthesis and modification of macromolecules (RNA and DNA)
32	Glutamine ABC transport- er ATP-binding protein	СМ	26/24	5.5/6.2	65	1	gi-489804998	4	Transport/binding protein
33	Polyphosphate kinase 2	С	36/37	8.6/8.7	370	4	gi-313669185	18	Energy metabolism
34	Imidazoleglycerol- phosphate dehydratase	С	33/34	7.8/8	273	4	gi-313668038	16	Histidine metabolism
35	Acetyl-CoA carboxylase carboxyl transferase subunit	С	35/33	6.4/7.5	403	3	gi-313668288	16	Fatty acid biosynthesis
36	Porin	OM	31/34	6.7/7.1	115	1	gi-1669708	4	Cell envelope (surface structures)
37	Inorganic pyrophosphatase	С	19/20	4.8/4,7	77	1	gi-313668872	5	Lipid metabolism ( synthesis and degradation), calcium absorption and bone formation, DNA synthesis, and other biochemical transformations
38	Not identified		/26	/7.2					
39	DNA-binding response regulator	C	24/24	5.4/5.6	202	2	gi-15676500	16	modification of macromolecules (DNA)
40	Macrophage infectivity potentiator	OM	28/27	5.7/5.3	302	3	gi-15677418	14	Surface structures (virulence factor)
41	Macrophage infectivity potentiator	ОМ	28 27	5.7/5.1	384	4	gi-15677418	18	Surface structures (virulence factor)
42	Amino acid permease substrate-binding protein	PS	28/24	5.6/5	620	6	gi-121634581	29	Transport/binding protein (amino acids and amines)
43	Putative disulfide oxidoreductase	PS	25/24	5.4/4.9	181	2	gi-6900471	10	Miscellaneous
44	Amino acid ABC transporter substrate- binding protein	PS	29/25	5/4.8	634	5	gi-15677462	37	Transport/binding proteins (cations)
45	Ketol-acid reductoisomerase	С	36/37	5.6/6	579	6	gi-15677424	26	Amino acid biosynthesis
46	D-lactate dehydrogenase	С	36/35	5.7/5.8	262	4	gi-15677533	14	Energy metabolism (respiration)
47	Outer membrane protein P1	ОМ	50/46	9.3/8	255	3	gi-15676020	9	Cell envelope (surface structures)
48	Succinyl-coa synthetase subunit alpha	С	30/33	5.7/5.6	170	2	gi-488142532	10	Energy metabolism (tricarboxylic acid cycle)

<sup>a</sup>Subcellular localization; C: cytoplasmic, OM: outer membrane, PS: periplasmic, CM: cytoplasmic membrane, U: unknown. The prediction of cellular localization was obtained using the PSORTb program (cut-off value 7.5). When all of the localization scores were equally presumable, the notation U (unknown) was selected. U (parenthesis, and small case) implicates a localization, which was more presumable than others.

#### Sheikhi R et al.



According to the mass spectrometry (MS) analysis, the identified proteins included the cytoplasmic, periplasmic, cytoplasmic, and outer membrane proteins (Table 1). The predicted cellular localization was obtained using the PSORTb program v.0.3 (25) with a cut-off value of 7.5. When all of the localization scores were equally presumable, the notation U (for unknown) was selected; this indicates that the protein may have multiple localization sites.

#### 5. Discussion

The identification of the *N. lactamica* whole proteome can be used for the determination of common proteins among *N. lactamica*, as a harmless commensal species, and NmB, as an obligate human pathogen, to develop an effective vaccine based on their common proteins, to prevent life-threatening bacterial meningitis. Investigation of the proteomic profiles of the *N. lactamica* strains and their comparison with proteome profiles of NmB strains should be considered, because there are only a few studies that have been conducted in this regard. There is no information about *N. lactamica* strains wholecell proteome profile, already available for NmB strain

MC58 (14). However, some studies have been performed, analyzing the outer membrane proteins (12, 15) and outer membrane vesicle vaccine antigens (16) in *N. lactamica* and NmB. For this reason, there is great interest in identification of *N. lactamica* proteome profile. In this study, the whole-cell extracts of NmB and *N. lactamica* strains were used for comparative proteomic analysis to identify common proteins between the two species by 2-DE and MOLDI-TOF/TOF-MS analysis.

We compared three replicas of the 2-DE gels for each of the studied bacterial strain. Most of the 800 protein spots yielded in any of the 2-DE gels were located in the range of pI 4.5 - 9.5 and  $M_r$  of 8 - 100 kDa. In our study, the proteins with  $M_r$  of more than 100 kDa were not separated in 2-DE gels because previous studies have shown that some of the proteins with high molecular masses, and highly hydrophobic proteins, particularly membrane proteins, may not be separated due to solubilization problems during the Isoelectric focusing (IEF) step and may not enter the gels (26-28). In contrast to the results of this study indicating  $M_r$  of 8 - 100 kDa, Abel et al. demonstrated  $M_r$ of 16 - 80 kDa in their study (12). These studies revealed that all membrane proteins still cannot be properly solubilized and separated using a unique method. In this study the theoretical molecular masses and pl values of the identified protein spots were very close to those values observed in the 2-DE gel. In our study, as shown in Table 1, among the identified protein spots, spot No. 47 (outer membrane protein P1) with theoretical  $M_r$  of 50 kDa was the largest membranous protein. In this study, the studied N. lactamica strains had approximately similar proteome to each other in the total number of spots and the general profile. The 2-DE gels showed that whole proteome maps of N. lactamica strains and NmB were slightly different. Consistent with this study, Abel et al. reported that different strains might show different proteome profiles despite having the same protein composition (12). In this study we did not aim to carry out identification of different protein spots between the studied bacterial strains.

In the current study, most of the protein identified spots were based on  $\geq$  2 matched peptides, although some of them were based on a single matched peptide (spots No. 3, 26, 29, 31, 32, 36 and 37). However, protein scores for all the identified protein spots were higher than the significance threshold (P < 0.05). According to the results obtained from peptide mass fingerprinting, among the identified protein spot sequences, 11 and 18 spots from the 2-DE gel, related to the reference strain of *N. lactamica*, matched with proteins belonging to *N. lactamica* ATCC 23970, and *N. lactamica*, respectively. Ten protein spot sequences from the 2-DE gel, related to the NmB, matched with proteins belonging to *N. meningitidis* MC58.

More than 200 common proteins were recognized between the two species in 2-DE gels. The analysis of 48 identified protein spots by using PSORTb program revealed that 19 protein spots were membrane proteins including periplasmic, cytoplasmic and outer membrane proteins. In contrast to our study, Abel et al. identified 17 common membrane protein spots in *N. lactamica* strains (12). Also, they detected a small number of common membrane proteins between NmB and N. lactamica strains. Among the identified outer membrane proteins, outer membrane protein P1, porin, Opa and competence lipoprotein were reported as common outer membrane proteins between the two species in previous studies, with the results being in agreement with the present study (15, 16). The PSORTb program showed that some proteins may have multiple localization sites in the cell, for example peptidoglycan-binding protein LysM (spot No. 30), as a membranous protein, may be present at any location within the cell membrane. Other identified protein spots were cytoplasmic proteins.

Eleven protein spots (No. 38 - 48) from the 2-DE gel of NmB were identified, as shown in Table 1. Comparison of the 2-DE gel of two NmB and the reference strain of *N. lactamica* revealed that these 11 identified proteins were common proteins between the two bacterial strains mentioned above. Similar to our results, amongst the above-mentioned 11 identified protein spots, four spots (No. 39, 45, 47, 48) were reported by Mignogna et al. (14).

Spots No. 40, 41, 42, 43, 44, and 46 were identified only by the present study. Spot No. 38 had no significant match and was not identified. Investigation of relative volume (%) of identified protein spots revealed that most of the protein spots, but not all, had similar spot volume in 2-DE gels. Some of the protein spots (spots No. 12, 13, 14) in the 2-DE gels related to NmB have a lower spot volume than the same spots in 2-DE gels related to *N. lactamica* strains.

The results indicated that the protein spots identified by MOLDI-TOF/TOF mass spectrometry, included cell surface, energy metabolism, amino acid transport and metabolism, coenzyme metabolism, defense, multifunctional, cellular processes, DNA, RNA and protein synthesis, ribosomal structure, regulatory functions, replication, transcription, translation, unknown and hypothetical proteins with unknown function. Cellular localization of these proteins in the cytoplasm, periplasm, cytoplasmic and outer membrane is shown in Table 1. The availability of genome sequences combined with protein databases of NmB strain MC58 (14) has led to research on the meningococcal proteome for developing an effective vaccine against this important human pathogen. In this context, it is important to consider that for performance of a reliable proteomics approach for *N*. *lactamica* strains and also NmB ATCC 13090, we must have the genome sequences of these bacteria. Therefore, this study focused on comparative proteome profiles and identification of some of the common proteins between non-pathogenic strains of *N. lactamica* and a pathogenic strain of NmB.

In conclusion, we identified several common proteins between *N. lactamica* strains and NmB using 2-DE-based MS analysis. Some of our identified proteins, but not all, were in line with previous studies that used 2-DE and MS analysis. This proteomic approach provides us with earlier reports of whole proteome maps of *N. lactamica* strains. Finally, we suggest that further studies are required to compare proteomics of this concept especially when genome sequencing of *N. lactamica* is still in progress.

## Acknowledgments

We specially thank Shiraz institute for cancer research, Shiraz university of medical sciences, Shiraz, Iran for their great support. We are grateful to Hashem Zare for his excellent technical assistance.

#### Footnotes

**Authors' Contribution:**Raheleh Sheikhi contributed to all parts of the research. Mansoor Amin was involved in the bacteriological part of the study. Maryam Hamidinia performed the 2-DE analysis and mass spectrometry. Mohammad Ali Assarehzadegan performed the data analysis. Soodabeh Rostami was involved in the bacteriological part of the study. Zahra Mojtahedi contributed to all parts of the research.

**Financial Disclosure:**This study was supported by a grant from the health research institute, infectious and

tropical diseases research center, Jundishapur university of medical sciences, Ahvaz, Iran (No. 91124).

**Funding/Support:**Health research institute, infectious and tropical diseases research center, Jundishapur university of medical sciences, Ahvaz, Iran and the department of microbiology of Ahvaz Jundishapur university of medical sciences, Ahvaz, Iran.

### References

- Williams JN, Skipp PJ, O'Connor CD, Christodoulides M, Heckels JE. Immunoproteomic analysis of the development of natural immunity in subjects colonized by Neisseria meningitidis reveals potential vaccine candidates. *Infect Immun.* 2009;77(11):5080–9. doi: 10.1128/IAI.00701-09. [PubMed: 19737898]
- Davenport V, Guthrie T, Findlow J, Borrow R, Williams NA, Heyderman RS. Evidence for naturally acquired T cell-mediated mucosal immunity to Neisseria meningitidis. *J Immunol.* 2003;**171**(8):4263-70. [PubMed:14530350]
- Hsu CA, Lin WR, Li JC, Liu YL, Tseng YT, Chang CM, et al. Immunoproteomic identification of the hypothetical protein NMB1468 as a novel lipoprotein ubiquitous in Neisseria meningitidis with vaccine potential. *Proteomics*. 2008;8(10):2115–25. doi: 10.1002/ pmic.200700574. [PubMed: 18491322]
- Bennett JS, Griffiths DT, McCarthy ND, Sleeman KL, Jolley KA, Crook DW, et al. Genetic diversity and carriage dynamics of Neisseria lactamica in infants. *Infect Immun*. 2005;73(4):2424–32. doi: 10.1128/IAI.73.4.2424-2432.2005. [PubMed: 15784588]
- Finney M, Vaughan T, Taylor S, Hudson MJ, Pratt C, Wheeler JX, et al. Characterization of the key antigenic components and preclinical immune responses to a meningococcal disease vaccine based on Neisseria lactamica outer membrane vesicles. *Hum Vaccin.* 2008;4(1):23–30. [PubMed: 17921703]
- Hill DJ, Griffiths NJ, Borodina E, Virji M. Cellular and molecular biology of Neisseria meningitidis colonization and invasive disease. *Clin Sci (Lond)*. 2010;**118**(9):547–64. doi: 10.1042/CS20090513. [PubMed: 20132098]
- Li Y, Zhang Q, Winterbotham M, Mowe E, Gorringe A, Tang CM. Immunization with live Neisseria lactamica protects mice against meningococcal challenge and can elicit serum bactericidal antibodies. *Infect Immun.* 2006;**74**(11):6348-55. doi: 10.1128/ IAI.01062-06. [PubMed: 16966413]
- Tunes CF, Ferraz AS, Scola MCG, De Gaspari EN. Intranasal Delivery of Whole Cells of Neisseria Species: Study of Cross - Reactive Antigens in Rabbits. *Open Vaccine J.* 2008;1(1):13-21. doi: 10.2174/1875035400801010013.
- Evans CM, Pratt CB, Matheson M, Vaughan TE, Findlow J, Borrow R, et al. Nasopharyngeal colonization by Neisseria lactamica and induction of protective immunity against Neisseria meningitidis. *Clin Infect Dis.* 2011;**52**(1):70-7. doi: 10.1093/cid/ciq065. [PubMed: 21148522]
- Sánchez S, Troncoso G, Ferreirós CM, Criado MT. Evaluation of cross-reactive antigens as determinants of cross-bactericidal activity in pathogenic and commensal Neisseria. *Vaccine*. 2001;19(25-26):3390-8. doi: 10.1016/s0264-410x(01)00077-9. [PubMed: 11348703]
- Wong HE, Li MS, Kroll JS, Hibberd ML, Langford PR. Genome wide expression profiling reveals suppression of host defence responses during colonisation by Neisseria meningitides but not N. lactamica. *PLoS One.* 2011;6(10):e26130. doi: 10.1371/journal. pone.0026130. [PubMed: 21980401]
- Abel A, Sanchez S, Arenas J, Criado MT, Ferreiros CM. Bioinformatic analysis of outer membrane proteome of Neisseria meningitidis and Neisseria lactamica. *Int Microbiol.* 2007;10(1):5–11. [PubMed: 17407055]
- 13. Williams JN, Skipp PJ, Humphries HE, Christodoulides M,

O'Connor CD, Heckels JE. Proteomic analysis of outer membranes and vesicles from wild-type serogroup B Neisseria meningitidis and a lipopolysaccharide-deficient mutant. *Infect Immun.* 2007;**75**(3):1364-72. doi: 10.1128/IAI.01424-06. [PubMed: 17158897]

- Mignogna G, Giorgi A, Stefanelli P, Neri A, Colotti G, Maras B, et al. Inventory of the proteins in Neisseria meningitidis serogroup B strain MC58. J Proteome Res. 2005;4(4):1361-70. doi: 10.1021/ pr0500511. [PubMed:16083288]
- Sanchez S, Arenas J, Abel A, Criado MT, Ferreiros CM. Analysis of outer membrane protein complexes and heat-modifiable proteins in Neisseria strains using two-dimensional diagonal electrophoresis. J Proteome Res. 2005;4(1):91–5. doi: 10.1021/ pr049846i. [PubMed: 15707362]
- Vaughan TE, Skipp PJ, O'Connor CD, Hudson MJ, Vipond R, Elmore MJ, et al. Proteomic analysis of Neisseria lactamica and N eisseria meningitidis outer membrane vesicle vaccine antigens. *Vaccine*. 2006;24(25):5277–93. doi: 10.1016/j.vaccine.2006.03.013. [PubMed: 16682101]
- Mendum TA, Newcombe J, McNeilly CL, McFadden J. Towards the immunoproteome of Neisseria meningitidis. *PLoS One*. 2009;4(6):e5940. doi: 10.1371/journal.pone.0005940. [PubMed: 19529772]
- Ferrari G, Garaguso I, Adu-Bobie J, Doro F, Taddei AR, Biolchi A, et al. Outer membrane vesicles from group B Neisseria meningitidis delta gna33 mutant: proteomic and immunological comparison with detergent-derived outer membrane vesicles. *Proteomics.* 2006;6(6):1856–66. doi: 10.1002/pmic.200500164. [PubMed: 16456881]
- Winn WC, Koneman EW. Koneman's color atlas and textbook of diagnostic microbiology. Philadelphia: Lippincott williams & wilkins; 2006.
- Alber D, Oberkotter M, Suerbaum S, Claus H, Frosch M, Vogel U. Genetic diversity of Neisseria lactamica strains from epidemiologically defined carriers. *J Clin Microbiol.* 2001;**39**(5):1710–5. doi: 10.1128/JCM.39.5.1710-1715.2001. [PubMed: 11325979]
- Bernardini G, Renzone G, Comanducci M, Mini R, Arena S, D'Ambrosio C, et al. Proteome analysis of Neisseria meningitidis serogroup A. *Proteomics*. 2004;4(10):2893–926. doi: 10.1002/ pmic.200400946. [PubMed: 15378741]
- 22. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.* 1976;**72**(1-2):248–54. doi: 10.1016/0003-2697(76)90527-3. [PubMed: 942051]
- Williams JN, Christodoulides M, Heckels JE. Analysis of the immune response to Neisseria meningitidis using a proteomics approach. *Methods Mol Biol*. 2012;**799**:343–60. doi: 10.1007/978-1-61779-346-2\_20. [PubMed: 21993655]
- 24. Neuhoff V, Arold N, Taube D, Ehrhardt W. Improved staining of proteins in polyacrylamide gels including isoelectric focusing gels with clear background at nanogram sensitivity using Coomassie Brilliant Blue G-250 and R-250. *Electrophoresis*. 1988;**9**(6):255–62. doi:10.1002/elps.1150090603. [PubMed: 2466658]
- Gardy JL, Laird MR, Chen F, Rey S, Walsh CJ, Ester M, et al. PSORTb v.2.0: expanded prediction of bacterial protein subcellular localization and insights gained from comparative proteome analysis. *Bioinformatics*. 2005;21(5):617–23. doi: 10.1093/bioinformatics/ bti057. [PubMed: 15501914]
- Gorg A, Weiss W, Dunn MJ. Current two-dimensional electrophoresis technology for proteomics. *Proteomics*. 2004;4(12):3665–85. doi:10.1002/pmic.200401031. [PubMed: 15543535]
- 27. Bernardini G, Braconi D, Santucci A. The analysis of Neisseria meningitidis proteomes: Reference maps and their applications. *Proteomics.* 2007;7(16):2933-46. doi: 10.1002/pmic.200700094. [PubMed: 17628027]
- Gorg A, Drews O, Luck C, Weiland F, Weiss W. 2-DE with IPGs. *Electrophoresis*. 2009;**30 Suppl 1**:S122–32. doi: 10.1002/ elps.200900051. [PubMed: 19441019]

www.SID in