

# Molecular and Morphometric Characterization of *Acanthamoeba* spp. from Different Water Sources of Northwest Iran as a Neglected Focus, Co-Bordered With the Country of Iraq

Aram Khezri,<sup>1,2,\*</sup> Esmael Fallah,<sup>1,3,4</sup> Mostafa Mostafazadeh,<sup>2,5</sup> Adel Spotin,<sup>1</sup> Abbas Shahbazi,<sup>1</sup>

Mahmoud Mahami-Oskouei,<sup>1</sup> and Taimuor Hazratian<sup>1</sup>

<sup>1</sup>Department of Parasitology and Mycology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

<sup>2</sup>Students' Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran

<sup>3</sup>Immunology Research Center, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

<sup>4</sup>Department of Laboratory Medicine, Ardabil Medical Sciences Branch, Islamic Azad University, Ardabil, Iran

<sup>5</sup>Department of Biochemistry and Clinical Laboratories, Tabriz University of Medical Sciences, Tabriz, Iran

\*Corresponding author: Aram Khezri, Department of Parasitology and Mycology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran. Tel: +98-9145583850, Fax: +98-4133373745, E-mail: dr.aram66@yahoo.com

Received 2016 April 15; Revised 2016 September 25; Accepted 2016 October 08.

## Abstract

**Background:** *Acanthamoeba* spp. is a free-living opportunistic protozoan parasites, which can be found in tap, fresh and bottled mineral waters, contact lens solutions, soil etc.

**Objectives:** The present study is aimed to determine the *Acanthamoeba* spp. on the basis of their morpho-molecular aspects in different water sources of the West Azerbaijan province, Northwest of Iran.

**Methods:** In this cross-sectional study, 60 water samples were collected from rivers and tap waters during June to September 2015. The water samples were filtered through a cellulose nitrate filter and cultured on non-nutrient agar medium. The extracted DNAs were amplified and some amplicons were sequenced using partial 18S rRNA for genotyping and phylogenetic analyses.

**Results:** Twenty-seven (45%) out of 60 water samples were positive to *Acanthamoeba* spp. using both culture and morphological examinations. In addition, 24 (40%) out of 27 positive samples in culture method were confirmed by PCR to be *Acanthamoeba* spp.

**Conclusions:** A relatively high prevalence of *Acanthamoeba* spp. in rivers reflects a risk alert for threatening human health in the region. However, well hygienic status of the tap waters considering *Acanthamoeba* spp. cannot be ignored in western co-border regions of Iran-Iraq. This study can also serve as a platform for further explorations of water sources in Iran and neighboring countries.

**Keywords:** West Azerbaijan, Water resources, PCR, Iran, *Acanthamoeba*

## 1. Background

Free-living amoebae (FLA), from the genera of *Acanthamoeba*, *Balamuthia* and *Naegleria* can be found all around the world (1). *Acanthamoeba* spp., are opportunistic protozoan parasites that can be isolated from the wide variety of environments such as tap, fresh, coastal, bottled mineral waters, contact lens solutions, eyewash stations, soil, dust, sewage, ventilators and air-conditioning units (1-4). Some species of *Acanthamoeba* have direct roles as opportunistic human pathogens causing amoebic keratitis, especially in contact lens wearers and life-threatening granulomatous amoebic encephalitis (GAE) in immunosuppressed/compromised patients, such as HIV/AIDS and organ transplant recipients (5). It is shown that more than 80% of the normal human population has antibodies

against *Acanthamoeba* in their body. This is clearly indicative of that the organism often comes into contact with humans (6, 7).

*Acanthamoeba* has two forms in its life cycle, active trophozoites and resistant cysts. The double-layered coats of cysts enable resistance against disinfectants, such as chlorine compounds and antibiotics. Moreover, it can survive in a wide range of temperature variations, -2°C to 45°C (5, 8). Based on the 18S rRNA gene sequences, the genus *Acanthamoeba* is classified into 17 different genotypes (T1 - T17). The majority (90%) of infection-causing isolates belong to the T4 genotype (9, 10) and more than ninety percent (> 90%) of the *Acanthamoeba* isolates that produce keratitis belong to the T3 and T4 genotypes (11, 12). Of the genotypes (T4, T3, and T2) associated with *Acanthamoeba*

keratitis (AK), T2 and T4 have been reported in Iran's water (10, 13).

The different species of *Acanthamoeba* are classified based on the size of cysts and the numbers of arms into the three distinct morphological groups (I, II, and III); group I: with an average diameter of cysts  $\geq 18 \mu\text{m}$ , fewer than six arms and smooth ectocyst and star shape endocysts, group II: it is possible for the ectocyst and endocyst to be close together or widely separated, ectocysts may be thick or thin and endocysts may be polygonal, triangular or round with a mean diameter of less than  $18 \mu\text{m}$ , group III: ectocysts are thin and endocysts may have 3 - 5 blunt corners with the mean cyst diameter of  $< 18 \mu\text{m}$  (9, 14, 15). In recent years, remarkable focus has been dedicated to *Acanthamoeba* due to (i) increasing *Acanthamoeba* infections, (ii) the potential role of *Acanthamoeba* in the ecosystem, (iii) model organism to study the molecular biology of motility and phagocytosis resemblance to macrophages and (iv) being a host or reservoir for microbial pathogens (15).

## 2. Objectives

Although several studies have been accomplished on molecular characterization of *Acanthamoeba* spp. in different regions of Iran (16, 17) there is no similar research that has been done in the our study area, therefore, information on the occurrence and distribution of *Acanthamoeba* spp. in environmental and drinking water in West Azerbaijan province remains unclear. The aim of the present study was to determine the epidemiological status of *Acanthamoeba* spp. based on their molecular and morphometric features in water sources of the West Azerbaijan province, northwest Iran.

## 3. Methods

### 3.1. Study Area

In this cross-sectional study, 60 water samples were collected during June to September 2015 from river sources and tap water in the southern parts of the West Azerbaijan province. The water samples were collected from different parts of Simineh Rud (28 samples) and Zarineh Rud (16 samples), the two main rivers located in the West Azerbaijan province, as well as from the tap waters of the cities of Boukan (four samples), Shahindezh (four samples), Mian-doab (four samples) and Mahabad (four samples) (Figure 1).

At the time of sampling, free residual chlorine was measured by the Cl D.P.D Amcor test Kit (HACH, Germany). Each sample (1000 mL) was filled in a sterile glass bottle

and transferred immediately to the Parasitology Department of Tabriz University of Medical Sciences, Iran, within 24 hours, in room temperature and stored at the same temperature for subsequent analysis.

### 3.2. *Acanthamoeba* spp. Isolation and Measurement

All volume of each water sample (1 liter) was passed through a cellulose nitrate filter,  $0.42 \mu\text{m}$  in diameter (Millipore, Germany) with a weak vacuum (flow rate, 1.3 mL/minute). The samples were divided into two parts and filtered separately using a filtration system. One of the filters was placed directly on the culture medium (1.5% non-nutrient agar plates) and the other was shaken in Page's Amoeba Saline and then the resulting solution was centrifuged for 10 minutes at 500 rpm. The sediments were cultured on 1.5% non-nutrient agar (Merck, Germany) plates seeded with heat-killed *Escherichia coli* and then incubated at  $37^\circ\text{C}$  for up to 14 days. During the incubation period, the amoebic growth was examined by inverted microscope (Olympus, Japan) on a daily basis. The positive amoebic growth cultures were examined based on morphological characteristics for the presence of trophozoites and cysts, particularly considering their two-layered cell wall cysts.

Isolated *Acanthamoeba* spp. were transferred into axenic culture, peptone yeast extract glucose broth (PYG) (Sigma-Aldrich, USA) [0.75% proteose peptone (wt/vol), 0.75% yeast extract (wt/vol), and 1.5% glucose (wt/vol)]. The identification of *Acanthamoeba* at the genus level was performed according to the distinctive features of double-walled cysts. Microscopic slides were prepared from positive cultures and stained by Giemsa's staining method as followed: the fixed smear was immersed in Giemsa (Merck, Germany) solution (1: 45 mL of  $\text{dH}_2\text{O}$ ) in a staining container for 60 minutes. The slide was then rinsed with  $\text{dH}_2\text{O}$  (dried at  $37^\circ\text{C}$  for 60 minutes) and studied under light microscope (BH-2, Olympus Corporation, Japan) using 100X objective with scale bar. The diameter of the cysts and the numbers of arms were recorded and classified based on the Pussard and Pons classification (18). Different characters of cysts were measured by ocular micrometer, which were previously calculated by dividing ocular micrometer to stage micrometer ( $100 \times \text{objective} = 1 \mu\text{m}$  per unit space).

### 3.3. DNA Extraction

DNA extraction kit (Takapou Zist, Iran) was used to extract the total genomic DNA in the concentrated debris according to the kit protocol. The purity of extracted DNA from each sample was determined by the NanoDrop (ND-1000, Termo Fisher Scientific, USA) in the wavelength of 260 nm.



Figure 1. Map of Iran Presenting Study Locations in West Azerbaijan Province, Northwest Iran Where Located With Co Border of Iraq Country

### 3.4. Polymerase Chain Reaction

A pair of primers (JDP1-JDP2); JDP1: 5'-GGCCCAGATCGTTACCGTGAA-3' and JDP2: 5'-TCTCACAAGCTGCTAGGGGAGTCA-3' was used to amplify a fragment of approximately 500 bp *Acanthamoeba* genus-specific 18S ribosomal DNA (19, 20). PCR was performed in a 30 µL volume reaction mixer containing 1.25 U of Taq DNA polymerase (Cinnagen, Iran), 2 µL of template DNA and 0.3 mM dNTP, 20 pmol of each primer, 1.5 mM MgCl<sub>2</sub> and 10 mM PCR buffer. PCR was performed using an initial denaturation condition for 3 minutes at 94°C, 32 amplification cycles (denaturing 94°C for 45 s, annealing at 55°C for 35 s, and extension at 72°C for 1 minute) and a final elongation step for 10 minutes at 72°C. The PCR products were fractionated by gel electrophoresis on 2% agarose gel that the DNA Safe stain (Invitrogen, UK) has added to the gel solution during preparation and the amplicons were visualized under UV light in a transilluminator device.

### 3.5. 18S rRNA Sequencing and Phylogenetic Analysis


In this study, in order to get the genotyping and phylogenetic analysis, the Genetic Analyzer 3130 ABI sequenced 14 isolates of *Acanthamoeba*. Ambiguous (heterozygous) sites were coded using the standard IUPAC codes for combinations of two or more bases. Contigs from all samples were aligned and edited in consensus positions compared to GenBank sequences of all regional species using the Sequencher Tmv.4.1.4 software for PC (Gene Codes Corpora-

tion). The MEGA 5.05 software and maximum likelihood algorithm with kimura 2-parameter model for phylogenetic analysis were used. The accuracy of phylogenetic tree was evaluated by 1000 bootstrap re-sampling.

## 4. Results

The mean concentration of free residual chlorine was about 0.3 - 1 mg/Liter. Based on the morphological identification, 27 (45%) out of 60 collected water samples were positive to *Acanthamoeba* spp. using both culture and morphological examinations. Based on Pussard and Pon's criteria (18), 19 cases belonged to group II (70.4%) and 8 cases were members of group III (29.6%). It should be mentioned that co-infections with both groups was identified in one sample. The ranges of identified cysts diameters were measured  $11.1 \pm 1.02$  to  $16.4 \pm 4.76$  µm (Table 1). Giemsa staining method was used for morphological study (Figure 2). All of the positive samples were from the rivers and the amoeba was not observed in the any of the tap water samples. PCR was performed using JDP1 and JDP2 primers for small subunit ribosomal DNA and detected approximately 500 bp band on agarose gel for all positive isolates. Fifteen of the positive PCR samples (53.5%) were taken from Simineh Rud river and nine samples (56.5%) were taken from Zarineh Rud river. There was no positive sample in the studied tap water samples (Table 2).

In phylogenetic analysis, 14 out of 24 amplicons were sequenced which among them, only 13 sequences (shown in



**Figure 2.** *Acanthamoeba* Cysts in Primary Morphology Assessment by Giemsa Staining (x100)

Table 1) were explicitly identified as T4 genotype and one sequence was identified as T5. Twelve out of 13 sequences have common haplotypes (AZE1-AZE3: Figure 3) compared to reference isolate and only one sequence has new haplotype (Buk 11, Accession no. KP940446) which in position 101, T (Thymine) was replaced by C (Cytosine). *Hartmannella* spp. (KF697197) was considered as a group branch in drawing the phylogeny tree (Figure 4).

## 5. Discussion


This study indicated relatively high prevalence of *Acanthamoeba* spp. (T4) with different morphometric aspects but no more heterogeneity traits in rivers of West Azerbaijan, Northwest Iran. The high percentage of *Acanthamoeba* spp., in different environmental sources represents a sanitary risk for public health, particularly contact lens users and immunocompromised/suppressed patients (21).

In the present study, *Acanthamoeba* spp. was isolated in 45% of cultured samples, which amongst them, 88.8% were confirmed by PCR. The difference between PCR and culture method is probably due to other free living amoeba cysts except *Acanthamoeba* spp. Based on the morphological classification of Pussard and Pons (18), 18 cases of the positive cultures were classified in group II, with the most frequency in the world. Although, 8 positive cases were members of group III and one case from group I and group II. The results of this study showed that none of the samples from tap water contain *Acanthamoeba* spp., which is in contrast with some of the previous studies (4, 22). This difference, maybe, resulted from the fact that the previous studies collected their samples from a storage tank of water while our samples were obtained directly from tap water (4, 22).


Tap water is usually taken from a river or dam that it is cleaned at a purification station, such as Rand Water's, and then supplied via pipelines to taps. Therefore, it can affect the *Acanthamoeba* trophozoites. Another study on the water storage tanks in UK has shown that these water supplies promote colonization of domestic water with free-living amoebae, including *Acanthamoeba*, and therefore increases the risk of infection with the amoebae. This accounts for the significantly greater incidence of *Acanthamoeba* infection in the UK and supports advice to avoid using tap water in contact lens care routines (23).

In the present study, in tap water samples the mean concentration of free residual chlorine was about 0.3 - 1 mg/L, which can be another reason that no *Acanthamoeba* spp. was isolated from tap water samples. It has been reported that the trophozoites of *Acanthamoeba* are sensitive to free residual chlorine, while the cysts are resistant to chlorine (24). On the other hand, in normal environments such as chlorine-free water, the parasite lives as trophozoite, however in frigid environments it forms cyst. In the optimum water condition, *Acanthamoeba* lives as a trophozoite since it is susceptible to chlorine. Therefore, it has been conceived that the growth of *Acanthamoeba* in water with chlorine is difficult. Given the fact that river water is chlorine-free, rivers are a significant source of the organism. In the case of the river waters, we have isolated *Acanthamoeba* species from 27 out of 44 (45%) river water samples by cultivation and morphologic assessment; also, 24 samples (40%) were reconfirmed by the molecular method. These results show that the contamination rate of river water in this area is very high and it maybe a result from the fact that these two rivers are exposed to air and soil, as well as other infectious sources, such as sewage, wastewater and feces of animals. These factors raise the contam-





**Figure 3.** Nucleotide Sequence Alignment of 18S rRNA Gene Based on Detected Novel and Common Haplotypes in this Study Compared to Retrieved Sequences From GenBank Database



**Figure 4.** Phylogenetic Tree of *Acanthamoeba* 18S rRNA Sequences From Surface Water of Northwest Iran and Other Previous Registered Sequences of Different areas Using c Algorithm With Kimura 2 Parameter Model and 1000 Bootstrap resampling

ination rate in the river water sources. Therefore, the residents of these areas often have an indirect contact with the *Acanthamoeba* infection.

In Iran, infections with *Acanthamoeba* spp. has increased in recent years, mainly due to poor hygiene in contact lens users (4). The ten-year study on patients with amoebic keratitis has shown that the *Acanthamoeba* infection is continually rising in Iran (21). Niyati et al., from drinking and recreational water, reported *Acanthamoeba* in 17 (25.4%) of the 67 collected samples in East Azerbaijan, Northwest Iran. Most genotypes in this study were also reported as T4 (25).

Bagheri et al. (2010) reported that half of their inves-

tigated samples collected from cold and warm tap water sources of hospitals in 13 cities of Iran were harboring *Acanthamoeba* spp. (4). In another study conducted by Hooshyar et al. (2013), it is shown that 32 (80%) out of 40 water samples collected from 20 stagnant waters in squares and parks in different regions of Qazvin city, Iran, were positive for free living amoebas by the culture method. Similar to our findings, PCR reactions were positive in 14 (43.8%) out of 32 culture samples (16). Maghsood et al. (2005) isolated *Acanthamoeba* spp. from water samples by molecular methods in different cities of Iran (10). In the study conducted by Rezaeian et al. (2008), *Acanthamoeba* spp. have been recovered from 58% collected samples from a variety

**Table 1.** Morphological Observations of the Isolated *Acanthamoeba* spp. Amongst Positive Samples on Non-Nutrient Agar Culture Media<sup>a</sup>

No.	Region	Cyst Diameter, $\mu\text{m}$	Group <sup>b</sup>	Culture	PCR	Genotype
1	Gharakand <sup>c,d</sup>	11.1 $\pm$ 1.02	II	+	+	T4
2	Amirabad <sup>d</sup>	14.2 $\pm$ 0.82	II	+	+	-
3	Hamamian <sup>d</sup>	11.1 $\pm$ 1.2	III	+	+	-
4	Hossinname <sup>d</sup>	12.1 $\pm$ 0.99	II	+	+	-
5	Miandoab <sup>e</sup>	11.8 $\pm$ 1.01	III	+	+	-
6	Kosteh <sup>d</sup>	11.3 $\pm$ 1.01	III	+	+	T4
7	Gharajoz <sup>d</sup>	11.1 $\pm$ 0.95	II	+	+	-
8	Miandoabz <sup>d</sup>	15.2 $\pm$ 1.08	III	+	+	T5
9	Khankandi <sup>e</sup>	13.3 $\pm$ 1.04	II	+	+	T4
10	Malar <sup>e</sup>	12.2 $\pm$ 1.02	II	+	+	T4
11	Tazeghalla <sup>d</sup>	14.2 $\pm$ 0.97	II	+	+	T4
12	Akhtata <sup>d</sup>	12.3 $\pm$ 1.1	II	+	+	T4
13	Shahrikand <sup>d</sup>	11.5 $\pm$ 0.89	II	+	+	T4
14	Simineh <sup>d</sup>	12.3 $\pm$ 1.02	III	+	+	T4
15	Gogjali <sup>e</sup>	13.4 $\pm$ 1.37	II	+	+	-
16	Gavmishgoli <sup>e</sup>	12.5 $\pm$ 0.96	II	+	+	T4
17	Ashigolan <sup>d</sup>	14.1 $\pm$ 1.3	II	+	-	-
18	Gharaghshlagh <sup>e</sup>	16.4 $\pm$ 1.76	II	+	+	T4
19	Hajihassan <sup>e</sup>	12.2 $\pm$ 0.88	III	+	+	-
20	Ghazali <sup>e</sup>	11.7 $\pm$ 1.32	III	+	+	T4
21	Ghomghala <sup>d</sup>	12.4 $\pm$ 0.97	II	+	+	T4
22	Vosokand <sup>d</sup>	16.4 $\pm$ 5.02	II,I	+	+	-
23	Ghzi <sup>d</sup>	14.6 $\pm$ 0.99	II	+	+	-
24	Darman <sup>d</sup>	12.2 $\pm$ 1.2	II	+	-	-
25	Kanikozalah <sup>e</sup>	13.4 $\pm$ 1.03	III	+	+	-
26	Sardarabad <sup>d</sup>	11.6 $\pm$ 1.47	II	+	+	T4
27	Kanirash <sup>d</sup>	11.1 $\pm$ 0.98	II	+	-	-

<sup>a</sup> Values are expressed as mean  $\pm$  SD.  
<sup>b</sup> Pussard and Pons' criteria (1977).  
<sup>c</sup> Buk II isolate, Accession no. KP940446.  
<sup>d</sup> Samples have been taken from Simineh Rud River  
<sup>e</sup> Samples have been taken from Zarineh Rud River

**Table 2.** Frequency of *Acanthamoeba* spp. Amongst Studied Water Samples

Source		Culture	Sample No	PCR
Tap water		0	16	0
River	Simineh Rud	17	28	15
	Zarineh Rud	10	16	9
Total (%)		27 (45%)	60	24 (40%)

of ecological habitats such as tap water, soil and dust in Tehran, Iran (21).

*Acanthamoeba* spp. has been detected in 3% of samples taken from the James River in Virginia, USA (26). In the Pusan province, Korea, *Acanthamoeba* has been isolated from 5.8% of tap water samples (22). In Spain, Lorenzo-Morales et al. (2005) reported that 59.5% of tap water samples were

harboring *Acanthamoeba* (8). The observed difference in the prevalence of *Acanthamoeba* in tap water in different countries might be due to the different water treatment strategies or the accuracy of the water treatment system in each country. In this study, no more new haplotypes were found in analyzed isolates, this can be associated with the structural features of 18S rRNA gene, which indicated as a conserved nucleus marker and it's to be diploid. The findings of the present study provide the evidence for high prevalence of *Acanthamoeba* spp. in rivers, which reflects a risk alert for threatening human health in northwest Iran, where placed with the co-border of Iraq country. In addition, a well hygienic status of tap water considering *Acanthamoeba* spp. cannot be ignored in the region. This study can also serve as a platform for further explorations of water sources in Iran.

**Acknowledgments**

The authors appreciate the staff of the department of parasitology and microbiology of Tabriz University of Medical Sciences for their help.

**Footnotes**

**Authors' Contribution:** Study concept and design, Esmaeel Fallah, Aram Khezri; molecular analysis, Mostafa Mostafazadeh; analysis and interpretation of data, Abbas Shahbazi, Mahmoud Mahami Oskouei, Taimuor Hazratian; drafting of the manuscript, Aram Khezri; critical revision of the manuscript for important intellectual content, Aram Khezri. All authors read and approved the final manuscript.

**Funding/Support:** This work was supported fully by the Tabriz immunology research center, Tabriz University of Medical Sciences, Tabriz Iran, (grant no: 92/91).

**References**

- Magnet A, Galvan AL, Fenoy S, Izquierdo F, Rueda C, Fernandez Vadillo C, et al. Molecular characterization of *Acanthamoeba* isolated in water treatment plants and comparison with clinical isolates. *Parasitol Res.* 2012;**111**(1):383-92. doi: [10.1007/s00436-012-2849-2](https://doi.org/10.1007/s00436-012-2849-2). [PubMed: 22395660].
- Lorenzo-Morales J, Ortega-Rivas A, Martinez E, Khoubbane M, Artigas P, Periago MV, et al. *Acanthamoeba* isolates belonging to T1, T2, T3, T4 and T7 genotypes from environmental freshwater samples in the Nile Delta region, Egypt. *Acta Trop.* 2006;**100**(1-2):63-9. doi: [10.1016/j.actatropica.2006.09.008](https://doi.org/10.1016/j.actatropica.2006.09.008). [PubMed: 17078918].
- Chen CJ, Yu MW, Liaw YF. Epidemiological characteristics and risk factors of hepatocellular carcinoma. *J Gastroenterol Hepatol.* 1997;**12**(9-10):S294-308. [PubMed: 9407350].
- Bagheri H, Shafiei R, Shafiei F, Sajjadi S. Isolation of *acanthamoeba* spp. From drinking waters in several hospitals of iran. *Iran J Parasitol.* 2010;**5**(2):19-25. [PubMed: 22347240].

5. Huang SW, Hsu BM. Isolation and identification of Acanthamoeba from Taiwan spring recreation areas using culture enrichment combined with PCR. *Acta Trop.* 2010;**115**(3):282-7. doi: [10.1016/j.actatropica.2010.04.012](https://doi.org/10.1016/j.actatropica.2010.04.012). [PubMed: [20434423](https://pubmed.ncbi.nlm.nih.gov/20434423/)].
6. Ashford JW, Coburn KL, Fuster JM. The elgiloy microelectrode: fabrication techniques and characteristics. *J Neurosci Methods.* 1985;**14**(4):247-52. [PubMed: [4058056](https://pubmed.ncbi.nlm.nih.gov/4058056/)].
7. Chappell CL, Wright JA, Coletta M, Newsome AL. Standardized method of measuring acanthamoeba antibodies in sera from healthy human subjects. *Clin Diagn Lab Immunol.* 2001;**8**(4):724-30. doi: [10.1127/CL109018](https://doi.org/10.1127/CL109018). [PubMed: [96134](https://pubmed.ncbi.nlm.nih.gov/96134/)].
8. Lorenzo-Morales J, Ortega-Rivas A, Foronda P, Martinez E, Valladares B. Isolation and identification of pathogenic Acanthamoeba strains in Tenerife, Canary Islands, Spain from water sources. *Parasitol Res.* 2005;**95**(4):273-7. doi: [10.1007/s00436-005-1301-2](https://doi.org/10.1007/s00436-005-1301-2). [PubMed: [15678350](https://pubmed.ncbi.nlm.nih.gov/15678350/)].
9. Khan NA. Acanthamoeba: biology and increasing importance in human health. *FEMS Microbiol Rev.* 2006;**30**(4):564-95. doi: [10.1111/j.1574-6976.2006.00023.x](https://doi.org/10.1111/j.1574-6976.2006.00023.x). [PubMed: [16774587](https://pubmed.ncbi.nlm.nih.gov/16774587/)].
10. Maghsood AH, Sissons J, Rezaian M, Nolder D, Warhurst D, Khan NA. Acanthamoeba genotype T4 from the UK and Iran and isolation of the T2 genotype from clinical isolates. *J Med Microbiol.* 2005;**54**(Pt 8):755-9. doi: [10.1099/jmm.0.45970-0](https://doi.org/10.1099/jmm.0.45970-0). [PubMed: [16014429](https://pubmed.ncbi.nlm.nih.gov/16014429/)].
11. Schuster FL, Visvesvara GS. Free-living amoebae as opportunistic and non-opportunistic pathogens of humans and animals. *Int J Parasitol.* 2004;**34**(9):1001-27. doi: [10.1016/j.ijpara.2004.06.004](https://doi.org/10.1016/j.ijpara.2004.06.004). [PubMed: [15313128](https://pubmed.ncbi.nlm.nih.gov/15313128/)].
12. Stothard DR, Schroeder-Diedrich JM, Awwad MH, Gast RJ, Ledee DR, Rodriguez-Zaragoza S, et al. The evolutionary history of the genus Acanthamoeba and the identification of eight new 18S rRNA gene sequence types. *J Eukaryot Microbiol.* 1998;**45**(1):45-54. [PubMed: [9495032](https://pubmed.ncbi.nlm.nih.gov/9495032/)].
13. Niyiyati M, Lorenzo-Morales J, Rezaie S, Rahimi F, Mohebbi M, Maghsood AH, et al. Genotyping of Acanthamoeba isolates from clinical and environmental specimens in Iran. *Exp Parasitol.* 2009;**121**(3):242-5. doi: [10.1016/j.exppara.2008.11.003](https://doi.org/10.1016/j.exppara.2008.11.003). [PubMed: [19059239](https://pubmed.ncbi.nlm.nih.gov/19059239/)].
14. Pussard M, Pons R. Morphology of cystic wall and taxonomy of genus Acanthamoeba (Protozoa, Amoebozoa). *Protistologica.* 1977;**13**(4):557-98.
15. Visvesvara GS, Moura H, Schuster FL. Pathogenic and opportunistic free-living amoebae: Acanthamoeba spp., Balamuthia mandrillaris, Naegleria fowleri, and Sappinia diploidea. *FEMS Immunol Med Microbiol.* 2007;**50**(1):1-26. doi: [10.1111/j.1574-695X.2007.00232.x](https://doi.org/10.1111/j.1574-695X.2007.00232.x). [PubMed: [17428307](https://pubmed.ncbi.nlm.nih.gov/17428307/)].
16. Hooshyar H, Hosseinbigi B, Saraei M, Alizadeh S, Eftakhar M, Rasti S, et al. Genotyping of acanthamoeba isolated from surface and stagnant waters of qazvin, central iran. *Iran Red Crescent Med J.* 2013;**15**(6):536-8. doi: [10.5812/ircmj.4121](https://doi.org/10.5812/ircmj.4121). [PubMed: [24349758](https://pubmed.ncbi.nlm.nih.gov/24349758/)].
17. Nazar M, Haghghi A, Niyiyati M, Eftekhari M, Tahvildar-Biderouni F, Taghipour N, et al. Genotyping of Acanthamoeba isolated from water in recreational areas of Tehran, Iran. *J Water Health.* 2011;**9**(3):603-8. doi: [10.2166/wh.2011.152](https://doi.org/10.2166/wh.2011.152). [PubMed: [21976207](https://pubmed.ncbi.nlm.nih.gov/21976207/)].
18. Page FC. A revised classification of the Gymnamoebia (Protozoa: Sarcodina). *Zool J Linn Soc.* 1976;**58**(1):61-77.
19. Booton GC, Visvesvara GS, Byers T, Kelly DJ, Fuerst PA. Identification and distribution of Acanthamoeba species genotypes associated with nonkeratitis infections. *J Clin Microbiol.* 2005;**43**(4):1689-93. doi: [10.1128/JCM.43.4.1689-1693.2005](https://doi.org/10.1128/JCM.43.4.1689-1693.2005). [PubMed: [15814986](https://pubmed.ncbi.nlm.nih.gov/15814986/)].
20. Boggild AK, Martin DS, Lee T, Yu B, Low DE. Laboratory diagnosis of amoebic keratitis: comparison of four diagnostic methods for different types of clinical specimens. *J Clin Microbiol.* 2009;**47**(5):1314-8. doi: [10.1128/JCM.00173-09](https://doi.org/10.1128/JCM.00173-09).
21. Rezaeian M, Niyiyati M, Farnia SH, Haghi AM. Isolation of Acanthamoeba spp. from different environmental sources. *Iranian J Parasitol.* 2008;**3**(1):44-7.
22. Jeong HJ, Yu HS. The role of domestic tap water in Acanthamoeba contamination in contact lens storage cases in Korea. *Korean J Parasitol.* 2005;**43**(2):47-50. [PubMed: [15951638](https://pubmed.ncbi.nlm.nih.gov/15951638/)].
23. Kilvington S, Gray T, Dart J, Morlet N, Beeching JR, Frazer DG, et al. Acanthamoeba keratitis: the role of domestic tap water contamination in the United Kingdom. *Invest Ophthalmol Vis Sci.* 2004;**45**(1):165-9. [PubMed: [14691169](https://pubmed.ncbi.nlm.nih.gov/14691169/)].
24. Joslin CE, Tu EY, McMahon TT, Passaro DJ, Stayner LT, Sugar J. Epidemiological characteristics of a Chicago-area Acanthamoeba keratitis outbreak. *Am J Ophthalmol.* 2006;**142**(2):212-7. doi: [10.1016/j.ajo.2006.04.034](https://doi.org/10.1016/j.ajo.2006.04.034). [PubMed: [16876498](https://pubmed.ncbi.nlm.nih.gov/16876498/)].
25. Behniafar H, Niyiyati M, Lasjerdi Z. Molecular Characterization of Pathogenic Acanthamoeba Isolated from Drinking and Recreational water in East Azerbaijan, Northwest Iran. *Environ Health Insights.* 2015;**9**:7-12. doi: [10.4137/EHI.S27811](https://doi.org/10.4137/EHI.S27811). [PubMed: [26157334](https://pubmed.ncbi.nlm.nih.gov/26157334/)].
26. Ettinger MR, Webb SR, Harris SA, McIninch SP, C. Garman G, Brown BL. Distribution of free-living amoebae in James River, Virginia, USA. *Parasitol Res.* 2003;**89**(1):6-15. doi: [10.1007/s00436-002-0707-3](https://doi.org/10.1007/s00436-002-0707-3). [PubMed: [12474037](https://pubmed.ncbi.nlm.nih.gov/12474037/)].