

# Clinical and Epidemiological Profile of Pandemic Influenza A H1N1, H3N2, and Type B in the Southeast of Caspian Sea, Iran

Naeme Javid,<sup>1</sup> Abdolvahab Moradi,<sup>1,\*</sup> Alijan Tabarraei,<sup>1,2</sup> and Masoud Bazouri<sup>2</sup>

<sup>1</sup>Department of Microbiology, Golestan University of Medical Sciences, Gorgan, IR Iran

<sup>2</sup>Infectious Diseases Research Center, Gorgan, IR Iran

\*Corresponding author: Abdolvahab Moradi, Department of Microbiology, School of Medicine, Golestan University of Medical Sciences, Gorgan, Iran. Tel: +98-9112733321; +98-1732421289, Fax: +98-1732440225, E-mail: Abmoradi@yahoo.com

Received 2015 December 19; Revised 2017 January 28; Accepted 2017 January 31.

## Abstract

**Background:** Timely diagnosis of influenza virus is important because this virus can cause severe illness. The 2009 pdmH1N1 influenza virus spread rapidly throughout the world as the first infectious pandemic of the 21st century.

**Objectives:** The aim of this study was the investigation of clinical and epidemiological figure of influenza virus A/H1N1, A/H3N2, and Influenza B infection among patients with respiratory syndrome in Golestan province, Southeast of Caspian sea, Iran.

**Methods:** This prospective, cross sectional study took place since November 2010 through March 2014. Demographic and clinical data were collected. Nasopharyngeal (NP) swabs were taken from patients with respiratory syndrome in virus transport medium and were extracted with High Pure Viral RNA Extraction Kit. Real time PCR were performed according to the CDC recommended protocol.

**Results:** A total of 790 suspected cases were assessed; pandemic A H1N1, A/H3N2, and influenza B virus were confirmed in 25 cases (3.2%), 21 cases (2.7%), and 22 (2.8%), respectively. The greatest number of confirmed cases occurred in the age group of 25 to 34 years. There was no significant association between positive cases and age, sex, residency, and clinical symptoms.

**Conclusions:** The prevalence of pandemic Influenza viruses in recent years has caused financial losses as well as mortalities around the world. This shows the importance of the rapid diagnosis of common serotypes in our society. Using real-time RT-PCR is recommended for the early diagnosis and the rapid identification of the individuals infected with pandemic influenza virus.

**Keywords:** Pandemic H1N1, Influenza A, H3N2, Influenza B, Real-Time RT-PCR, Iran

## 1. Background

Influenza pandemics occur when new influenza A subtypes get widespread among the population. Over the past few years, the global spread of highly pathogenic avian influenza A (H5N1) and the beginning of the pandemic influenza A (pH1N1) have raised concerns about the global ranking of influenza. The use of antiviral drugs (mainly oseltamivir) combined with vaccines is the response to influenza pandemics (1). Influenza A (H1N1) virus infects young and healthy adults suggesting that this group are more susceptible to the disease. It is possible that some level of cross protection antibodies exist in older people. High risk groups for this disease include pregnant women, old people, and people with serious medical illness (2).

Symptoms of pandemic influenza consist of fever, cough, body pain, headache, sore throat, and gastrointestinal complications. This illness is mostly self-limited, thus, a large proportion of infected patients do not get registered in health centers and the Statistics estimating the disease are not correct (3). The definition of influenza A (H1N1) was having high grade fever (upper than 38°C) or at least having two symptoms of respiratory disease. A con-

firmed case of influenza A (H1N1) was defined as a patient with high grade fever (> 38°C) or at least two of respiratory symptoms. The patient have H1N1 viral infection should be also confirmed by reverse transcriptase PCR (RT-PCR) (4-6).

Some cases have a high incidence of local transmission (such as in New York), but many other cases have led to low transmission or no secondary transmission. The mortality of influenza virus is low (2 % of hospitalized patients), but its morbidity is high (7). To control the influenza A H1N1 infection, early diagnosis with a cost-effective, rapid, and effectual assay is necessary, exclusively in developing countries. Using real time RT-PCR assay compared with other molecular techniques is recommended for the early diagnosis and the rapid identification of individuals infected with pandemic influenza virus. Fast and accurate detection of influenza improves medical handling by proper conditions of prophylaxis, rapid treatment, and appropriate management plans for public health responses to the outbreaks and the avoidance of unnecessary treatment (8).

Influenza virus infection is not detectable only with symptoms. The clinical picture of the disease is similar to respiratory syncytial virus, parainfluenza viruses, ade-

noviruses, coronaviruses, and metapneumovirus (9). Correct diagnosis is urgent for the identification of pandemic influenza, surveillance, and public health interventions. Without practical laboratory tests, it is not possible to provide appropriate response efforts (10).

## 2. Objectives

The aim of the present study was the investigation of clinical and epidemiological figure of influenza virus A/H1N1, A/H3N2, and influenza B infection among patients with respiratory syndrome in Golestan province, the Southeast of Caspian sea, Iran, 2011 to 2014.

## 3. Methods

### 3.1. Specimens

This prospective, cross sectional study took place since November 2010 through March 2014. The study population was inclusive of all suspected specimens of pandemic influenza A (H1N1) virus infection who had attended hospitals and healthcare centers in Golestan province over the study period. All patients with symptoms like influenza were included in this study and often had high grade fever ( $> 38^{\circ}\text{C}$ ) or any of respiratory symptoms.

Suspected cases of influenza were confirmed using a Real-time reverse-transcriptase-polymerase-chain-reaction (rRT-PCR) assay at the laboratory of influenza research center in Golestan University of Medical Sciences according to the recommended protocol of the U.S. center for disease control and prevention (CDC) (11).

In this study, the center of influenza received 790 specimens. Acceptable specimen types were given nasopharyngeal (NP) swab. Following the specimen collection in the health care center, Specimens were placed in viral transport medium and were transported to the Influenza center on cold packs.

### 3.2. Sample Processing and RNA Extraction

Specimens were processed for RNA extraction. The extractions were performed following the protocols supplied with high pure viral RNA extraction kit (Roche Diagnostics, Germany).

### 3.3. One Step Real-Time PCR Testing

After RNA extraction with High pure viral RNA extraction kit (Roche, Germany), specimens were tested for influenza A and B viruses using TaqMan probes real-time RT-PCR (rRT-PCR) according to the manual of Centers for Disease Control and Prevention (CDC) and by Invitrogen

SuperScript III Platinum one step qRT-PCR kit. The subtype of influenza virus was determined for positive influenza A specimen. The subtyping consisted of A/H3 influenza (hemagglutinin gene), A pandemic (H1N1) 2009 virus (hemagglutinin gene), and Influenza B virus (nucleoprotein gene).

PCR was performed in 25  $\mu\text{L}$  volumes containing 12.5  $\mu\text{L}$  2rxn Buffer, 10  $\mu\text{M}$  of each primer (40  $\mu\text{M}$  of Influenza A primers), 5 pmol of probe, 0.5  $\mu\text{L}$  of Rox, 2.5 unit of Enzyme mix and 5  $\mu\text{l}$  of RNA template. The reaction was carried out in a 7300 ABI Real time PCR and with the following settings:  $50^{\circ}\text{C}$  for 30 minutes,  $95^{\circ}\text{C}$  for 5 minutes, followed by 40 cycles of denaturation at  $95^{\circ}\text{C}$  for 15 seconds and annealing and extension  $60^{\circ}\text{C}$  for 30 seconds (Table 1).

## 4. Results

Among a total of 790 suspected cases, the following results were obtained: the number of influenza cases was 68 (8.6%) which among the proportion of influenza A/H3N2 was 21 (30.8%), influenza A/pdmH1N1 25 (36.7%), and influenza B 22(32.3%) (Table 2). Among the 790 suspected cases, male cases accounted for 72 and female cases for 718 patients. In confirmed cases 7 (10.3%) were male and 61 (89.7%) were female. The mean age was 29.79 years (SD = 16.0, ranging from  $< 1$  days old to 92 years old). The greatest number of confirmed cases occurred in the age group of 25 to 34 years, accounting for 354 (44.8%) of all cases, followed by 227 (28.7%) in 15 to 24 years and the lowest number was in  $\geq 65$  years, with 22 cases (2.8%) of all cases.

Among the total cases, 244 (30.9%) were hospitalized of which 16 (23.5%) were confirmed cases of influenza. From a total of 790 suspected cases, 283 (35.8%) were pregnant women including 28 (41.2%) confirmed cases (Tables 3 and 4). The most signs were cough (88.7%) and fever (84.3%). As shown in Figures 1 and 2, the highest incidence of confirmed cases occurred in December.

## 5. Discussion

Influenza virus infection can spread rapidly and that is responsible for morbidity and mortality each year in the world. Mutations in the genes of the influenza virus lead to diseases with different symptoms. Therefore, it is important to identify the symptoms in different areas. Checking the symptoms can help in the diagnosis of the disease and its treatment. Chronic respiratory diseases as well as hypertension, diabetes mellitus, and pregnancy were the most common risk factors in the sever disease. Moreover, acute respiratory distress syndrome and viral pneumonia were the most common clinical causes of mortality (7, 12).

**Table 1.** The List of Used Primer and Probe Set

Name		Sequence (5' > 3')	Working Concentration, μM
Influenza A	Primer F	GACCRATCCTGTCACCTCTGAC	40
	Primer R	AGGGCATTYGGACAAAKCGTCTA	40
	Probe <sup>a</sup>	TGCAGTCCTCGCTCACTGGGCACG	10
SW InfA	Primer F	GCACGGTCAGCACITTATYCTRAG	10
	Primer R	GTGRGCTGGGTTTTCATTGGTC	10
	Probe <sup>b</sup>	CYACTGCAAGCCCA <sup>T</sup> ACACACAAGCAGGCA	10
H3N2	Primer F	AAGCATTCCYAATGACAAACC	10
	Primer R	ATTGCRCCRAATATGCCTCTAGT	10
	Probe	CAGGATCACATATGGSCCTGTCCAG	10
Influenza B	Primer F	GAGACACAATTGCCTACTGCTT	10
	Primer R	TTCTTTCCACCGAACCAAC	10
	Probe	AGAAGATGGAGAAGGCAAAGCAGAACTAGC	10
RNAase P	Primer F	AGATTGGACCTGCGAGCG	10
	Primer R	GAGCGGCTGTCTCCACAAGT	10
	Probe	TTCTGACCTGAAGGCTCTGCGCG	10

<sup>a</sup>TaqMan probes consist of FAM reporter (5'-end with the reporter molecule 6-carboxyfluorescein) and BHQ1 quencher (Blackhole Quencher 1) at the 3'-end.

<sup>b</sup>Quenched internally at a modified<sup>T</sup> residue with BHQ1, to prevent probe extension by Taq polymerase (11).

**Table 2.** The Frequency of Influenza Viruses and the Proportion for Influenza Type /Subtype by Year and Demographic Characteristics<sup>a</sup>

Characteristics	No. Samples Tested	No. of Influenza Cases	No. of Influenza Cases			
			A (H3N2)	A (H1N1) pdm09	Type B	Type A and B
<b>Year</b>						
2010 - 2014	790	68 (8.6)	21 (30.8)	25 (36.7)	22 (32.3)	0
<b>Gender</b>						
Male	72	7 (10.3)	4 (57.1)	1 (14.3)	2 (28.6)	0
Female	718	61 (89.7)	17 (27.9)	24 (39.3)	20 (32.8)	0
<b>Age group, y</b>						
1 to 14	32	2 (2.9)	1 (50)	1 (50)	0	0
15 to 24	227	20 (29.4)	5 (25)	8 (40)	7 (35)	0
25 - 34	354	28 (41.2)	8 (28.6)	12 (42.8)	8 (28.6)	0
35 to 44	83	9 (13.2)	5 (55.6)	3 (33.3)	1 (11.1)	0
45 to 54	42	4 (5.9)	0	0	4 (100)	0
55 to 64	30	3 (4.4)	2 (66.7)	1 (33.3)	0	0
Over 65	22	2 (2.9)	0	0	2 (100)	0
Total	790	68 (8.6)	21 (30.8)	25 (36.7)	22 (32.3)	0

<sup>a</sup>Values are expressed as No. (%).

In this study, out of 790, the suspected number of influenza cases was 68 (8.6%) among which the proportion of influenza A/ pdmH1N1 was 25 (36.7%). Although in Monsef's

study (2013) in Hamadan out of 180 patients, 63.8% were H1N1 positive. In Dashti-Khavidaki's study (2009, Tehran), 34.5%, which was consistent with our results. In Afrasi-

**Table 3.** The Comparison of Variable Features in Confirmed and Unconfirmed Cases of Pandemic Influenza A and B Virus<sup>a</sup>


Variable		Confirmed Cases (68 Cases)	Unconfirmed Cases (722 Cases)	P Value
Sex	Male	7 (10.3)	65 (9)	0.07
	Female	61 (89.7)	657 (91)	
Residency	Urban	21 (30.9)	216 (29.9)	0.5
	Rural	47 (69.1)	506 (70.1)	
Hospitalization	Yes	16 (23.5)	228 (31.6)	0.2
	No	52 (76.5)	494 (48.4)	
Pregnancy	Yes	28 (41.2)	255 (35.3)	0.01
	No	40 (58.8)	467 (64.7)	

<sup>a</sup>Values are expressed as No. (%).

**Table 4.** Clinical Symptoms and the Tests of 68 Cases of Influenza A and B<sup>a</sup>

Positive Sample	Clinical Symptoms and Tests				
	Fever	Cough	Myalgia	Sore throat	Rhinorrhea
pH1N1	24 (96)	24 (96)	22 (88)	10 (40)	15 (60)
H3N2	20 (95.2)	19 (90.4)	16 (76.1)	11 (52.3)	12 (57.1)
B	20 (90.9)	21 (95.4)	16 (72.7)	16 (72.7)	11 (50)


<sup>a</sup>Values are expressed as No. (%).



**Figure 1.** Frequency of Unconfirmed and Confirmed Cases of Pandemic Influenza A (pH1N1), H3N2, and Influenza B Based on the Month of Occurrence

abian’s study (2009, Kurdistan), 14.8%; Jedary Seifi’s study (2014 Tabriz), 20.3%; and Gao et al.’s (2009, USA), 19.5% were H1N1 positive (3, 7, 12-14). After several years of the 2009 pan-

dem ic outbreaks, H1N1pdm09 was, still, the dominant circulating strain. This was supported by world health organization in 2010 which says that the pandemic strains may



**Figure 2.** Age Distribution of Influenza Virus Types and Subtypes

circulate as seasonal viruses.

Influenza A/H3N2, in our result, was 21 (30.8%), also, in the study by Haghshenas (2011-2013) in which a total number of 201 patients (35.2%) were diagnosed with influenza A1 H3N2 infection. Contrary to Hajikhezri’s study which was 6.8%. Burden of influenza B in our study was 32.3%, also, in Gaini et al.’s in global influenza B study (2000 to 2013) was 22.6%, and in Qi et al.’s (2011-2015), influenza B was 34.1% (15-18). In the current research, fever (96%), cough (96%), and myalgia (88%) were the most common symptoms among the infected patients with pH1N1, but Rhino rhea and sore throat were present only in 60% and 40% of the patients, respectively (Table 5). These observations were in accordance with the results obtained in other studies, finally, it was found that fever, cough, and myalgia were the best diagnostic model for H1N1infection (19).

In this study, among the confirmed cases, 7 (10.3%) were male and 61 (89.7%) were female. Also, Li et al. in 2011 in a their study reported that female subjects with influenza A are higher than males (20). Also, this point was reported by other researchers (21-23). And in Cao et al.’s study, male subjects were higher, however, Caini et al. presented that sexes were equally distributed (15). And in Cao et al.’s study, there was no sex difference in the incidence of confirmed pandemic influenza A (H1N1) (24). The mean age of the confirmed cases in the present study was 29.7 years and this was similar to other studies. In this study, there were 22 people aged > 65 years. The lower level of infection and susceptibility to influenza virus in this age group compared with younger age groups can be explained by pre-existing immunity, lower chances of being in crowded places, and fewer contacts with other people (2).

In this study, the peak incidence of the disease occurred in December and, considering the age group of samples, these peaks can be attributed to the transmission

of the disease in universities and schools (2). In the last of the pandemic influenza, fast and accurate patient identification remains crucial in preventing the extensive transmission (2).

In conclusion, the outbreak of pandemic Influenza viruses in recent years has created a lot of deaths and financial losses around the world. This result shows the importance of the rapid identification of common serotypes in our society. The detection of antigenic types of circulating viruses has a great importance in vaccine preparation for high-risk individuals. Therefore, using RT-PCR is recommended for early diagnosis and rapid identification of individuals infected with pandemic influenza virus. In addition, it can accelerate vaccine manufacture. Molecular methods should be used in vaccine manufacturing in cases of acute infections. Moreover, Scheduling the urgent influenza A vaccination program is suggested, which can definitely prevent the spread of the virus.

### Acknowledgments

The authors thank all staff of the influenza reference laboratory in Golestan University of Medical Sciences.

**Table 5.** The Comparison of the Frequency of Influenza Viruses and the Proportion of Influenza Type

Study	Year		Influenza Cases, %		
			A (H3N2)	A (H1N1) pdm09	Type B
Dashti-Khavidaki et al.	2009	Iran-Tehran	-	34.5	-
Afrasiabian et al.	2009	Iran-Kurdestan	-	14.8	-
Gao et al.	2009	USA	-	19.5	-
Monsef et al.	2013	Iran-Hamedan	-	63.8	-
Jedary Seifi et al.	2014	Iran-Tabriz	-	20.3	-
Haghshenas et al.	2011 - 2013	Iran-Mazandaran	35.2	-	-
Hajikhezri et al.	2009 - 2010	Iran-Ahvaz	6.8	-	0.7
Gaini et al.	200 - 2013	Global study	-	-	22.6
Qi et al.	2011 - 2015	china	44.7	21	34.1

**References**

- Steain MC, Dwyer DE, Hurt AC, Kol C, Saksena NK, Cunningham AL, et al. Detection of influenza A H1N1 and H3N2 mutations conferring resistance to oseltamivir using rolling circle amplification. *Antiviral Res.* 2009;**84**(3):242-8. doi: [10.1016/j.antiviral.2009.09.010](https://doi.org/10.1016/j.antiviral.2009.09.010). [PubMed: 19800370].
- Panning M, Eickmann M, Landt O, Monazahian M, Olschlager S, Baumgarde S, et al. Detection of influenza A(H1N1)v virus by real-time RT-PCR. *Euro Surveill.* 2009;**14**(36) [PubMed: 19758541].
- Afrasiabian S, Mohsenpour B, Bagheri KH, Barari M, Ghaderi E, Hashemi R, et al. Epidemiological survey on pandemic influenza A (H1N1) virus infection in Kurdistan province, Islamic Republic of Iran, 2009. *East Mediterr Health J.* 2014;**20**(3):169-74. [PubMed: 24950074].
- Bakhshayeshkaram M, Saidi B, Tabarsi P, Zahirifard S, Ghofrani M. Imaging Findings in Patients With H1N1 Influenza A Infection. *Iran J Radiol.* 2011;**8**(4):230-4. doi: [10.5812/iranjradiol.4554](https://doi.org/10.5812/iranjradiol.4554). [PubMed: 23329946].
- Natoghian A, Dadras M, Gouya MM, Nabavi M, Soroush M, Hooman N, et al. Pandemic Flu in Islamic Republic of Iran; A Review of Health System Response From July to November. *Arch Pediatr Infect Dis.* 2013;**1**(2):80-6. doi: [10.5812/pedinfect.9074](https://doi.org/10.5812/pedinfect.9074).
- Yavarian J, Naseri M, Shadab A, Shafiei Jandaghi NZ, Mokhtari Azad T. Epidemiological aspects of pandemic influenza A(H1N1) virus from 2009 to 2011 in Iran. *Influenza Other Respir Viruses.* 2012;**6**(6):e74-6. doi: [10.1111/j.1750-2659.2012.00357.x](https://doi.org/10.1111/j.1750-2659.2012.00357.x). [PubMed: 22487173].
- Monsef F, Esmaeili R, Eghbalian F, Pourhossein A, Khiabanchian O, Monsef A. Is there yet influenza virus A/H1N1 infection in Hamadan, Iran? : A cross-sectional serological study in lower the age of 16. *Life Sci J.* 2013;**10**:123-6.
- Ma XJ, Shu YL, Nie K, Qin M, Wang DY, Gao RB, et al. Visual detection of pandemic influenza A H1N1 Virus 2009 by reverse-transcription loop-mediated isothermal amplification with hydroxynaphthol blue dye. *J Virol Methods.* 2010;**167**(2):214-7. doi: [10.1016/j.jviromet.2010.03.027](https://doi.org/10.1016/j.jviromet.2010.03.027). [PubMed: 20381535].
- Dwyer DE, Smith DW, Catton MG, Barr IG. Laboratory diagnosis of human seasonal and pandemic influenza virus infection. *Med J Aust.* 2006;**185**(10 Suppl):S48-53. [PubMed: 1715952].
- Jernigan DB, Lindstrom SL, Johnson JR, Miller JD, Hoelscher M, Humes R, et al. Detecting 2009 pandemic influenza A (H1N1) virus infection: availability of diagnostic testing led to rapid pandemic response. *Clin Infect Dis.* 2011;**52** Suppl 1:S36-43. doi: [10.1093/cid/ciq020](https://doi.org/10.1093/cid/ciq020). [PubMed: 21342897].
- Geneva F. CDC protocol of real time RT-PCR for swine influenza A (H1N1); 2009.
- Jedary Seifi S, Sabouri Ghannad M. Comparison of Real-Time Polymerase Chain Reaction and Conventional Cell Culture for Detection of Influenza A in Tabriz, Iran. *Avicenna J Clin Microbiol Infect.* 2014;**1**(2) doi: [10.17795/ajcmi-21034](https://doi.org/10.17795/ajcmi-21034).
- Dashti-Khavidaki S, Khalili H, Gholamalipour F, Soudbakhsh A, Talasaz AH, Hajabdolbaghi M, et al. Approach to Pandemic 2009 influenza: first report from a main referral hospital for Pandemic H1N1 influenza care in Iran. *J Infect Dev Ctries.* 2010;**4**(10):629-35. [PubMed: 21045355].
- Gao F, Loring C, Laviolette M, Bolton D, Daly ER, Bean C. Detection of 2009 pandemic influenza A(H1N1) virus Infection in different age groups by using rapid influenza diagnostic tests. *Influenza Other Respir Viruses.* 2012;**6**(3):e30-4. doi: [10.1111/j.1750-2659.2011.00313.x](https://doi.org/10.1111/j.1750-2659.2011.00313.x). [PubMed: 22114876].
- Caini S, Huang QS, Ciblak MA, Kuszniarz G, Owen R, Wangchuk S, et al. Epidemiological and virological characteristics of influenza B: results of the Global Influenza B Study. *Influenza Other Respir Viruses.* 2015;**9** Suppl 1:3-12. doi: [10.1111/irv.12319](https://doi.org/10.1111/irv.12319). [PubMed: 26256290].
- Haghshenas M, Jafarian E, Babamahmoodi F, Tabrizi A, Nandoost S, Alizadeh-Navaei R. Prevalence of influenza A/H3N2 virus in northern Iran from 2011 to 2013. *Caspian J Intern Med.* 2015;**6**(2):116-9. [PubMed: 26221512].
- Hajikhezri Z, Makvandi M, Samarraf-zadeh AR, Neisi N, Ahmadi K. Relative frequency of seasonal influenza A and B in Khuzestan province by reverse transcription-polymerase chain reaction (RT-PCR) during 2009-2010. *Jundishapur J Microbiol.* 2013;**6**(4):e5312. doi: [10.5812/jjm.5312](https://doi.org/10.5812/jjm.5312).
- Qi L, Xiong Y, Xiao B, Tang W, Ling H, Long J, et al. Epidemiological and Virological Characteristics of Influenza in Chongqing, China, 2011-2015. *PLoS One.* 2016;**11**(12):e0167866. doi: [10.1371/journal.pone.0167866](https://doi.org/10.1371/journal.pone.0167866). [PubMed: 27936139].
- Kim CO, Nam CM, Lee DC, Han SH, Lee JW. Clinical predictors of novel influenza A (H1N1) infection in Korea. *Yonsei Med J.* 2010;**51**(6):895-900. doi: [10.3349/ymj.2010.51.6.895](https://doi.org/10.3349/ymj.2010.51.6.895). [PubMed: 20879057].
- Li T, Liu Y, Di B, Wang M, Shen J, Zhang Y, et al. Epidemiological investigation of an outbreak of pandemic influenza A (H1N1) 2009 in a boarding school: serological analysis of 1570 cases. *J Clin Virol.* 2011;**50**(3):235-9. doi: [10.1016/j.jcv.2010.11.012](https://doi.org/10.1016/j.jcv.2010.11.012). [PubMed: 21195022].
- Afzali H, Nematian M, Rajabi J, Soleimani Z, Momen-Heravi M, Salehi A, et al. Epidemiological survey of confirmed influenza A (H1N1) in Kashan, Aran and Bidgol cities during 2009-10. *Fez J Kashan Univ Med Sci.* 2011;**15**(3).
- Saleh P, Noshad H, Naghili B. Demographic and Paraclinical Findings of Patients with Novel H1N1 Infection Hospitalized in Infectious Dis-

- ease Ward, Sina Hospital, Tabriz, Iran. *ZUMS J.* 2011;**19**(75):84-93.
23. Shiley KT, Nadolski G, Mickus T, Fishman NO, Lautenbach E. Differences in the epidemiological characteristics and clinical outcomes of pandemic (H1N1) 2009 influenza, compared with seasonal influenza. *Infect Control Hosp Epidemiol.* 2010;**31**(7):676-82. doi: [10.1086/653204](https://doi.org/10.1086/653204). [PubMed: 20500086].
24. Cao B, Li XW, Mao Y, Wang J, Lu HZ, Chen YS, et al. Clinical features of the initial cases of 2009 pandemic influenza A (H1N1) virus infection in China. *N Engl J Med.* 2009;**361**(26):2507-17. doi: [10.1056/NEJMoa0906612](https://doi.org/10.1056/NEJMoa0906612). [PubMed: 20007555].