



# Multicenter Identification and Antifungal Susceptibility Patterns of *Candida* Species Isolated from Clinical Samples

Parisa Badiie,<sup>1</sup> Hamid Badali,<sup>2</sup> Kambiz Diba,<sup>3</sup> Hadis Jafarian,<sup>1\*</sup> Rasoul Mohammadi,<sup>4</sup> Hossein Mirhendi,<sup>5</sup> and Mohammad Javad Najafzadeh<sup>6</sup>

<sup>1</sup>Prof. Alborzi Clinical Microbiology Research Center, Shiraz University of Medical Sciences, Shiraz, IR Iran

<sup>2</sup>Department of Medical Mycology and Parasitology/Invasive Fungi Research Centre (IFRC), School of Medicine, Mazandaran University of Medical Sciences, Sari, IR Iran

<sup>3</sup>Cellular and Molecular Research Center, School of Medicine, Urmia University of Medical Sciences, Urmia, IR Iran

<sup>4</sup>Department of Medical Parasitology and Mycology, Faculty of Medicine, Isfahan University of Medical Sciences, Isfahan, IR Iran

<sup>5</sup>Department of Medical Mycology and Parasitology, School of Public Health and Institute of Health Research, Tehran University of Medical Sciences, Tehran, IR Iran

<sup>6</sup>Department of Parasitology and Mycology, School of Medicine, Mashhad University of Medical Sciences, Mashhad, IR Iran

\*Corresponding author: Hadis Jafarian, Prof. Alborzi Clinical Microbiology Research Center, Nemazi Hospital, Zand Ave, Shiraz, IR Iran. Tel: +98-71-36474304, Fax: +98-71-36474303, E-mail: hdsjafarian@yahoo.com

Received 2017 June 25; Revised 2017 October 08; Accepted 2017 October 18.

## Abstract

**Background:** Invasive fungal infections without proper treatment could lead to high mortality rate, especially in immunocompromised patients. *Candida* species distribution and drug susceptibility patterns vary in different areas. Understanding the etiologic agents and drug susceptibility patterns in each region are required for the best management of patients with *Candida* infections.

**Objectives:** The aim of this study was to identify *Candida* species isolated from clinical samples of six university hospitals in Iran and detect their susceptibility patterns to seven antifungal agents.

**Methods:** Clinical samples from patients with fungal infections were cultured on Sabouraud dextrose agar. Isolated yeasts were identified by API 20C AUX kit, according to the manufacturer's instructions. Drug susceptibility patterns to amphotericin B, caspofungin, voriconazole, fluconazole, posaconazole, itraconazole and ketoconazole were determined, according to CLSI M27-A3 and S4.

**Results:** In total, 428 species of *Candida* were isolated from clinical samples (1950 samples). Most isolated species were *Candida albicans*, followed by *C. tropicalis*, *C. parapsilosis*, *C. kefyr*, *C. famata*, *C. glabrata*, *C. krusei*, *C. dubliniensis*, *C. guilliermondii* and *C. lusitaniae*. Sensitivity rate of *C. albicans* to amphotericin B, caspofungin, voriconazole, fluconazole, and itraconazole was 96.6%, 99.5%, 88.6%, 90.6%, and 52% with MIC<sub>90</sub> values equal to 0.25 µg/mL, 0.125 µg/mL, 0.125 µg/mL, 2 µg/mL, and 1 µg/mL, respectively. The MIC<sub>90</sub> values for ketoconazole and posaconazole were 0.125 µg/mL and 0.064 µg/mL, respectively. Different sensitivity to antifungal agents was present in non-*albicans Candida* species especially in *C. krusei*, *C. glabrata*, and *C. tropicalis*.

**Conclusions:** According to this study, *C. albicans* is the most prevalent etiologic agent in infected patients and caspofungin is the most effective antifungal agent. Knowledge about etiologic agents and their susceptibility patterns in each region is helpful for successful treatment of the patients.

**Keywords:** *Candida albicans*, Fluconazole, Amphotericin B, Itraconazole, *Candida tropicalis*

## 1. Background

Systemic candidiasis in immunocompromised patients is associated with high morbidity and mortality rates, especially in those not responsive to antifungals (1, 2). *Candida albicans* is the most prevalent isolate from human infections, however other *Candida* species have been reported as well (3, 4). The relative frequency of non-*albicans Candida* species varies in different areas, for example, in North America *C. glabrata* is the second and in Iran is the third species isolated after *C. albicans* (5,

6). The epidemiology of candidemia varies, according to geographical region. Also, a variety in the distribution of *Candida* species in different areas was reported (2-6). The reason of emerging non-*albicans* species is not clear yet and can be associated with improvements in more sensitive methods for the identification of *Candida* species and indiscriminate use of antifungal drugs for prophylaxis and treatment (7, 8).

In the recent years, treatments of systemic candidiasis are a challenge due to resistant etiologic agents and the emergence of infections caused by species other than *C. al-*

*bicans* (3, 5, 6). Adequate knowledge about the etiologic agents and drug susceptibility patterns are required for treating the hospitalized patients with risk factors of systemic candidiasis.

## 2. Objectives

This study aimed at identifying *Candida* species isolated from clinical samples in six university hospitals (Shiraz, Isfahan, Tehran, Urmia, Sari, and Mashhad) and determined their susceptibility patterns to seven antifungal agents.

## 3. Methods

### 3.1. Isolates Sources

Clinical samples (1950 samples) including abscess, wound, blood, cerebrospinal fluid, bronchoalveolar lavage, vaginal discharge, nails, and sputum were sent to mycology laboratories of tertiary hospitals in Shiraz, Isfahan, Tehran, Urmia, Sari, and Mashhad. They were cultured on Sabouraud dextrose agar (Merck, Germany), containing chloramphenicol (Merck, Germany), according to standard protocols related to each sample and incubated for 48 hours at room temperature for rapidly growing species and up to seven days for slow growing. All *Candida* species isolated from hospitalized patients with signs and symptoms of fungal infections were entered in this study. Isolated *Candida* species were transferred to Prof. Alborzi Clinical Microbiology Research Center, Shiraz University of Medical Sciences, and identified by API 20C AUX kit (BioMerieux, France), according to the manufacturer's protocol.

### 3.2. Antifungal

Antifungal susceptibility test to amphotericin B (AMB, Sigma-Aldrich, Germany), caspofungin (CAS, Sigma-Aldrich, USA), voriconazole (VRC, Sigma-Aldrich, USA), fluconazole (FLU, Sigma-Aldrich, USA), posaconazole (POS, Sigma-Aldrich, Germany), itraconazole (ITR, Sigma-Aldrich, India), and ketoconazole (KET, Sigma-Aldrich, China) was performed according to Clinical and Laboratory Standards Institute (CLSI) M27-A3 and CLSI M27-S4 (9, 10). Briefly, RPMI medium with L-Glutamine and without sodium bicarbonate (Sigma, Germany) was buffered with 0.165 mole per liter of 3-(N-Morpholino) propanesulfonic acid, 4 morpholinepropanesulfonic acid (Sigma, Germany). pH was adjusted to 7 and filtered in sterile conditions.

### 3.3. Susceptibility Tests

Stock concentration of each drug was prepared in a suitable solvent (water for CAS and DMSO (Merck, Germany) for other drugs) and diluted with RPMI to obtain the working concentration solution. Serial dilutions from AMB, CAS, VOR, POS, ITR, and KET, ranging from 0.032 to 16  $\mu\text{g}/\text{mL}$  and for FLU from 0.125 to 64  $\mu\text{g}/\text{mL}$ , were prepared. For each series, positive (well without antifungal) and negative (well without yeast) controls were considered. In a 96-well plate (JETBIOFIL, China), 100  $\mu\text{L}$  of RPMI and serial dilutions were poured in each well. The suspensions with a concentration equal to 0.5 McFarland were prepared from 24 - to 48 - hour incubated colonies ( $1 \times 10^6$  to  $5 \times 10^6$  cells/mL) and diluted at 1:1000 with RPMI ( $1 \times 10^3$  to  $5 \times 10^3$  cells/mL). Yeast suspensions (100  $\mu\text{L}$ ) were added to each well, except the control negative well. The final yeast concentration in each well was  $0.5 \times 10^3$  to  $2.5 \times 10^3$  cells/mL. Plates were kept at 35°C for 24 to 48 hours and were read visually after the incubation time. Minimum Inhibitory Concentrations (MIC) for CAS, VOR, FLU, POS, ITR, and KET were described as the lowest concentration of the drug that could reduce fungal growth by 50% to 80%, compared to positive controls. For AMB, MIC was described as the lowest concentration of the drug that could stop any visible yeast growth.

### 3.4. Statistical Analysis

Data were analyzed using the WHONET 5.6 software, and MIC50, MIC90 (MIC values which inhibit 50% and 90% of the isolates) and geometric mean was calculated for each drug and each isolate. New *Candida* species breakpoint provided by the CLSI was used in the present study. These new breakpoints are drug and species specific; POS and KET have no breakpoint in the new CLSI (10).

## 4. Results

Overall, 428 *Candida* were isolated from clinical samples, the prevalent species was *C. albicans* (273, 63.78%), followed by *C. tropicalis* (38, 8.87%), *C. parapsilosis* (35, 8.17%), *C. kefyr* (20, 4.67%), *C. famata* (20, 4.67%), *C. glabrata*, (18, 4.2%), *C. krusei* (10, 2.34%), and other species (14, 3.27%). The second most frequent isolate from Shiraz, Mashhad, and Sari was *C. tropicalis* while *C. parapsilosis* in Isfahan and Tehran, and *C. famata* in Urmia were the second most frequently isolated. Distributions of the isolated species are presented in Table 1.

Susceptibility patterns of the common *Candida* species isolates are shown in Table 2 and the uncommon *Candida* species isolates are shown in Table 3. The sensitivity rates of *C. albicans* species to AMB, CAS, VRC, and FLU were

**Table 1.** Distribution of *Candida* Species Isolated From Six University Hospitals in Iran

Species	Shiraz	Mashhad	Isfahan	Urmia	Tehran	Sari	Total <sup>a</sup>
<i>Candida albicans</i>	47	52	55	38	41	40	273 (63.78)
<i>Candida tropicalis</i>	7	6	9	5	4	7	38 (8.87)
<i>Candida parapsilosis</i>	3	5	10	6	7	4	35 (8.17)
<i>Candida kefyr</i>	3	2	5	4	3	3	20 (4.67)
<i>Candida famata</i>	2	2	-	9	5	2	20 (4.67)
<i>Candida glabrata</i>	3	4	6	1	2	2	18 (4.20)
<i>Candida krusei</i>	5	1	3	-	1	-	10 (2.34)
Others <sup>b</sup>	1	1	4	4	2	2	14 (3.27)
<b>total</b>	<b>71</b>	<b>73</b>	<b>92</b>	<b>67</b>	<b>65</b>	<b>60</b>	<b>428</b>

<sup>a</sup> Data are presented as No (%)

<sup>b</sup> *Candida dubliniensis*, *Candida guilliermondii* and *Candida lusitanae*

96.6% (271.9/273), 99.5% (272/273), 88.6% (242/273), and 88.3% (241/273), respectively. The resistance rate of *C. albicans* to ITR was 12.7% with susceptible dose dependence of 35.3% and sensitivity rate of 52%. The MIC90 values and geometric means for POS and KET in this species were 0.064 µg/mL and 0.027 µg/mL, and 0.125 µg/mL and 0.028 µg/mL, respectively.

*Candida tropicalis* was the second most isolated species from patients; it was more sensitive to CAS, POS and KET with MIC90 value of 0.125 µg/mL. The MIC90 value of *C. glabrata* for AMB and CAS, VRC, FLU, POS, ITR and KET was 4 µg/mL, 2 µg/mL, 2 µg/mL, 16 µg/mL, 2 µg/mL, 4 µg/mL, and 2 µg/mL, respectively. The resistance rates of *C. guilliermondii* to both AMB and ITR were 20%, with MIC90 value of 16 µg/mL for KET.

The most resistance rate among isolated species to AMB and CAS was in *C. krusei* (MIC90 8 and 2 µg/mL, respectively). *Candida krusei* sensitivity rates to AMB, VOR and ITR were 70% (7/10), 50% (5/10), and 33.3% (3/10), respectively. *Candida parapsilosis* was completely sensitive to all antifungal agents and susceptible dose dependent rate to ITR was 60%. *Candida kefyr* was sensitive to all the antifungal drugs and 10% (2/20) of *C. famata* were resistant to ITR and 5% (1/20) resistant to AMB.

## 5. Discussion

*Candida* species could cause severe infections with high morbidity and mortality in hospitalized patients (2-4). *Candida albicans* was reported as the most isolated species from infected patients in this study and other studies in Asian, European, and American countries (3, 5, 11-14). A global increase was seen in the number of infections

caused by species other than *C. albicans*, including *C. tropicalis*, *C. parapsilosis*, *C. glabrata*, and *C. krusei* (8, 15). *Candida tropicalis* was the second isolated *Candida* species in this study. The epidemiology of this species was reported as 15.3% in Brazil (16), 13.4% in Iran (17), and 8.4% in Italy (18).

Amphotericin B is a polyene antifungal drug with high activity against a wide variety of fungal pathogens. The sensitivity rates of *C. albicans* to AMB in this study was 96.6% (271.9/273), and in other studies, this was reported as 97.4% (37/38), 96.6% (113/117), 100% (93/93), and 93% (160/172) (11, 15, 19, 20). The sensitivity rate of *C. glabrata* in this study was 88.9% (16/18) and in other studies, this was reported as 85% (34/40) and 93.8% (15/16) (6, 11). This rate for *C. krusei* was 70% (7/10) in the present study and 90% (56/62) in the other studies (6). The differences in sensitivity rates could be due to patient population or type of the study. Given the limited number of isolates in non-*albicans* species, high resistant rate may not be reliable enough.

Echinocandins are a new line of antifungal drugs. According to Espinel-Ingroff et al. (2013), use of the CLSI species-specific CAS breakpoint could lead to results indicating an excessive number of sensitive isolates (such as *C. glabrata* and *C. krusei*) as resistant. Therefore, they suggested that routine testing or reporting of CLSI CAS MICs for *Candida*, according to CLSI (2012), is not suitable (21). Generally, in this study, CAS was an effective agent against *Candida* species, except some *C. glabrata* and *C. krusei* species (MIC90 = 2 µg/mL). The MIC90 values of this drug for *C. glabrata* and *C. krusei* were reported as 0.19 and 0.75 µg/mL in colonized isolates in 2014 (15); 4 and 1 µg/mL in patients hospitalized at ICUs and urology wards (22); and 0.125 and 0.25 µg/mL in patients with cancer (11). In this study, all isolates were from infected patients and this is the reason for the differentiation between the studies.

**Table 2.** Sensitivity Pattern of Common *Candida* Species Isolated From Clinical Samples in Six University Hospitals of Iran

<i>Candida</i>	Antibiotic Name	Breakpoints	%R <sup>a</sup>	%I <sup>b</sup>	%S <sup>c</sup>	MIC50, µg/mL	MIC9, µg/mL	MIC Range, µg/mL
<i>Candida albicans</i>	Amphotericin B	S ≤ 1, R ≥ 1	3.4	0	96.6	0.032	0.25	0.032 - 16
	Caspofungin	S ≤ 0.25, I = 0.5, R ≥ 1	0.5	0	99.5	0.032	0.125	0.032 - 1
	Voriconazole	S ≤ 0.12, I = 0.25, - 0.5 R ≥ 1	6.9	4.5	88.6	0.032	0.125	0.032 - 2
	Fluconazole	S ≤ 2, SDD = 4, R ≥ 8	4.9	4.5	90.6	0.125	2	0.032 - 64
	Posaconazole	WBP <sup>d</sup>	0	0	0	0.032	0.064	0.01 - 1
	Itraconazole	S ≤ 0.12, SDD = 0.25, - 0.5 R ≥ 1	12.7	35.3	52	0.064	1	0.032 - 2
	Ketoconazole	WBP	0	0	0	0.032	0.125	0.01 - 1
	Amphotericin B	S ≤ 1, R ≥ 1	19	0	81	0.064	4	0.032 - 8
<i>Candida tropicalis</i>	Caspofungin	S ≤ 0.25, I = 0.5, R ≥ 1	0	4.8	95.2	0.032	0.125	0.032 - 0.5
	Voriconazole	S ≤ 0.12, I = 0.25, - 0.5 R ≥ 1	14.3	14.3	71.4	0.032	1	0.032 - 16
	Fluconazole	S ≤ 2, SDD = 4, R ≥ 8	9.5	0	90.5	0.25	2	0.032 - 64
	Posaconazole	WBP	0	0	0	0.032	0.125	0.032 - 0.25
	Itraconazole	S ≤ 0.12, SDD = 0.25, - 0.5 R ≥ 1	14.3	38.1	47.6	0.125	1	0.032 - 2
	Ketoconazole	WBP	0	0	0	0.032	0.125	0.032 - 16
<i>Candida parapsilosis</i>	Amphotericin B	S ≤ 1, R ≥ 1	0	0	100	0.032	0.032	0.032 - 0.5
	Caspofungin	S ≤ 2, I = 4, R ≥ 8	0	0	100	0.032	0.125	0.032 - 0.25
	Voriconazole	S ≤ 0.12, I = 0.25, - 0.5 R ≥ 1	0	0	100	0.032	0.032	0.032 - 0.032
	Fluconazole	S ≤ 2, SDD = 4, R ≥ 8	0	0	100	0.125	0.25	0.064 - 2
	Posaconazole	WBP	0	0	0	0.032	0.032	0.032 - 0.032
	Itraconazole	S ≤ 0.12, SDD = 0.25, - 0.5 R ≥ 1	0	60	40	0.125	0.5	0.032 - 0.5
	Ketoconazole	WBP	0	0	0	0.032	0.032	0.032 - 0.064
<i>Candida kefyr</i>	Amphotericin B	S ≤ 1, R ≥ 1	0	0	100	0.032	1	0.032 - 1
	Caspofungin	S ≤ 2	0	0	100	0.032	0.125	0.032 - 0.125
	Voriconazole	WBP	0	0	0	0.032	0.032	0.032 - 0.032
	Fluconazole	S ≤ 8, R ≥ 64	0	0	100	0.5	0.5	0.25 - 0.5
	Posaconazole	WBP	0	0	0	0.032	0.032	0.032 - 0.032
	Itraconazole	S ≤ 0.12, R ≥ 1	0	16.7	83.3	0.032	0.125	0.032 - 0.125
	Ketoconazole	WBP	0	0	0	0.032	0.032	0.032 - 0.032

<sup>a</sup> Resistant

<sup>b</sup> Intermediate

<sup>c</sup> Sensitive

<sup>d</sup> This drug doesn't have any breakpoint (without any breakpoint)

Antifungal resistance may occur due to cross-reactivity of azole antifungal agents. Such a trend was reported using FLU and ITR as prophylaxis or treatment in patients, leading to resistant species, and treatment with other azoles failed. Resistance rate to FLU varied in different studies and for *C. glabrata* it was reported as 95% (38/40), 6.2% (1/16) and 10.3% (6/58 with 52/58 susceptible dose dependent) (6, 11, 23). These rates for *C. krusei* were 95.2% (59/62) (6). *Candida*

*krusei* is naturally resistant to antifungal drugs, especially FLU (24). In the current study, ITR resistance rate in *C. albicans* was 12.7% (34.7/273) with dose dependent susceptibility of 35.3% (96/273). This rate was reported as 15.1% (26/172), 5.4% (2/38), 28% (36/117) and 11.9% (18/167) in other studies (6, 11, 15, 25). Itraconazole resistance rates in *C. glabrata* and *C. krusei* in the present study were 77.8% (14/18) and 33.3% (3/10), and in other studies, these were 85% (34/40) and

**Table 3.** Sensitivity Pattern of Uncommon Candida Species Isolated From Clinical Samples in Six University Hospital in Iran

Candida	Antibiotic name	Breakpoints	%R	%I	%S	MIC50	MIC90	MIC Range
<i>Candida famata</i>	Amphotericin B	$S \leq 1, R \geq 1$	5	0	95	0.032	0.125	0.032 - 2
	Caspofungin	$S \leq 2$	0	0	94.7	0.032	0.5	0.032 - 16
	Voriconazole	WBP	0	0	0	0.032	0.25	0.032 - 0.5
	Fluconazole	$S \leq 8, R \geq 64$	0	0	100	0.25	0.5	0.064 - 8
	Posaconazole	WBP	0	0	0	0.032	0.5	0.032 - 1
	Itraconazole	$S \leq 0.12, R \geq 1$	10	35	55	0.032	0.5	0.032 - 1
	Ketoconazole	WBP	0	0	0	0.032	0.032	0.032 - 0.25
<i>Candida glabrata</i>	Amphotericin B	$S \leq 1, R \geq 1$	11.1	0	88.9	0.032	16	0.032 - 16
	Caspofungin	$S \leq 0.12, I = 0.25, R \geq 0.5$	22.2	11.1	66.7	0.032	8	0.032 - 8
	Voriconazole	ECV <sup>a</sup> = 0.5	0	0	0	0.032	4	0.032 - 4
	Fluconazole	SDD $\leq 32, R \geq 64$	11.1	0	88.9	0.5	64	0.064 - 64
	Posaconazole	WBP	0	0	0	0.125	8	0.032 - 8
	Itraconazole	$S \leq 0.12, SDD = 0.25, - 0.5 R \geq 1$	77.8	0	22.2	1	16	0.032 - 16
	Ketoconazole	WBP	0	0	0	0.064	4	0.032 - 4
<i>Candida krusei</i>	Amphotericin B	$S \leq 1, R \geq 1$	30	0	70	1	8	0.032 - 8
	Caspofungin	$S \leq 0.25, I = 0.5, R \geq 1$	33.3	0	66.7	0.125	2	0.032 - 2
	Voriconazole	$S \leq 0.5, I = 1, R \geq 2$	33.3	16.7	50	0.064	2	0.032 - 2
	Fluconazole	NR <sup>b</sup>	0	0	0	32	64	2 - 64
	Posaconazole	WBP	0	0	0	0.125	0.5	0.032 - 0.5
	Itraconazole	$S \leq 0.12, SDD = 0.25, - 0.5 R \geq 1$	33.3	33.3	33.3	0.125	1	0.064 - 1
	Ketoconazole	WBP	0	0	0	0.064	4	0.064 - 4
<i>Candida spp</i> <sup>c</sup>	Amphotericin B	$S \leq 1, R \geq 1$	0	0	100	0.032	0.032	0.032 - 0.032
	Caspofungin	$S \leq 2$	0	0	100	0.032	0.064	0.032 - 0.064
	Voriconazole	WBP	0	0	0	0.032	0.032	0.032 - 0.032
	Fluconazole	$S \leq 8, R \geq 64$	0	0	100	0.25	4	0.125 - 4
	Posaconazole	WBP	0	0	0	0.032	0.032	0.032 - 0.032
	Itraconazole	$S \leq 0.12, R \geq 1$	0	0	80	0.032	0.125	0.032 - 0.125
	Ketoconazole	WBP	0	0	0	0.032	0.032	0.032 - 0.032

<sup>a</sup> Epidemiological cut off value

<sup>b</sup> Naturally resistant (This species is resistant to fluconazole with every MICs)

<sup>c</sup> *Candida dubliniensis, Candida guilliermondii* and *Candida lusitanae*

85.5% (53/62), and 50% (7/14) and 30% (6/18), respectively (6, 15). Voriconazole is a drug of choice for the treatment of filamentous fungi and it has a good activity against fluconazole-resistant *C. glabrata* strains (26). According to literature, low resistance rate of VOR for *Candida* species was reported (6, 14, 27).

Ketoconazole showed significant toxicity as a systemic drug, so it is only available as a topical drug (cream and shampoo) for the treatment of cutaneous fungal infections (26). Posaconazole and KET do not have any break-

point mentioned in CLSI M27-S4 reference (10). Previously, highest MIC values for KET were reported in *C. krusei* (4 µg/mL) (15) and *C. albicans* (32 and 2 µg/mL) (15, 28). The difference could be explained by the sample size and study population, e.g., in immunocompromised cases, like HIV patients, the resistance rate to anti-fungal agents was high (28). In this study, highest MIC90 value for POS was observed in *C. glabrata* (2 µg/mL), as in a previous study (15). In a study from 70 medical centers around the world, the highest MIC90 value for POS was reported in *C. glabrata*

with 2 µg/mL (29).

## 6. Conclusion

Selection of the most appropriate drug and effective treatment for patients is critical in clinical practice. According to this study, *C. albicans* is the most prevalent etiologic agent in infected patients and caspofungin is the most effective antifungal agent. Regional data may not be applicable to other regions and may lead to failure in treatment. Knowledge about etiologic agents and their susceptibility patterns in each region is required for successful treatment of patients.

## Acknowledgments

The authors thank Hassan Khajehei for his help with language copy editing. The authors thank their colleague Zahra Jafarpour, who provided insight and expertise that greatly assisted the research.

## Footnotes

**Authors' Contribution:** Parisa Badiie Study concept and design, Parisa Badiie, Hamid Badali, Kambiz Diba, Rasoul Mohammadi, Hossein Mirhendi, and Mohammad Javad Najafzadeh Acquisition of data, Parisa Badiie and Hadis Jafarian Analysis and interpretation of data, Parisa Badiie and Hadis Jafarian Drafting of the manuscript, Parisa Badiie, Hamid Badali, Kambiz Diba, Rasoul Mohammadi, Hossein Mirhendi and Mohammad Javad Najafzadeh Critical revision of the manuscript for important intellectual content, Parisa Badiie and Hadis Jafarian Statistical analysis, Parisa Badiie and Hadis Jafarian Administrative, technical, and material support, Parisa Badiie Study supervision.

**Conflict of Interest:** The authors declare no conflict of interest.

**Financial Disclosure:** The authors declare no competing financial interests.

## References

- Brown GD, Denning DW, Gow NA, Levitz SM, Netea MG, White TC. Hidden killers: human fungal infections. *Sci Transl Med*. 2012;**4**(165):165rv13. doi: [10.1126/scitranslmed.3004404](https://doi.org/10.1126/scitranslmed.3004404). [PubMed: 23253612].
- Badiie P, Alborzi A, Vojdani R, Shakiba E, Rasouli M, Ravanfar P, et al. Early diagnosis of systemic candidiasis in bone marrow transplant recipients. *Exp Clin Transplant*. 2010;**8**(2):98-103. [PubMed: 20565365].
- Colombo AL, Guimaraes T, Sukienik T, Pasqualotto AC, Andreotti R, Queiroz-Telles F, et al. Prognostic factors and historical trends in the epidemiology of candidemia in critically ill patients: an analysis of five multicenter studies sequentially conducted over a 9-year period. *Intensive Care Med*. 2014;**40**(10):1489-98. doi: [10.1007/s00134-014-3400-y](https://doi.org/10.1007/s00134-014-3400-y). [PubMed: 25082359].
- Lortholary O, Renaudat C, Sitbon K, Madec Y, Denoed-Ndam L, Wolff M, et al. Worrying trends in incidence and mortality of candidemia in intensive care units (Paris area, 2002-2010). *Intensive Care Med*. 2014;**40**(9):1303-12. doi: [10.1007/s00134-014-3408-3](https://doi.org/10.1007/s00134-014-3408-3). [PubMed: 25097069].
- Pfaller MA, Diekema DJ. Epidemiology of invasive mycoses in North America. *Crit Rev Microbiol*. 2010;**36**(1):1-53. doi: [10.3109/10408410903241444](https://doi.org/10.3109/10408410903241444). [PubMed: 20088682].
- Badiie P, Alborzi A, Shakiba E, Farshad S, Japoni A. Susceptibility of Candida species isolated from immunocompromised patients to antifungal agents. *East Mediterr Health J*. 2011;**17**(5):425-30. [PubMed: 21796956].
- Lockhart SR, Messer SA, Gherna M, Bishop JA, Merz WG, Pfaller MA, et al. Identification of Candida nivariensis and Candida bracarenensis in a large global collection of Candida glabrata isolates: comparison to the literature. *J Clin Microbiol*. 2009;**47**(4):1216-7. doi: [10.1128/JCM.02315-08](https://doi.org/10.1128/JCM.02315-08). [PubMed: 19193845].
- Horn DL, Neofytos D, Anaissie EJ, Fishman JA, Steinbach WJ, Olyaei AJ, et al. Epidemiology and outcomes of candidemia in 2019 patients: data from the prospective antifungal therapy alliance registry. *Clin Infect Dis*. 2009;**48**(12):1695-703. doi: [10.1086/599039](https://doi.org/10.1086/599039). [PubMed: 19441981].
- CLSI reference method for broth dilution antifungal susceptibility testing of yeasts; approved standard third edition. CLSI document M27-A3. Wayne, PA: Clinical and Laboratory Standards Institute; 2008.
- CLSI reference method for broth dilution antifungal susceptibility testing of yeasts; fourth informational supplement. CLSI document M27-S4. Wayne, PA: Clinical and Laboratory Standards Institute; 2012.
- Shokohi T, Bandalizadeh Z, Hedayati MT, Mayahi S. In vitro antifungal susceptibility of Candida species isolated from oropharyngeal lesions of patients with cancer to some antifungal agents. *Jundishapur J Microbiol*. 2011;**4**(Supplement 1):S19-S26.
- Badiie P, Amirghofran AA, Nour MG, Shafa M, Nemati MH. Incidence and outcome of documented fungal endocarditis. *Int Cardiovasc Res J*. 2014 Dec;**8**(4):152. doi: [25614858](https://doi.org/10.25614858).
- Ghasemi Z, Hashemi SJ, Rezaei S, Kordbache P, Khosravi M, Mortazavi H, et al. Molecular Analysis of Candida species with Emphasis on Pre-disposing Factors in Cutaneous Candidiasis Patients. *Jundishapur J Microbiol*. 2016;**10**(2):e41030. doi: [10.5812/jjm.41030](https://doi.org/10.5812/jjm.41030).
- Wang H, Xiao M, Chen SC, Kong F, Sun ZY, Liao K, et al. In vitro susceptibilities of yeast species to fluconazole and voriconazole as determined by the 2010 National China Hospital Invasive Fungal Surveillance Net (CHIF-NET) study. *J Clin Microbiol*. 2012;**50**(12):3952-9. doi: [10.1128/JCM.01130-12](https://doi.org/10.1128/JCM.01130-12). [PubMed: 23035204].
- Haddadi P, Zareifar S, Badiie P, Alborzi A, Mokhtari M, Zomorodian K, et al. Yeast colonization and drug susceptibility pattern in the pediatric patients with neutropenia. *Jundishapur J Microbiol*. 2014;**7**(9):e11858. doi: [10.5812/jjm.11858](https://doi.org/10.5812/jjm.11858). [PubMed: 25485060].
- Doi AM, Pignatari AC, Edmond MB, Marra AR, Camargo LF, Siqueira RA, et al. Epidemiology and Microbiologic Characterization of Nosocomial Candidemia from a Brazilian National Surveillance Program. *PLoS One*. 2016;**11**(1):e0146909. doi: [10.1371/journal.pone.0146909](https://doi.org/10.1371/journal.pone.0146909). [PubMed: 26808778].
- Pakshir K, Zomorodian K, Zakaei A, Motamedi M, Rahimi Ghiasi M, Karamitalab M. Molecular identification and in-vitro antifungal susceptibility testing of Candida species isolated from patients with onychomycosis. *Curr Med Mycol*. 2015;**1**(4):26-32. doi: [10.18869/acadpub.cmm.1.4.26](https://doi.org/10.18869/acadpub.cmm.1.4.26). [PubMed: 28681001].
- Tortorano AM, Prigitano A, Lazzarini C, Passera M, Deiana ML, Cavinato S, et al. A 1-year prospective survey of candidemia in Italy and changing epidemiology over one decade. *Infection*. 2013;**41**(3):655-62. doi: [10.1007/s15010-013-0455-6](https://doi.org/10.1007/s15010-013-0455-6). [PubMed: 23559357].

19. Zomorodian K, Rahimi MJ, Pakshir K, Motamedi M, Ghiasi MR, Reza-shah H. Determination of antifungal susceptibility patterns among the clinical isolates of *Candida* species. *J Glob Infect Dis*. 2011;**3**(4):357-60. doi: [10.4103/0974-777X.91059](https://doi.org/10.4103/0974-777X.91059). [PubMed: [2223999](https://pubmed.ncbi.nlm.nih.gov/2223999/)].
20. Parisa B, Alborzi A, Shakiba E, Ziyaeyan M, Rasuli M. Molecular Identification and In-Vitro Susceptibility of *Candida albicans* and *C. dubliniensis* Isolated from Immu-nocompromised Patients. *Iran Red Crescent Med J*. 2009;**2009**(4):391-7.
21. Espinel-Ingroff A, Arendrup MC, Pfaller MA, Bonfietti LX, Bustamante B, Canton E, et al. Interlaboratory variability of Caspofungin MICs for *Candida* spp. Using CLSI and EUCAST methods: should the clinical laboratory be testing this agent? *Antimicrob Agents Chemother*. 2013;**57**(12):5836-42. doi: [10.1128/AAC.01519-13](https://doi.org/10.1128/AAC.01519-13). [PubMed: [24018263](https://pubmed.ncbi.nlm.nih.gov/24018263/)].
22. Zarei Mahmoudabadi A, Rezaei-Matehkolaei A, Ghanavati F. The susceptibility patterns of *Candida* species isolated from urine samples to posaconazole and caspofungin. *Jundishapur J Microbiol*. 2015;**8**(3):e24298. doi: [10.5812/jjm.24298](https://doi.org/10.5812/jjm.24298). [PubMed: [25861442](https://pubmed.ncbi.nlm.nih.gov/25861442/)].
23. Won EJ, Shin JH, Choi MJ, Lee WG, Park YJ, Uh Y, et al. Antifungal susceptibilities of bloodstream isolates of *Candida* species from nine hospitals in Korea: application of new antifungal breakpoints and relationship to antifungal usage. *PLoS One*. 2015;**10**(2):e0118770. doi: [10.1371/journal.pone.0118770](https://doi.org/10.1371/journal.pone.0118770). [PubMed: [25706866](https://pubmed.ncbi.nlm.nih.gov/25706866/)].
24. Kanafani ZA, Perfect JR. Antimicrobial resistance: resistance to antifungal agents: mechanisms and clinical impact. *Clin Infect Dis*. 2008;**46**(1):120-8. doi: [10.1086/524071](https://doi.org/10.1086/524071). [PubMed: [18171227](https://pubmed.ncbi.nlm.nih.gov/18171227/)].
25. Bassetti M, Taramasso L, Nicco E, Molinari MP, Mussap M, Viscoli C. Epidemiology, species distribution, antifungal susceptibility and outcome of nosocomial candidemia in a tertiary care hospital in Italy. *PLoS One*. 2011;**6**(9):e24198. doi: [10.1371/journal.pone.0024198](https://doi.org/10.1371/journal.pone.0024198). [PubMed: [21935385](https://pubmed.ncbi.nlm.nih.gov/21935385/)].
26. Nett JE, Andes DR. Antifungal Agents: Spectrum of Activity, Pharmacology, and Clinical Indications. *Infect Dis Clin North Am*. 2016;**30**(1):51-83. doi: [10.1016/j.idc.2015.10.012](https://doi.org/10.1016/j.idc.2015.10.012). [PubMed: [26739608](https://pubmed.ncbi.nlm.nih.gov/26739608/)].
27. Lyon GM, Karatela S, Sunay S, Adiri Y. Candida Surveillance Study I. Antifungal susceptibility testing of *Candida* isolates from the Candida surveillance study. *J Clin Microbiol*. 2010;**48**(4):1270-5. doi: [10.1128/JCM.02363-09](https://doi.org/10.1128/JCM.02363-09). [PubMed: [20129963](https://pubmed.ncbi.nlm.nih.gov/20129963/)].
28. Katirae F, Teifoori F, Soltani M. Emergence of azole-resistant *Candida* species in AIDS patients with oropharyngeal candidiasis in Iran. *Curr Med Mycol*. 2015;**1**(3):11-6. doi: [10.18869/acadpub.cmm.1.3.11](https://doi.org/10.18869/acadpub.cmm.1.3.11). [PubMed: [28680991](https://pubmed.ncbi.nlm.nih.gov/28680991/)].
29. Pfaller MA, Rhomberg PR, Messer SA, Jones RN, Castanheira M. Isavuconazole, micafungin, and 8 comparator antifungal agents' susceptibility profiles for common and uncommon opportunistic fungi collected in 2013: temporal analysis of antifungal drug resistance using CLSI species-specific clinical breakpoints and proposed epidemiological cutoff values. *Diagn Microbiol Infect Dis*. 2015;**82**(4):303-13. doi: [10.1016/j.diagmicrobio.2015.04.008](https://doi.org/10.1016/j.diagmicrobio.2015.04.008). [PubMed: [25986029](https://pubmed.ncbi.nlm.nih.gov/25986029/)].