



Antiviral Activity of Thiazolide Derivatives Against Dengue Virus in Huh-7 Cell Line

Farkhanda Yasmin,¹ Tahir Yaqub,^{1,*} Idrees Khan,² Wasim Shahzad,¹ Abu Saeed Hashmi,¹ Zarfishan Tahir,^{1,3} Nadia Mukhtar,¹ and Sajid Umar⁴

¹Institute of Biochemistry and Biotechnology, University of Veterinary and Animal Sciences, Lahore, Pakistan

²Department of Microbiology, University of Veterinary and Animal Sciences, Lahore, Pakistan

³National Centre of Excellence in Molecular Biology, Lahore, Pakistan

⁴Department of Pathobiology, PMAS Arid Agriculture University Rawalpindi, Pakistan

*Corresponding author: Tahir Yaqub, University of Veterinary and Animal Sciences, Out Fall Road, Lahore, Pakistan. Tel: +92-3006950418, E-mail: tahiryaqub@uvas.edu.pk

Received 2017 April 14; Revised 2017 October 07; Accepted 2017 November 05.

Abstract

Background: Dengue is one of the main health problems worldwide with dramatic increases in reported cases in the last few decades. Now, dengue is a major threat to half of the world's population and approximately 50 million infections occur every year.

Objectives: The major aim of this study was to screen out different synthetic antiviral compounds against dengue virus. As treatments used for dengue virus infections are nonspecific and effectless, it is necessary to find specific antiviral compounds that may be helpful in the treatment of dengue fever. In cell culture models, thiazolides are active against anaerobic bacteria, protozoa, and a range of viruses.

Methods: Samples from positive dengue fever patients were included in the study. For dengue serotype-2, serum samples were further confirmed by serotyping protocol.

Results: After toxicological analysis, a sharp significant antiviral response was shown by compounds A and B decreasing the viral titer of serotype 2 as 46% and 53%, respectively. A greater antiviral activity was indicated by compound "B" against NS3 gene as compared to compound "A". Against NS3 gene, a comparable antiviral activity was observed for compounds "A" and "B".

Conclusions: From our results, it can be concluded that compounds A and B (Thiazolides) as dengue antiviral drugs can reduce viral load in patients if their inhibitory effects are reproduced in vivo. To identify the specific active ingredients, further studies are required by using various techniques to determine the efficacy of these compounds in animal model.

Keywords: Dengue Virus, Dengue Fever, Human Hepatoma Cell Line, Thiazolide Derivatives, Antiviral Activity

1. Background

Dengue virus (DENV) is transmitted to human by *Aedes* mosquitoes; it causes dengue fever (DF) and dengue hemorrhagic fever (DHF), which is a self-limiting febrile illness (1, 2). Dengue fever is relatively mild, but DHF leads to the life-threatening dengue shock syndrome (DSS). The world health organization (WHO) considers dengue as a major global public health challenge in tropic and subtropic regions. Dengue was seen a 30-fold upsurge worldwide between 1960 and 2010 due to increased population growth rate, global warming, unplanned urbanization, inefficient mosquito control, frequent air travel, and lack of health care facilities (2-5). Two and a half billion people reside in dengue-endemic regions and 400 million infections roughly occur per year, with a mortality rate surpassing 5% - 20% in some areas (6, 7). Dengue infection affects more than 100 countries, including European countries and the

United States (USA) (7). In Pakistan, all the serotypes of dengue virus are reported to present especially during the monsoon period, making it a dengue hyper endemic region (8, 9).

The dengue virus, a member of the genus *Flavivirus* of the family *Flaviviridae*, is an arthropod-borne virus that includes four different serotypes (DENV1-4) (4, 8, 10). These viruses are enveloped and have a single-stranded positive-sense RNA genome of approximately 11 kb (7). A single long open reading frame of the viral RNA encodes a polyprotein that is processed by cellular and viral proteases into three structural (C, prM, and E) and seven nonstructural (NS) proteins (NS1, NS2A, NS2B, NS3, NS4A, NS4B, and NS5). The structural proteins form the virus particles and play roles in receptor binding, virus fusion, and virion assembly. Nonstructural proteins are responsible for the replication of viral genome and evasion from host immunity. Many antiviral compounds have been reported to inhibit

DENV replication *in vitro* and *in vivo* (6, 11, 12). However, no approved antiviral drug is yet available for the treatment of DENV infectious disease. Thiazolidine is a class of compounds that merits special attention because it belongs to a group of substances with activity in medicinal chemistry. This nucleus is associated with antibacterial, antifungal, antiviral, antituberculosis, anticancer, and antiparasite biological activities.

2. Objectives

The present study was designed to study the antiviral activity of thiazolidines against NS3 gene (non-structural) of a dengue serotype-2 isolate (Pakistani isolate). These findings suggest that thiazolidines may inhibit DENV replication by targeting the NS3 protein.

3. Methods

3.1. Ethics Statement

The study was approved by the institutional ethics review committee of the centre for applied molecular biology (CAMB), Lahore, Pakistan, and institutional review board (IRB), general hospital, Lahore. All experiments were carried out according to the guidelines of the committee (CEMB/IRB XGT3456). Before sample collection, a written informed consent was taken from all the patients or their attendant.

3.2. Collection of Serum Sample

For each patient, all the demographics, clinical data, and other relevant information were recorded in separate data sheets. Serum samples were collected from 50 patients suffering from dengue serotype-2 and stored at -20°C until processed for further study.

3.3. Serotype Determination

Only those samples from DF patients were considered which were declared positive according to the WHO criteria. For dengue serotype-2, serum samples were further confirmed by serotyping protocol developed by CEMB (13). The selected DENV-2 samples were collected during three different years (2011, 2012, and 2013) and processed for further experiments. For serotyping analysis, nested PCR was used. For each sample, the amplified bands were gel eluted and sequence analysis was performed. For serotyping analysis, junction of C-prM gene of dengue virus isolates was selected. For serotype 2, the amplified product was 403 base pairs (bp) and for serotype 3, it was 453 bp in length.

3.4. Cell Line

Human hepatoma Huh 7 cells was obtained from CEMB, University of the Punjab, Lahore, and maintained in Dulbecco's modified Eagle's medium (DMEM; life technologies, USA) supplemented with 10% FBS and antibiotics under 5% CO_2 at 37°C . Hybridoma-producing anti-E antibody (HB- 112) was maintained in Roswell Park Memorial Institute 1640 medium (RPMI1640; Life Technologies, USA) supplemented with 10% FBS and antibiotics under 5% CO_2 at 37°C , and its supernatant was centrifuged at 1000 g for 5 minutes.

3.5. Thiazolide Derivatives

Two Thiazolidines derivatives (A and B) obtained from NIBGE Faisalabad (Voucher number 234) were dissolved in a solvent dimethyl sulfoxide (DMSO), following modified procedure described previously (11). Chemical structures of the thiazolidines derivatives A (DD2) and B (DD10) used in this study are shown in Figure 1.




Figure 1. Chemical Structures of the Thiazolidines Derivatives A (DD2) and B (DD10) Used in This Study

3.6. Splitting of Cells

Using DMEM (life technologies, USA), the cell line was grown accompanied by $100\mu\text{g}/\text{mL}$ penicillin, $100\mu\text{g}/\text{mL}$ streptomycin, and 10% fetal bovine serum (FBS) (Wisent INC., Montreal, Canada) incubated at 37°C with 5% CO_2 . Huh-7 is highly efficient for DENV replication. For toxicological analysis using DMEM media, the Huh-7 cell line was

grown with the suitable antibiotic, 10% FBS, and CO₂, split and then plated. 5 mL of DMEM with antibiotic was added to dislodge the cells into single cell suspension (20 to 30 times). Then, 10 μ L of the mixture was taken and the cells were counted on haemocytometer. With different concentrations of compounds, these cells were treated; the first well was remained as control while different concentrations of the compound from lowest to highest were added to the other wells. The cells were trypsinized after 24 hours, and were counted on haemocytometer.

3.7. Dengue RNA Detection in Huh -7 Cells

For the detection of Huh-7 cells expressing NS3 gene, total RNA from grown cell line of NS3 gene was extracted. Through gene specific primers (NS3F GCAAGCTTGCCATGGCCGCTGGAGTATTGTCGGC, NS3R - GCGCGCCGCTTTCTTCCACTGCAAACTCTTGTTTC), the RNA of NS3 gene was reverse transcribed by RT PCR into cDNA by utilizing MML-V. Reverse transcriptase un-transfected Huh-7 cells were kept as negative (-ve) control. RT-PCR reaction was done using gene specific primers for NS3 gene. The approximate band of NS3 (1904 bp) was observed compared to control Huh-7 cells.

3.8. Anti-Dengue Analysis of Thiazolides in Cells

After 24 hours, incubation was done at 37°C in an atmosphere of 5% CO₂ and dengue infected serum was added. To check the antiviral activity of compounds after 24 hours incubation, total RNA was extracted from dengue serum infected Huh-7 cells using Pure-script RNA Isolation kit (life technologies, USA) (12, 14-18). Through quantitative RT-PCR, viral titer was determined. The dengue RNA in human plasma was checked through the quantitative assay using the provided procedure in the kit Maxima TMSYBER Green/ROX qPCR Master Mix kit (Fermantas, Hanover, MD, USA).

4. Results

4.1. Toxicological Analysis of Compounds

In Huh-7 cells, toxicity of compounds was determined. The cell culture was treated with variable concentrations of the thiazolides derivatives. Trypsinization of cells was done after 24 hours of treatment, and using haemocytometer, cells were counted. At different concentrations, the cells viability was checked through trypan blue exclusion test. Non-toxic concentration was the concentration that gave 60 - 70 percent of the cells survival. DMSO was used as control. The concentration of compounds up to 40 - 50 μ M/mL was non-toxic to the cells (Table 1).

4.2. Antiviral Activity

In huh-7 cells, the activity of antiviral compounds was checked at different concentrations ranging from 5 to 90 μ M/mL. 40 μ M dose of the compounds was found nontoxic to the cells. DMSO was used as a negative control and for positive control, dengue infected sera were used. Before the addition of compounds, the cells were infested with positive serum of dengue serotype 2. Viral titer of infected Huh-7 cell line was reduced up to 46% by using compound "A" and 53% by treatment with compound "B" compared to the control samples. Decrease in dengue RNA was seen up to 45% for compound A and 53% for compound B, respectively. It indicates that compound "B" is comparatively more active against dengue, which proposes its medicinal importance (Figure 2).

5. Discussion

To control the growing cases of dengue disease in the endemic region, new treatments are urgently needed. In coming years, Pakistan is at high risk of dengue virus epidemic, since it has a history of dengue with the 2011 outbreak in Lahore, which recorded more than 15,000 dengue cases with high mortality rate (19). Different strategies have been discovered, and one of the best approaches is the use of antiviral compounds to inhibit the virus replication against dengue protease gene. The identification and subsequent use of anti-dengue active compounds are potentially useful for developing new antiviral treatments against dengue. Currently, vaccine development is underway against DENV. The practical applications of these vaccines are always a big challenge, such as the potential of antibody-dependent enhancement (ADE) and non-neutralizing antibodies produced by the human body during a primary infection of dengue virus. The current treatment used for dengue virus infections is nonspecific and effectless. Therefore, the current study was designed to find specific antiviral compounds that may be helpful in the treatment of dengue fever.

The functional inhibition of viral NS proteins is mainly based on the anti-flavivirus drug development (10, 20). In fact, the NS3 protein is multifunctional enzyme, which has three important regions including serine protease, helicase, and a nucleotide triphosphatase (7, 21). Structural analysis of NS3 protein has revealed that a fragment of 180 amino acids at N-terminus of NS3 is homologous to serine protease active sites and therefore this region is important for proteolytic cleavage and processing of dengue polyprotein (3, 22) as the precursor protein is cleaved using the NS2b-NS3 protease of virus and also with the proteases of the host cell to release the individual proteins (23). The

Table 1. Toxicity Analysis of Different Concentrations of Thiazolide Derivatives in Huh-7 Cells

S.No	Concentration of Compounds	Thiazolide Derivative A		Thiazolide Derivative B	
		Average No of Cells, Count /mL	Standard Error	Average No of Cells, Count /mL	Standard Error
1	0	6.215×10^6	8.459×10^4	1.905×10^6	1.686×10^4
2	5	5.998×10^6	6.840×10^4	1.551×10^6	2.335×10^4
3	15	4.265×10^6	4.692×10^4	1.440×10^6	2.002×10^4
4	30	3.015×10^6	3.818×10^4	1.024×10^6	1.761×10^4
5	45	2.742×10^6	2.389×10^4	7.127×10^5	9.917×10^3
6	60	1.784×10^6	3.289×10^4	2.851×10^5	3.668×10^3
7	75	1.684×10^6	1.935×10^4	1.535×10^5	2.563×10^3
8	90	1.502×10^6	2.038×10^4	1.308×10^5	2.233×10^3




Figure 2. Antiviral analysis of compounds A and B in Huh-7 cell line; Infected Huh-7 cells were treated with different concentrations of compounds A and B. NS3 genome levels of dengue virus are presented as a percentage in comparison with the levels of dengue RNA in cells incubated with control.

virus replication complex is formed by NS3 and NS5 proteins and replication complex assembly is on the intracellular membrane for the amplification of the viral genome (24, 25). Moreover, the NS3 structural studies are helpful to develop the novel antiviral compounds that target mainly dengue enzymes playing important roles in viral replication.

The present study describes antiviral activity of synthetic active antiviral compounds against dengue in Huh-7 cell line. Different solvents were used to determine the

compounds solubility such as DMSO and water. Huh-7 cell line is significant in the estimation of antiviral activity (12, 18). Antiviral response was directly indicated by viral titer. The concentrations of 5 µg/mL to 50 µg/mL with volume of 5 - 90 µM of compounds were used. In Huh-7 cell, cytotoxicity of synthetic compounds with different concentrations was examined. With the administration of 30 µM (50 µg of synthetic compounds/ml), the compounds were non-toxic. Significant cytotoxicity was not shown by the cells. Using haemocytometer and trypanblue stain method, cell

viability was calculated. The current study delivers investigational data for the treatment of DENV infection. In this study, anti-dengue activity of selected thiazolides was checked in Huh-7 cells, in order to provide experimental data for clinical treatment of dengue. Adopting the same strategy, antiviral screening of thiazolides compound was performed in infected Huh-7 cells with known titer of virus and serotype.

A sharp significant antiviral response was shown by the compounds "A and B" decreasing the viral titer of serotype 2 as 46% and 53%, respectively, after 48 hours of incubation. Furthermore, compound "B" presented a greater antiviral response against NS3 gene as compared to compound "A". Against NS3 gene, a comparable antiviral activity was observed for compounds "A" and "B". The viral titer increased against the antiviral compounds. Compound "B" showed enhanced antiviral activity against NS3 gene as compared to compound "A". To monitor anti-DENV compounds, many viral replicon-based studies were carried out (4, 12, 15-18). The current study is different from a previous report on DENV-2 strain replicon that was constructed in different cells, in which the genetic material of whole virus is used instead of active protease gene, resulting in a less clarification of mechanism. Similarly, Qing Min et al. (16) also performed antiviral compound analysis through the infection of Vero and Huh-7 cell lines with dengue. DENV replicon constructs showed higher rate of replication in human Huh7.5 cells than in Chinese hamster BHK21 cells (1). A study was carried out to explore anti dengue effects in plants (26). The positive results showed that compounds A and B have potent DENV antiviral properties. Thus, A and B are new antiviral compounds that might be significant compounds in the treatment of dengue fever.

6. Conclusions

This may suggest that by using compounds A and B (Thiazolide derivatives) as antiviral drugs, dengue virus viral load can be reduced in patients if their antiviral activities are reproduced in vivo. To find out the specific active ingredients, further studies are needed by using various techniques to determine the efficacy of these compounds in animal model.

Acknowledgments

We are thankful to the laboratory staff of institute of biochemistry and biotechnology, university of veterinary and animal sciences, Lahore, Pakistan, for their technical help throughout the study. Farkhanda Yasmin's PhD scholarship was supported by the higher education commission, Pakistan.

Footnotes

Authors' Contribution: Farkhanda Yasmin collected all the required data, processed the samples, and drafted the manuscript; Tahir Yaqub and Nadia Mukhtar designed the study and analyzed the data; Idrees Khan, Wasim Shahzad, Abu Saeed Hashmi, and Zarfishan Tahir interpreted the data; Sajid Umar critically and substantially revised the manuscript; all authors read and approved the final manuscript.

Conflict of Interest: Authors declare that there is no conflict of interest in this study.

Funding/Support: Financial support for this study was provided by the center of excellence in molecular biology, Lahore, and partially by higher education commission of Pakistan.

References

1. Yang CC, Tsai MH, Hu HS, Pu SY, Wu RH, Wu SH, et al. Characterization of an efficient dengue virus replicon for development of assays of discovery of small molecules against dengue virus. *Antiviral Res.* 2013;**98**(2):228-41. doi: [10.1016/j.antiviral.2013.03.001](https://doi.org/10.1016/j.antiviral.2013.03.001).
2. Das S, Pingle MR, Munoz-Jordan J, Rundell MS, Rondini S, Granger K, et al. Detection and serotyping of dengue virus in serum samples by multiplex reverse transcriptase PCR-ligase detection reaction assay. *J Clin Microbiol.* 2008;**46**(10):3276-84. doi: [10.1128/JCM.00163-08](https://doi.org/10.1128/JCM.00163-08). [PubMed: [18685000](https://pubmed.ncbi.nlm.nih.gov/18685000/)].
3. Bazan JF, Fletterick RJ. Detection of a trypsin-like serine protease domain in flaviviruses and pestiviruses. *Virology.* 1989;**171**(2):637-9. [PubMed: [2548336](https://pubmed.ncbi.nlm.nih.gov/2548336/)].
4. Gibbons RV, Vaughn DW. Dengue: an escalating problem. *BMJ.* 2002;**324**(7353):1563-6. [PubMed: [12089096](https://pubmed.ncbi.nlm.nih.gov/12089096/)].
5. Gubler DJ. Dengue and dengue hemorrhagic fever. *Clin Microbiol Rev.* 1998;**11**(3):480-96. [PubMed: [9665979](https://pubmed.ncbi.nlm.nih.gov/9665979/)].
6. Leardkamolkarn V, Srigulpanit W, Phurimsak C, Kumkate S, Himakoun L, Sripanidkulchai B. The inhibitory actions of Houltuynia cordataaqueous extract on dengue virus and dengue-infected cells. *J Food Biochem.* 2012;**26**:86-92.
7. Wilder-Smith A, Ooi EE, Vasudevan SG, Gubler DJ. Update on dengue: epidemiology, virus evolution, antiviral drugs, and vaccine development. *Curr Infect Dis Rep.* 2010;**12**(3):157-64. doi: [10.1007/s11908-010-0102-7](https://doi.org/10.1007/s11908-010-0102-7). [PubMed: [21308524](https://pubmed.ncbi.nlm.nih.gov/21308524/)].
8. Jahan F. Dengue Fever (DF) in Pakistan. *Asia Pac Fam Med.* 2011;**10**(1):1. doi: [10.1186/1447-056X-10-1](https://doi.org/10.1186/1447-056X-10-1). [PubMed: [21349169](https://pubmed.ncbi.nlm.nih.gov/21349169/)].
9. Khan E, Kismet M, Khan N, Nasir A, Ayub S, Hasan R. Demographic and clinical features of dengue fever in Pakistan from 2003-2007: a retrospective cross-sectional study. *PLoS One.* 2010;**5**(9):12505. doi: [10.1371/journal.pone.0012505](https://doi.org/10.1371/journal.pone.0012505). [PubMed: [20856935](https://pubmed.ncbi.nlm.nih.gov/20856935/)].
10. Bollati M, Alvarez K, Assenberg R, Baronti C, Canard B, Cook S, et al. Structure and functionality in flavivirus NS-proteins: perspectives for drug design. *Antiviral Res.* 2010;**87**(2):125-48. doi: [10.1016/j.antiviral.2009.11.009](https://doi.org/10.1016/j.antiviral.2009.11.009). [PubMed: [19945487](https://pubmed.ncbi.nlm.nih.gov/19945487/)].
11. Korba BE, Montero AB, Farrar K, Gaye K, Mukerjee S, Ayers MS, et al. Nitazoxanide, tizoxanide and other thiazolides are potent inhibitors of hepatitis B virus and hepatitis C virus replication. *Antiviral Res.* 2008;**77**(1):56-63. doi: [10.1016/j.antiviral.2007.08.005](https://doi.org/10.1016/j.antiviral.2007.08.005). [PubMed: [17888524](https://pubmed.ncbi.nlm.nih.gov/17888524/)].
12. Shum D, Smith JL, Hirsch AJ, Bhinder B, Radu C, Stein DA, et al. High-content assay to identify inhibitors of dengue virus infection.

- Assay Drug Dev Technol.* 2010;**8**(5):553-70. doi: [10.1089/adt.2010.0321](https://doi.org/10.1089/adt.2010.0321). [PubMed: [20973722](https://pubmed.ncbi.nlm.nih.gov/20973722/)].
13. Fatima Z, Idrees M, Bajwa MA, Tahir Z, Ullah O, Zia MQ, et al. Serotype and genotype analysis of dengue virus by sequencing followed by phylogenetic analysis using samples from three mini outbreaks-2007-2009 in Pakistan. *BMC Microbiol.* 2011;**11**:200. doi: [10.1186/1471-2180-11-200](https://doi.org/10.1186/1471-2180-11-200). [PubMed: [21906394](https://pubmed.ncbi.nlm.nih.gov/21906394/)].
 14. Hsu YC, Chen NC, Chen PC, Wang CC, Cheng WC, Wu HN. Identification of a small-molecule inhibitor of dengue virus using a replication system. *Arch Virol.* 2012;**157**(4):681-8. doi: [10.1007/s00705-012-1224-z](https://doi.org/10.1007/s00705-012-1224-z). [PubMed: [22249364](https://pubmed.ncbi.nlm.nih.gov/22249364/)].
 15. Masse N, Davidson A, Ferron F, Alvarez K, Jacobs M, Romette JL, et al. Dengue virus replicons: production of an interserotypic chimera and cell lines from different species, and establishment of a cell-based fluorescent assay to screen inhibitors, validated by the evaluation of ribavirin's activity. *Antiviral Res.* 2010;**86**(3):296-305. doi: [10.1016/j.antiviral.2010.03.010](https://doi.org/10.1016/j.antiviral.2010.03.010). [PubMed: [20307577](https://pubmed.ncbi.nlm.nih.gov/20307577/)].
 16. Qing M, Liu W, Yuan Z, Gu F, Shi PY. A high-throughput assay using dengue-1 virus-like particles for drug discovery. *Antiviral Res.* 2010;**86**(2):163-71. doi: [10.1016/j.antiviral.2010.02.313](https://doi.org/10.1016/j.antiviral.2010.02.313). [PubMed: [20153777](https://pubmed.ncbi.nlm.nih.gov/20153777/)].
 17. Noble CG, Chen YL, Dong H, Gu F, Lim SP, Schul W, et al. Strategies for development of Dengue virus inhibitors. *Antiviral Res.* 2010;**85**(3):450-62. doi: [10.1016/j.antiviral.2009.12.011](https://doi.org/10.1016/j.antiviral.2009.12.011). [PubMed: [20060421](https://pubmed.ncbi.nlm.nih.gov/20060421/)].
 18. Wengler G. The carboxy-terminal part of the NS 3 protein of the West Nile Flavivirus can be isolated as a soluble protein after proteolytic cleavage and represents an RNA-stimulated NTPase. *Virology.* 1991;**184**(2):707-15. doi: [10.1016/0042-6822\(91\)90440-m](https://doi.org/10.1016/0042-6822(91)90440-m).
 19. Rai MA. Epidemic: Control of dengue fever in Pakistan. *Nature.* 2011;**479**(7371):41. doi: [10.1038/479041d](https://doi.org/10.1038/479041d). [PubMed: [22051664](https://pubmed.ncbi.nlm.nih.gov/22051664/)].
 20. Lim SP, Wang QY, Noble CG, Chen YL, Dong H, Zou B, et al. Ten years of dengue drug discovery: progress and prospects. *Antiviral Res.* 2013;**100**(2):500-19. doi: [10.1016/j.antiviral.2013.09.013](https://doi.org/10.1016/j.antiviral.2013.09.013). [PubMed: [24076358](https://pubmed.ncbi.nlm.nih.gov/24076358/)].
 21. Warrener P, Tamura JK, Collett MS. RNA-stimulated NTPase activity associated with yellow fever virus NS3 protein expressed in bacteria. *J Virol.* 1993;**67**(2):989-96. [PubMed: [8380474](https://pubmed.ncbi.nlm.nih.gov/8380474/)].
 22. Gorbalenya AE, Donchenko AP, Koonin EV, Blinov VM. N-terminal domains of putative helicases of flavi- and pestiviruses may be serine proteases. *Nucleic Acids Res.* 1989;**17**(10):3889-97. [PubMed: [2543956](https://pubmed.ncbi.nlm.nih.gov/2543956/)].
 23. Lindenbach BD, Pragai BM, Montserret R, Beran RK, Pyle AM, Penin F, et al. The C terminus of hepatitis C virus NS4A encodes an electrostatic switch that regulates NS5A hyperphosphorylation and viral replication. *J Virol.* 2007;**81**(17):8905-18. doi: [10.1128/JVI.00937-07](https://doi.org/10.1128/JVI.00937-07). [PubMed: [17581983](https://pubmed.ncbi.nlm.nih.gov/17581983/)].
 24. Murray CL, Jones CT, Rice CM. Architects of assembly: roles of Flaviviridae non-structural proteins in virion morphogenesis. *Nat Rev Microbiol.* 2008;**6**(9):699-708. doi: [10.1038/nrmicro1928](https://doi.org/10.1038/nrmicro1928). [PubMed: [18587411](https://pubmed.ncbi.nlm.nih.gov/18587411/)].
 25. Paul D, Bartenschlager R. Architecture and biogenesis of plus-strand RNA virus replication factories. *World J Virol.* 2013;**2**(2):32-48. doi: [10.5501/wjv.v2.i2.32](https://doi.org/10.5501/wjv.v2.i2.32). [PubMed: [24175228](https://pubmed.ncbi.nlm.nih.gov/24175228/)].
 26. Rothan HA, Zulqarnain M, Ammar YA, Tan EC, Rahman NA, Yusof R. Screening of antiviral activities in medicinal plants extracts against dengue virus using dengue NS2B-NS3 protease assay. *Trop Biomed.* 2014;**31**(2):286-96. [PubMed: [25134897](https://pubmed.ncbi.nlm.nih.gov/25134897/)].