

An updated conspectus of grasses of Punjab (India)

Amarjit Singh Soodan *, Ashok Kumar and Sushant Sharma

Department of Botanical and Environmental Sciences, Guru Nanak Dev University, Amritsar, Punjab, India

Abstract

Poaceae is the fourth largest family of the flowering plants. It includes about 700-800 genera and 11000-13000 species distributed worldwide. The family has unmatched ecological and economic importance. With its origin in the early cretaceous and major diversification in the mid Cenozoic, the family at present covers nearly a fifth of land surface and occurs in nearly all the habitats of the world. With a sub tropical ecology and an agrarian economy of Punjab, grasses comprise the most significant group in the region. Despite an overwhelming significance, taxonomic studies in grasses have not received sufficient attention in the region. It is only in the work of Sharma and Khosla (1989) that grass species have been classified into subfamilies and tribes. However, after the establishment of the Grass Phylogeny Working Group (GPWG) the world has witnessed a renaissance in grass systematics. But, India remains an 'undercollected' country as far as grass diversity is concerned. Our work on the exploration and systematics of the grass flora of the region is an effort to consolidate and update the information on the diversity of grasses of the studied area. The present studies have brought the cumulative species number to 192 including seven new reports. The species representation of subfamilies is: Aristidoideae (5) Arundinoideae (5) Bambusoideae (4), Centothecoideae (1), Chloridoideae (55) Erhartoideae (3), Panicoideae (98) and Pooideae (21).

Key words: Cumulative, India, Poaceae, Punjab, Systematics, Taxonomy, Tropical

Introduction

The grass family Poaceae (R. Br.) Barnh. is the fourth largest family of the flowering plants. It includes about 700-800 genera and 11000-13000 species distributed worldwide (Clayton and Renvoize, 1986; Watson and Dallwitz, 1999). In the classification of grasses proposed by Grass Phylogeny Working Group (GPWG) the genera have been put into forty two tribes and twelve subfamilies besides a group of uncertain affinities, Incertae Sedis. Apart from a high degree of taxonomic diversity, the family has unmatched ecological and economic importance. With its origin in the early cretaceous and major diversification in the mid cenozoic, the family at present covers nearly a fifth of land surface (Arabaci and Yildiz, 2004; Ture and Bell, 2004) and occurs in nearly all the habitats of the world (Clayton and Renvoize, 1986; Ture and Bocuk, 2007). All the cereals and millets are cultivated grasses. Sugarcane, the main source of sugar around the world is also a cultivated grass species. Besides, grasses constitute the main source of forage and fodder for livestock. Apart from food and fodder several grasses are used to extract of aromatic oils

* Corresponding Author: assoodan@gmail.com

and scents (Kaul and Vats, 1998; Khanuja *et al.*, 2005; Kim *et al.*, 2005; Bhuiyan *et al.*, 2008; Sujatha, 2010). Grasses also comprise the main source of green cover of our lawns and landscape for tourism and sports. Also, their use in handicraft and cottage industry is well known.

Grasses assume even a greater importance in areas like Punjab which has a tropical/subtropical ecology and an agrarian economy. But grasses have not been given sufficient attention in the floristic compilations of the region. Some of the earliest works have made no mention of the group (Bamber, 1916; Kashyap, 1936). Others have given only an alphabetic list of species with sketchy descriptions (Stewart, 1869; Sabnis, 1940; Nair, 1978; Sharma and Bir, 1978; Sharma, 1990). It is only in the work of Sharma and Khosla (1989) that grass species have been classified into sub families and tribes. With the establishment of Grass Phylogeny Working Group and their publication of a revised and phylogenetic classification (GPWG, 2001), we have witnessed a great upsurge in research on various aspects of grasses across the globe. Amidst a scenario of renewed interest in various aspect of grasses, India has been identified as a 'seriously under-collected country' as far as inventorization of grass diversity is considered (Kellogg, 2006).

We have initiated work on a systematic inventorization, description and classification of the grass species of Punjab and adjoining hills according to the latest format and classification system proposed by GPWG (2001). In the paper, we present a comparative and updated account of the reports of grasses from the region.

Materials and Methods

Area of study

The entire state of Punjab is located between 29° 30' to 30° 0' N and 73° 55' to 76° 50' E along with the adjoining hills were surveyed and explored for species diversity of grasses in the entire range of habitats occupied by them. According to the classification of forest types proposed by Champion and Seth (1968), the vegetation type of Punjab falls under the subgroup 5B Northern Tropical Dry Deciduous Forests. The average elevation in the plain region is 200-300 AMSL and the annual rainfall ranges from 250mm to 1000 mm at different locations in the regions. In the hilly areas, there are small stretches of Subtropical Pine Forest and Himalayan Moist Forest in which the ground is covered by the stoloniferous grasses. During the extensive exploration and collection surveys, notes were taken on the economic and ethnobotanical significance, distribution range and the flowering and fruiting phenology of the species.

Taxonomic Description

Taxonomic description of the species was done in sufficient details with special emphasis on diagnostic characters for identification and classification. Grasses have a special morphology. They have a fibrous root system and the stem has a horizontal portion below (rhizome) or above (stolon) the soil surface. The erect portion, the culm bears characteristic leaves with a sheath and a blade. The inflorescence is a panicle, a raceme or a spike modified in various ways. The basic unit of the inflorescence is the spikelet which and typically, consists of two glumes enclosing one or more florets. Each bisexual floret consists of a lemma, a palea, 2-3 lodicules, 3 stamens and a pistil with a bifid plumose stigma. Besides, the spikelet may bear male and reduced florets. The description was based on characters of morphology and micromorphology of the vegetative (culm, leaf, blade, ligule, collar etc.) and reproductive (inflorescence, spikelet, floret, caryopsis etc.) parts. Stereoscopic study was followed by drawings which were inked to prepare the plates which were computer scanned

to prepare the final prints.

Identification

Identification of species was done with the help of the floras and compilations mentioned above as well as those of Clayton and Renvoize (1986) and Cope (1982). Specimens were deposited in the Departmental Herbarium.

Results and Discussion

Table 1 presents a comparative statement of grass species reported in some of the important floristic compilations of the region. Irrespective of the schemes of classification followed by these authors, we have annotated the species into the latest system of the GPWG (2001). Compared to just 5-6 subfamilies in the earlier systems of classification (Prat, 1960; Caro, 1982; Clayton and Renvoize, 1986; Watson and Dallwitz, 1992) the GPWG system recognizes 12 subfamilies with the thirteenth group 'Incertae Sedis' of uncertain affinities. Furthermore, the subfamilies have been classified into 42 tribes. Therefore, reclassification into the GPWG system has involved a revision in subfamilial and tribal affiliation of several species listed in Table 1.

A close scrutiny of various columns of Table 1 clearly shows that grasses received scant attention in the earliest floristic compilations and that work on their exploration and inventorization has made a staggered progress.

Stewart (1869) listed only 40 grass species, whereas Kashyap (1936) did not include grasses in his work. Sabnis (1940) prepared an alphabetic list of 74 species taking the cumulative total to 93 which further increased to 133 with Nair (1978) and 147 with Sharma and Bir (1978). However, the compendium by Sharma and Khosla (1989) registered addition of thirty eight new species taking the cumulative total to 185.

However, since the last revision of the grass flora of the region, grass systematics has picked momentum across the globe mainly after the establishment of the Grass Phylogeny Working Group and their system of classification (GPWG, 2001). This has aroused interest and provided invaluable guidance in systematics and evolution of grasses of the world. Several researchers have studied this interesting and promising group of plants. These studies relate to the fundamental questions relating to grass organography and development (Kellogg, 2000), morphological nature of the spikelets and its parts (Ambrose *et al.*, 2000) and origin of C₄ anatomy (Gaut and Doebley, 1997; Kellogg, 2000). Systematic analysis of grasses has also picked momentum (Linder and Rudall, 2005; Soreng *et al.*, 2007). Within subfamilies, large and complex genera have been revised (Salamini *et al.*, 2002; Saarela *et al.*, 2003; Spangler, 2003; Molina and De Aerasar, 2004; Finot *et al.*, 2005; Zuloaga and Morrone, 2005). Owing to their cosmopolitan distribution and occurrence in varied habitats and sharply defined phases in vegetative and reproductive phenology, grasses are being considered as ideal indicator of climate change (Yuan *et al.*, 2007).

Work on Indian grasses has resulted in new species reports (Gopalan and Chandrasekran, 2001; Veldkamp and Salunkhe, 2000; Kiranraj, 2008; Takhar and Katewa, 2008; Sur, 2001; Ravi *et al.*, 2001; Kumar *et al.*, 2008) documentation of intraspecific diversity through molecular and other methods (Chandra *et al.*, 2004; Saxena and Chandra, 2006) and ethanobotanical uses (Sahu *et al.*, 2010). Ecological studies relate to species association (Soodan *et al.*, 2009) habitat preference and dispersion (Bazzaz, 1991; Sharma *et al.*, 2010).

Even though research in grasses around the world and within the country has extended to several aspects, basic exploration and inventorization is far from complete.

Our studies brings the cumulative number of species to 192 with 7 new species reports for the region *Briza minor*, *Cymbopogon citratus*, *Elytrophorus spicatus*, *Polygonatherum crinitum*, *Themeda anathera*, *Tripogon jacquemontii*, *Zoysia matrella* (Table 1).

Among the subfamilies, Panicoideae is the best represented in the region. Within this subfamily 4 tribes with 98 species is the most well represented. It is expectedly so since Panicoid grasses are known to dominate the warmer regions like the area of present investigations. The species representation of subfamilies is: Bambusoideae (4), Ehrartoideae (3), Centothecoideae (1), Pooideae (21), Aristidoideae (5), Arundinoideae (5) and Chloridoideae (55).

Table 1. A comparative list of grass species of Punjab

Numbers (1, 2, 3 etc.) refer to authors named in the head row who reported under these synonyms, (+) = Reported; (-) = Not reported; (*) = First reports; [-] = Species not recovered.

TAXON	1 Stewart (1869)	2 Sabnis (1940)	3 Nair (1978)	4 Sharma and Bir (1978)	5 Sharma and Khosla (1989)	6 Present Work
Subfamily: Bambusoideae						
Tribe: Bambuseae						
<i>Bambusa glaucescens</i> (Willd.) Siebold ex Munro (1868) 89 [Syn. <i>B. nana</i> Roxb. (1832) 199 (2)] [#]	-	+*	-	+	-	+
<i>B. nutans</i> G.C.Wall. ex Munro (1868) 92	-	-	-	+*	+	+
<i>B. vulgaris</i> Schrad. ex J.C. Wendl. (1808) 26 [Syn. <i>B. arundinacea</i> (Retz.) Willd. (1799) 245 (1)]	+	-	-	+	+	+
<i>Dendrocalamus strictus</i> (Roxb.) Nees (1834) 476-477	+	-	-	-	+	+
<i>Drepanostachyum falcatum</i> (Nees) Keng f. (1983) 16 [Syn. <i>Arundinaria falcata</i> Nees (1834) 478 (1)]	+	-	-	-	-	+
Subfamily: Ehrhartoideae						
Tribe: Oryzeae						
<i>Leersia hexandra</i> Sw. (1788) 21	-	-	-	-	+*	[-]
<i>Oryza rufipogon</i> Griff. (1851) 5, t. 144	-	-	-	+*	+	+
<i>O. sativa</i> L. (1753) 353	+	+	+	+	+	+
Subfamily: Centothecoideae						
Tribe: Thysanolaenae						
<i>Thysanolaena maxima</i> (Roxb.) Kuntze (1891) 794	-	-	-	+*	+	+
Subfamily: Pooideae						
Tribe: Triticeae						
<i>Secale cereale</i> L. (1753) 84	+	-	-	-	-	+
<i>Hordeum aegiceras</i> Nees ex Royle (1839-1840) 418	+	-	-	-	-	+
<i>H. vulgare</i> L. (1753) 84-85 [Syn. <i>H. coeleste</i> (L.) P. Beauv. (1812) 114 (1)]	+	-	+	+	+	+
[<i>H. hexastichum</i> L. (1753) 85 (1)]						
<i>Triticum aestivum</i> L. (1753) 85	+	+	+	+	+	+
[Syn. <i>T. vulgare</i> Vill. (1787) 153 (2)]						
<i>T. compactum</i> Host (1809) 4,7	-	-	-	+*	-	+
<i>T. durum</i> Desf. (1798) 114	+	-	-	+	-	+
<i>T. sphaerococcum</i> Percival (1921) 157, 321 f. 202	-	-	-	+*	-	[-]
Tribe: Bromeae						
<i>Bromus tectorum</i> L. (1753) 77	-	+*	-	-	-	+
Tribe: Poeae						
<i>Alopecurus nepalensis</i> Trin. ex Steud. (1854) 148	-	+*	+	-	+	+
<i>Avena fatua</i> L. (1753) 80	+	+	+	+	+	+
<i>A. sativa</i> L. (1753) 79	+	-	+	-	-	+
<i>A. sterilis</i> L. (1762) 118	-	-	+*	-	+	+
<i>Agrostis gigantea</i> Roth (1788) 31 [<i>A. alba</i> L. (1753) 63 (2)]	-	+*	-	-	-	+

TAXON	1 Stewart (1869)	2 Sabnis (1940)	3 Nair (1978)	4 Sharma and Bir (1978)	5 Sharma and Khosla (1989)	6 Present Work
<i>Briza minor</i> L. (1753) 70	-	-	-	-	-	+*
<i>Lolium temulentum</i> L. (1753) 83	-	+*	+	+	+	+
<i>Phalaris minor</i> Retz. (1783) 8	-	+*	+	-	-	+
<i>Poa annua</i> L. (1753) 68	-	+*	-	+	+	+
<i>Polypogon fugax</i> Nees ex Steud. (1854) 184	-	-	-	-	+*	+
<i>P. monspeliensis</i> (L.) Desf. (1798) 67	-	+*	+	+	+	+
<i>Rostraria cristata</i> (L.) Tzvelev (1970, 1971) 47	-	-	+*	+	+	+
[Syn. <i>Lophochloa phleoides</i> (Vill.) Rchb. (1830) 42 (3)]	-	-	+*	+	+	+
<i>R. pumila</i> (Desf.) Tzvelev (1970) 48	-	-	+*	-	+	+
[Syn. <i>Lophochloa pumila</i> (Desf.) Bor (1960) 445 (3)]	-	-	+*	-	+	+
Tribe: Meliceae						
<i>Melica</i> sp.	+	-	-	-	-	[-]
Tribe: Stipeae						
<i>Stipa orientalis</i> Trin. (1829) 83	-	+*	-	-	-	[-]
Tribe : Nardeae						
<i>Nardus stricta</i> L. (1753) 53	+	-	-	-	-	[-]
Subfamily: Arundinoideae						
Tribe: Arundineae						
<i>Arundo donax</i> L. (1753) 81	+	+	-	+	+	+
<i>Elytrophorus spicatus</i> Willd A. Camus (1923) 547	-	-	-	-	-	+*
<i>Phragmites australis</i> (Cav.) Trin. ex Steud. (1840) 143	-	-	-	-	-	
[Syn. <i>Arundo phragmites</i> L. (1753) 81 (1)]	+	+	-	-	+	+
[<i>P. communis</i> Trin. (1820,1822) 134 (2)]	-	-	-	-	-	
<i>P. karka</i> (Retz.) Trin. ex Steud. (1841) 324	-	+*	+	+	+	+
Sub family: Danthonoideae						
Tribe: Danthoniae						
<i>Schismus arabicus</i> Nees (1841) 422	-	-	+*	-	-	[-]
Subfamily : Aristidoideae						
Tribe : Aristideae						
<i>Aristida adscensionis</i> L. (1753) 82	+	+	+	+	+	+
[Syn. <i>A. depressa</i> Retz. (1786) 22 (1)]	-	-	-	-	-	
<i>A. funiculata</i> Trin. et Rupr. (1842) 159	-	+*	+	-	+	+
<i>A. hystericula</i> Edgew. (1862) 208	-	+*	+	-	-	+
<i>A. hystricula</i> L. f. (1781,1782) 113	-	+*	+	-	-	+
<i>A. mutabilis</i> Trin. & Rupr. (1842)150-151	-	-	+*	-	-	+
<i>Stipagrostis hirtigluma</i> (Steud. ex Rupr. & Trin.) De Winter (1963) 134, 136	-	-	-	-	-	
[Syn. <i>Aristida hirtigluma</i> Steud. ex Trin. & Rupr.(1842) 171-172 (2)]	-	+*	-	-	-	+
Subfamily: Chloridoideae						
Tribe: Pappophoreae						
<i>Enneapogon cenchroides</i> (Licht. ex Roem. et Schult.) CE. Hubb (1934) 119	-	-	+*	-	-	+
<i>E. desvauxii</i> P. Beauv. (1812) 82, t. 16, f. 11	-	-	-	-	-	
[Syn. <i>Pappophorum brachystachyum</i> Jaub. & Spach.(1850) 365 (3)]	-	-	+*	-	-	+
<i>E. persicus</i> Boiss. (1844) 71	-	-	-	-	-	
[Syn. <i>Pappophorum aucheri</i> Jaub. & Spach. (1851)32,323 (2)]	-	+*	+	-	-	+
Tribe: Cynodontae						
<i>Aeluropus lagopoides</i> (L.) Trin. ex Thwaites (1864) 374	-	-	-	-	-	
[Syn. <i>A. repens</i> Trin. (1848) 462 (1)]	+	+	-	-	-	+
[<i>A. villosus</i> Trin. ex C.A. (1896) 334 (2)]	-	-	-	-	-	
<i>Chloris barbata</i> Sw. (1797) 200	-	-	-	-	+*	+
<i>C. dolichostachya</i> Lag. (1816)	-	-	+*	-	+	+
<i>C. gayana</i> Kunth (1830) 293, 58	-	-	-	-	+*	+
<i>C. montana</i> Roxb. (1820) 331	-	-	+*	-	+	+
<i>C. virgata</i> Sw. (1797) 203	-	-	+*	+	+	+
<i>Crypsis schoenoides</i> L. (1791) 166, 42, f. 1	-	-	+*	-	-	-

TAXON	1 Stewart (1869)	2 Sabnis (1940)	3 Nair (1978)	4 Sharma and Bir (1978)	5 Sharma and Khosla (1989)	6 Present Work
<i>Cynodon barberi</i> Rang. & Tadul. (1916) 846	-	-	-	-	+*	+
<i>C. dactylon</i> (L.) Pers. (1805) 85	+	+	+	+	+	+
<i>Melanocenchrus abyssinica</i> (R. Br. ex Fresen.) Hochst. (1855) 274	-	-	+	-	+	+
<i>M. jacquemontii</i> Jaub. et Spach. (1851) 36, t.325	-	+*	+	-	-	+
[Syn. <i>Gracilea royleana</i> Hook. f. (1896,1897) 284 (2)]						
<i>Ochthochloa compressa</i> (Forssk.) Hilu (1981) 560						
[<i>Eleusine flagellifera</i> Nees (1842) 220 (2)]						
[Syn. <i>E. compressa</i> (Forssk.) Asch. & Schweinf. ex C. Chr. (1922) 12 (3)]						
<i>Oropetium biflorus</i> Stapf (1820) 98	-	+*	-	-	-	[-]
<i>O. thomaeum</i> (L. f.) Trin. (1820) 98 pl. 3	-	+*	+	-	+	+
<i>Perotis hordeiformis</i> Nees (1838, 1841) 247- 248	-	-	-	+*	+	+
<i>P. indica</i> (L.) Kuntze (1891) 787	-	-	+*	-	-	+
<i>Tetrapogon tenellus</i> (J. König ex Roxb.) Chiov. (1908) 352	-	+*	+	-	-	+
[Syn. <i>Chloris tenella</i> J. König ex Roxb. (1820) 330 (2)]						
<i>T. villosus</i> Desf. (1799) 389, pl. 255	-	+*	-	-	-	+
[Syn. <i>Chloris villosa</i> (Desf.) Pers. (1805) 87 (2)]						
<i>Tragus racemosus</i> (L.) All. (1785) 241						
[Syn. <i>T. roxburghii</i> Panigrahi (1974) 496 (3)]	-	+*	+	+	+	+
[<i>T. biflorus</i> Schult. (1824) 205 (4)]						
<i>Tripogon jacquemontii</i> Nees ex Steud. (1892) 85	-	-	-	-	-	+*
<i>Zoysia matrella</i> (L.) Merr. (1912) 20, 230	-	-	-	-	-	+*
Tribe: Eragrostideae						
<i>Acraчne racemosa</i> (Heyne. ex Roem. & Schult.) Ohwi (1947) 1	-	+*	-	-	+	+
[Syn. <i>Eleusine verticillata</i> (Roxb.) (1820) 346 (2)]						
<i>Cleistogenes gatacrei</i> (Stapf) Bor (1960) 487	-	-	-	-	+*	+
<i>Dactyloctenium aegyptium</i> (L.) Willd. (1809) 1029	+	+	+	+	+	+
[Syn. <i>Eleusine aegyptia</i> (L.) Desf. (1798) 85 (2)]						
<i>D. aristatum</i> Link (1827) 59	-	-	-	-	+*	+
<i>D. sindicum</i> Boiss. (1859) 131	-	-	+*	-	+	+
<i>Desmostachya bipinnata</i> (L.) Stapf (1900) 632						
[Syn. <i>Erogrostis cynosuroides</i> (Retz.) P. Beauv. (1812) 71, 162, 174 (1)]	+	-	+	+	+	+
<i>Dinebra retroflexa</i> (Vahl) Panz. (1813) 59-60	-	-	-	+*	+	+
<i>Eleusine coracana</i> (L.) Gaertn. (1788) 8	+	-	-	+	+	+
<i>E. indica</i> (L.) Gaertn. (1788) 8	-	-	-	-	+*	+
<i>Eragrostiella nardoides</i> (Trin.) Bor (1940) 270	-	-	-	-	+*	+
<i>Eragrostis atrovirens</i> (Desf.) Trin. ex Steud. (1840) 562	-	-	-	+*	+	[-]
<i>E. cilianensis</i> (Bellardi) Vignolo ex Janch. (1907) 110	-	-	+*	-	+	[-]
[Syn. <i>E. major</i> L. (1809) 14,24 (2)]	-	-	+*	-	+	[-]
<i>E. ciliaris</i> (L.) R.Br. (1818) 478	-	-	+	+	+	+
<i>E. coarctata</i> Stapf 313 (1897).	-	-	+*	-	-	+
<i>E. diarrhena</i> (Schult. & Schult f.) Steud. (1854) 266	-	-	+*	+	+	+
<i>E. diplachnoides</i> Steud. (1854) 268	-	-	-	-	+*	+
<i>E. gangetica</i> (Roxb.) Steud. (1854) 266	-	-	+*	-	+	+
<i>E. japonica</i> (Thunb.) Trin. (1830) 405	-	-	+*	-	+	+
[Syn. <i>E. interrupta</i> (Thunb.) Trin. (1812) 71, 162,175 (2)]	-	+*	+	+	+	+
<i>E. minor</i> Host (1809) 15						
[Syn. <i>E. poaeoides</i> P. Beauv. (1812) 162 (3)]	-	+*	+	+	+	+
<i>E. nutans</i> (Retz.) Nees ex Steud. (1840) 563	-	-	+*	-	-	+
<i>E. pilosa</i> (L.) P. Beauv. (1812) 71, 162, 175	-	+*	+	+	+	+
<i>E. tenella</i> (L.) P. Beauv. Roem. & Schult. (1817) 576	-	-	+*	+	+	+
<i>E. tremula</i> Hochst. ex Steud. (1854) 269	-	+*	+	+	+	+
<i>Leptochloa chinensis</i> (L.) Nees (1824) 4	-	-	-	-	+*	+
<i>L. paniceae</i> (Retz.) Ohwi (1941) 311	-	+*	+	+	+	+
[Syn. <i>L. filiformis</i> (Pers.) P Beauv. (1812) 163,166 (2)]						
<i>Neyraudia arundinacea</i> (L.) Henrard (1929) 8	-	-	-	-	+*	+

TAXON	1 Stewart (1869)	2 Sabnis (1940)	3 Nair (1978)	4 Sharma and Bir (1978)	5 Sharma and Khosla (1989)	6 Present Work
<i>Sporobolus coromandelianus</i> (Retz.) Kunth ((1829) 68	-	+*	-	+	+	+
<i>S. diandrus</i> (Retz.) P. Beauv. (1812) 26, 147, 178	-	+*	+	+	+	+
<i>S. fertilis</i> Steud. (1965) 291	-	+*	-	-	+	+
[Syn. <i>S. indicus</i> (L.) R. Br. (1810) 170 (2)]	-	+*	-	-	+	+
<i>S. helvolus</i> (Trin.) T. Durand. & Schinz (1895) 820	-	+*	+	-	+	+
[Syn. <i>S. glaucifolius</i> (Hochst. ex Steud.) Hochst. ex T. Durand & Schinz (1854) 154 (2)]	-	+*	+	-	+	+
<i>S. ioclados</i> (Nees ex Trin.) Nees (1841) 161	-	-	+*	+	+	+
[Syn. <i>S. marginatus</i> Hochst. ex A. Rich. (1850) 397 (3)]	-	-	+*	+	+	+
<i>S. tenuissimum</i> (Mart. ex Schrank) Kuntze (1898) 369	-	+	+	-	-	+
[Syn. <i>S. minutiflorus</i> (Trin.) Link (1827) 88 (2)]	-	+	+	-	-	+
Subfamily : Panicoideae						
Tribe: Isachneae						
<i>Isachne albens</i> Trin. (1828) 8, 85	-	-	+*	-	-	[-]
<i>I. himalaica</i> Hook. f. 1897(1896) 23	-	+*	-	-	-	+
[Syn. <i>I. australis</i> R. Br. (1810) 196 (2)]	-	+*	-	-	-	+
Tribe: Paniceae						
<i>Alloteropsis cimicina</i> (L.) Stapf (1919) 487	-	-	-	-	+*	+
<i>Brachiaria brizantha</i> (Hochst. ex A. Rich.) Stapf (1919) 531	-	-	-	-	+*	+
<i>B. distachya</i> (L.) Stapf (1919) 565	-	-	-	-	+*	+
<i>B. mutica</i> (Forssk.) Stapf (1919) 526	-	-	-	-	+*	+
<i>B. ramosa</i> (L.) Stapf (1919) 542-544	-	+*	+	+	+*	+
[Syn. <i>Panicum ramosum</i> L. (1767) 29-30 (2)]	-	+*	+	+	+*	+
<i>B. reptans</i> (L.) C. Gardner & C.E. Hubb. (1938) pl 3363, f. 3	-	+*	+	-	+	+
[Syn. <i>Panicum prostratum</i> Lam. (1791) 171 (2)]	-	+*	+	-	+	+
<i>Cenchrus biflorus</i> Roxb. (1820) 238	-	+*	+	+	+	+
[Syn. <i>C. catharticus</i> Delile. (1839) 4 (2)]	-	+*	+	+	+	+
<i>C. ciliaris</i> L. (1771) 302	+	+	+	+	+	+
[Syn. <i>Pennisetum cenchroides</i> Rich. ex pers. (1805) 72 (1)]	+	+	+	+	+	+
<i>C. echinatus</i> L. (1753) 1050	+	-	-	-	-	+
<i>C. pennisetiformis</i> (Hosch.) & Steud.) (1854) 109	-	-	-	+*	+	+
<i>C. prieurii</i> (Kunth) Maire (1931) 523	-	-	+*	-	-	+
<i>C. setigerus</i> Vahl (1805) 395	-	-	+*	+	+	+
<i>Digitaria abludens</i> (Roem. & Schult.) Veldkamp (1973) 53-55	-	-	-	-	+*	+
<i>D. bicornis</i> (Lam.) Roem. & Schult. (1817) 470	-	-	-	-	+*	+
<i>D. ciliaris</i> (Retz.) Koeler (1802) 27	-	-	-	-	-	-
[Syn. <i>D. bifloris</i> Willd. (1809) 92 (3)]	-	-	+*	+	+	+
[Syn. <i>D. adscendens</i> (Kunth) Henrard (1934) 92 (4)]	-	-	-	-	-	-
<i>D. longiflora</i> (Retz.) Pers. (1805) 85	-	-	-	-	+*	+
<i>D. nodosa</i> Parl. (1842) 39	-	-	+*	-	-	+
<i>D. radicosa</i> (J. Presl.) Miq. (1857) 437	-	-	-	-	+	+
<i>D. sanguinalis</i> Pers. (1771) 52	+	-	-	-	-	+
<i>D. setigera</i> Roth (1817) 474	-	-	-	-	+*	+
<i>D. stricta</i> Roth ex Roem. & Schult. (1817) 474	-	-	+*	+	+	+
<i>Echinochloa colonum</i> (L.) Link. ((1833) 209	-	+*	+	+	+	+
[Syn. <i>Panicum colonum</i> L. (1759) 870 (2)]	-	+*	+	+	+	+
<i>E. crusgalli</i> (L.) P. Beauv. (1812) 161	-	+*	+	-	+	+
[Syn. <i>Panicum crusgalli</i> L. (1753) 56 (2)]	-	+*	+	-	+	+
<i>E. stagnina</i> (Retz.) P. Beauv. (1812) 171	-	-	-	-	+*	+
<i>Eriochloa fatmensis</i> (Hosch.) W. D. Clayton (1975) 108	-	-	-	-	-	-
[Syn. <i>E. nubica</i> (Steud.) Hack. & Stapf ex Thell. (1919) 697 (4)]	-	-	-	+*	+	+
<i>E. procera</i> (Retz.) C.E. Hubb. (1930) 256	-	-	+*	-	-	+
<i>Oplismenus burmanii</i> (Retz.) P. Beauv. (1812) 168,169	-	-	-	+*	+	+
<i>O. compositus</i> (L.) P. Beauv. (1812) 168,169	-	-	+*	+	+	+
<i>Panicum antidotale</i> Retz. (1786) 17	+	+	+	+	+	+
[Syn. <i>P. miliare</i> Tam. (1791) 173 (1)]	+	+	+	+	+	+
<i>P. atrosanguineum</i> Hochst. ex A. Rich. (1851) 375	+	+	+	-	-	+
[Syn. <i>P. hydaspicum</i> Edgew. (1862) 207 (1)]	+	+	+	-	-	+

TAXON	1 Stewart (1869)	2 Sabnis (1940)	3 Nair (1978)	4 Sharma and Bir (1978)	5 Sharma and Khosla (1989)	6 Present Work
<i>P. maximum</i> Jacq. (1781) 2, 13	-	-	+*	+	+	+
<i>P. miliaceum</i> L. (1753) 58	+	-	+	-	+	+
<i>P. paludosum</i> Roxb. (1820) 310	-	-	-	-	+	+
<i>P. virgatum</i> Roxb. ex Steud. (1841) 262	-	-	+*	-	+	+
[Syn. <i>P. psilopodium</i> Trin. (1826) 217 (5)]	-	-	+*	-	+	+
<i>P. repens</i> L. (1762) 86	-	-	-	-	+*	+
<i>P. tenellum</i> Roxb. (1854) 21	-	-	+*	-	+	+
[Syn. <i>P. trypheron</i> Schult. (1824) 244 (3)]	-	-	+*	-	+	+
<i>Paspalidium flavidum</i> (Retz.) A. Camus (1922) 419	-	-	+*	+	+	+
<i>Paspalum dilatatum</i> Poir. (1804) 35	-	-	-	-	+*	+
<i>P. longifolium</i> Roxb. (1820) 283	-	+*	+	-	-	+
[Syn. <i>P. longiflorum</i> Retz. (1786) 15 (2)]	-	+*	+	-	-	+
<i>P. notatum</i> Fluegge. (1810) 106	-	-	-	-	+*	+
<i>P. scrobiculatum</i> L. (1767) 29	+	-	+	-	+	+
<i>P. vaginatum</i> Sw. (1788) 21	-	+*	+	-	-	+
[Syn. <i>P. sanguinale</i> L. (1896) 15 (2)]	-	+*	+	+	+	+
[<i>P. paspaloides</i> (Michx.) Scribn. (1894) 29 (4)]	-	+*	+	+	+	+
<i>Pennisetum glaucum</i> (L.) R. Br. (1810) 195	-	-	-	-	-	-
[Syn. <i>Penicillaria spicata</i> Willd. (1809) 1037 (1)]	+	+	+	+	+	+
[<i>P. typhoides</i> (Burm.) Stapf et C.E. Hubb. (1933) 271 (2)]	-	+*	+	-	-	-
<i>P. orientale</i> (L.) C. Rich. (1805) 72	-	+*	+	-	+	+
<i>P. polystachyon</i> (L.) Schult. (1824) 146	-	-	-	-	+*	+
<i>P. purpureum</i> Schumach. (1827) 44	-	-	-	+*	+	+
<i>Setaria intermedia</i> Roem. & Schultz. (1817) 489	-	-	-	+*	+	+
[Syn. <i>S. tomentosa</i> (Roxb.) Kunth (1829) 47 (4)]	-	-	-	+*	+	+
<i>S. italicica</i> (L.) P. Beauv. (1812) 170, 178	-	-	-	-	+	+
[Syn. <i>Panicum italicum</i> R.Br. (1753) 56 (2)]	-	-	-	-	+	+
<i>S. pumila</i> (Poir.) Roem. & Schultz. (1817) 891	-	+*	+	-	+	+
[Syn. <i>S. glauca</i> (L.) P. Beauv (1812) 168, 169.(3)]	-	-	-	-	+	+
<i>S. sphacelata</i> (Schumach.) Stapf et C.E. Hubb. (1929) 195	-	-	-	-	+	+
<i>S. verticillata</i> (L.) P. Beauv. (1812) 178	-	-	+*	+	+	+
[Syn. <i>Panicum verticillatum</i> L. (1762) (3)]	-	-	+*	+	+	+
<i>Urochloa panicoides</i> P. Beauv. (1812) 53	-	-	+*	+	+	+
Tribe: Arundinelleae						
<i>Arundinella nepalensis</i> Trin. (1826) 62	-	-	-	-	+*	+
Tribe: Andropogoneae						
<i>Andropogon glomeratus</i> (Walter) Britton (1888) 67	+	-	-	-	-	+
[Syn. <i>Chrysopogon glaucoptis</i> Stend (1934) 139 (1)]	-	-	-	-	-	-
<i>A. pumilus</i> Roxb. (1894) 496	-	-	+*	-	-	+
<i>Apluda mutica</i> L. (1753) 82	-	-	-	-	-	-
[Syn. <i>A. varia</i> Hack. Subspecies <i>aristata</i> Hack. (1889 (199) (2)]	-	-	+*	-	+	+
[<i>A. aristata</i> L. (1756) 303 (5)]	-	-	-	-	-	-
<i>Arthraxon lancifolius</i> (Trin.) Hochst. (1856) 188	-	+*	-	-	+	+
[Syn. <i>Andropogon monticola</i> Schult. (1827) 665 (2)]	-	-	-	-	+*	+
<i>A. prionoides</i> Steud. (1956) 399	-	-	-	-	+*	+
<i>Bothriochloa insculpta</i> (Hochst.) A. Camus (1931) 165	-	-	-	-	+*	+
<i>B. bladhii</i> (Retz.) S.T. Blake. (1969) 62	-	-	-	-	+*	+
[<i>B. odorata</i> (Lisboa) A. Camus (1931) 165 (5)]	-	-	-	-	+*	+
<i>B. ischaemum</i> L. (1936) 201	-	-	+*	-	-	+
<i>B. pertusa</i> (L.) A. Camus (1931) 164	-	-	+*	+	+	+
<i>Capillipedium huegelii</i> (Hack.) A. Camus (1921) 308	-	-	-	-	+*	+
<i>C. parviflorum</i> (R.Br.) Stapf (1917) 169	-	-	-	-	+*	+
<i>Chrysopogon serrulatus</i> Trin. (1832) 318	-	-	-	-	-	-
[Syn. <i>Andropogon monticola</i> Schult. (1896) 193 (2)]	-	+*	+	-	+	+
[<i>C. fulvus</i> (Spreng.) Chiov. (1929) 327 (3)]	-	-	-	-	-	-
<i>Coix lacryma-jobi</i> L. ((1753) 972	-	-	-	-	+*	+
<i>Cymbopogon commutatus</i> (Steud.) Stapf (1907) 211	-	-	+*	-	+	+
[Syn. <i>C. parkeri</i> Stapf. (1929) 10 (3)]	-	-	-	-	-	-

TAXON	1 Stewart (1869)	2 Sabnis (1940)	3 Nair (1978)	4 Sharma and Bir (1978)	5 Sharma and Khosla (1989)	6 Present Work
<i>C. citratus</i> (DC.) Stapf. (1906) 357	-	-	-	-	-	+*
<i>C. jwarancusa</i> (Jones) Schult. (1824) 458	+	+	+	-	+	+
[Syn. <i>Andropogon schoenanthus</i> (L.) Spreng. (1896) 204. (2)]						
<i>C. martinii</i> (Roxb.) Watson (1882) 392	-	-	+*	-	+	+
<i>C. nardus</i> (L.) Rendle (1899) 155	-	-	-	-	+*	+
<i>Dichanthium annulatum</i> (Forssk.) Stapf (1917) 178	+	+	+	+	+	+
[Syn. <i>Andropogon annulatus</i> Forssk. (1775) 173 (1)]						
<i>D. caricosum</i> L. A. Camus (1921) 459	-	-	-	-	+*	+
<i>D. foveolatum</i> (Delile) Roberty (1960) 170		+*	-	-	+	+
[Syn. <i>Andropogon foveolatus</i> Delile (1812) 16 t. 82. (2)]						
<i>Eulaliopsis binata</i> (Retz.) C. E. Hubb. (1935) 3262	-	+*	-	-	+	+
[Syn. <i>Ischaemum angustifolium</i> Hook. (1889) 241 (2)]						
<i>Hackelochloa granularis</i> (L.) O. kuntze. (1891) 776	-	-	-	-	+*	+
<i>Hemarthria compressa</i> (L. f.) R. Br. (1810) 207		+*	+	+	+	+
[Syn. <i>Rottboellia compressa</i> L. (1781) 114 (2)]						
<i>Heteropogon contortus</i> (L.) P. Beauv. ex Roem. & Schult. (1817) 836	+	+	+	+	+	+
[Syn. <i>Andropogon contortus</i> L. (1753) 1045 (2)]						
<i>Imperata cylindrica</i> (L.) P. Beauv. (1812) 165	-	+*	+	+	+	+
[Syn. <i>I. arundinacea</i> Cypri. (1792) 27, t. 11 (2)]						
<i>Ischaemum rugosum</i> Salisb. (1791) 1	-	-	-	-	+*	+
<i>Iseilema prostratum</i> (L.) Andersson (1856) 251						
[Syn. <i>I. wightii</i> Andersson (1856) 251 (2)]	-	+*	+	-	+	+
[<i>Andropogon prostratus</i> L. (1856) 251 (3)]						
<i>Lasiurus sindicus</i> (Boiss.) Henr. (1941) 414	-	+*	+	-	+	+
[Syn. <i>Elionurus hirsutus</i> auct. (1881) 68 (2)]						
<i>Pogonatherum crinitum</i> (Thunb.) Kunth. (1906) 178	-	-	-	-	-	+*
<i>P. Paniceum</i> (Lam.) Hack. (1906) 178						
[Syn. <i>P. saccharoideum</i> P. Beauv. (1812) 176, 177 (2)]	-	+*	+	-	+	+
<i>Rottboellia cochinchinensis</i> (Lour.) Clayton (1981) 817	-	-	-	-	+*	+
[<i>R. exaltata</i> L. f. (1779) 40 (5)]						
<i>Mnesithea laevis</i> (Retz.) Kunth (1829) 154	-	+*	-	-	-	[-]
[Syn. <i>Rottboellia perforata</i> Roxb. (2)]						
<i>Saccharum bengalense</i> Retz. (1789) 16						
[Syn. <i>S. sara</i> Roxb. (1820) 249 (1)]	+	+	+	+	+	+
[<i>S. arundinaceum</i> Hook. f. (1786 -1787) 14 (2)]						
<i>S. officinarum</i> L. (1753) 54	+	-	+	+	+	+
<i>S. ravennae</i> L. (1774) 88	-	+*	+	+	+	+
[Syn. <i>Erianthus ravennae</i> Beauv. (1812) 162, 177 (2)]						
<i>S. spontaneum</i> L. (1771) 183	+	+	+	+	+	+
<i>Sorghum bicolor</i> (L.) Moench (1794) 207						
[Syn. <i>S. vulgare</i> Pers. (1805) 101 (1)]	+	-	+	+	+	+
[<i>S. cernuum</i> (Ard.) Host (1809) 2 (3)]						
<i>S. halepense</i> (L.) Pers. (1805) 101	+	+	+	+	+	+
[<i>Andropogon halpensis</i> Brot. (1804) 89 (2)]						
<i>Themeda anathera</i> (Nees ex Steud.) Hack. (1889) 669	-	-	-	-	-	+*
<i>T. quadrivalvis</i> (L.) O. Kuntze. (1891) 794	-	-	-	+*	+	+
[Syn. <i>Anthistiria ciliata</i> L. (1781-1782) 113 (1)]						
<i>Vetiveria zizanioides</i> (L.) Nash (1903) 67						
[Syn. <i>Anatherum muricatum</i> Beauv. (1812) 150 (1)]	+	+	+	+	+	+
[<i>Andropogon squarrosus</i> of Hook. f. (1896) 186 (2)]						
<i>Zea mays</i> L. (1753) 971	+	-	+	+	+	+
Species Number (Cumulative)	40	74 (93)	105 (133)	73 (147)	146 (185)	187 (192)

Conclusion

The reclassification of grasses of the region proposed in the present paper shall provide new direction to future studies in the group. This is in consonance with a world wide renewal of interest in systematics and evolution of this group. Apart from their conventional uses, grasses have emerged as model plant species for events in plant development and as indicators of environmental changes. Intensive explorations need to be carried out to identify candidate grass species for these areas in the modern biology. The usefulness of intensive explorations is indicated by the fact that the seven new species have been identified in the present study.

References

- Ambrose, B. A., Lerner, D. R., Ciceri, P., Padilla, C. M., Yajznofsky, M. F. and Schmidt, R. J. (2000) Molecular and genetic analysis of the *silky1* gene reveal conservation in the floral organ specification between dicots and monocots. *Molecular Cell* 5 (3): 569-579.
- Arabaci, T. and Yildiz, B. (2004) A floristical study on Poaceae spp., growing naturally in Malatya province. *Turkey Journal of Botany* 28: 361-368.
- Bamber, C. J. (1916) Plants of the Punjab. Superintendent, Government Printing Agency, Lahore.
- Bazzaz, F. A. (1991) Habitat selection in plants. *The American Naturalist* 137: S116-S130.
- Bhuiyan, M. N. I., Chowdhary, J. U. and Begum, J. (2008) Essential oil in roots of *Vetiveria Zizanioides* (L.) Nash Ex small from Bangladesh. *Bangladesh Journal of Botany* 37(2): 213-215.
- Caro, J. A. (1982) Sinopsis taxonomica de las gramineas argentinas. *Dominguezia* 4: 1-51.
- Champion, H. G. and Seth, S. K. (1968) A revised survey of the forest types of India. Government of India Press, Nasik.
- Chandra, A., Saxena, R., Roy, A. K. and Pathak, P. S. (2004) Estimation of genetic variation in *Dichanthium annulatum* genotypes by the RAPD technique. *Tropical Grasslands* 38: 245-252.
- Clayton, W. D. and Renvoize, S. A. (1986) Genera Graminum. Her Majesty's Stationery Office, London.
- Cope, T. (1982) Flora of Pakistan. In: Poaceae (eds. Nasir, E. and Ali, S. I.) No. 143. National Herbarium, Agricultural Research Council, Islamabad.
- Finot, V. L., Peterson, P. M., Zuloaga, F. O., Soreng, R. J. and Matthei, O. (2005) A revision of *Trisetum* (Poaceae: Pooideae: Aveniae) in South America. *Annals of Missouri Botanical Garden* 92: 533-568.
- Gaut, B. S. and Doebley, J. F. (1997) DNA sequence evidence for the segmental allotetraploid origin of maize. *Proceeding of Natural Academy of Sciences, USA* 94: 6809-6814.
- Gopalan, R. and Chadrasekaran, V. (2001) *Isachne jayachandranii* (Gramineae) - a new grass from southern Western Ghats, India. *Kew Botanical Garden, Kew*.
- GPWG (Grass Phylogeny Working Group) (2001) Phylogeny and subfamilial classification of the grasses (Poaceae). *Annals of the Missouri Botanical Garden* 88: 373-457.
- Kashyap, S. R. (1936) Lahore district flora. The University of Panjab, Lahore.
- Kaul, V. K. and Vats, S. K. (1998) Essential oil composition of *Bothriochloa pertusa* and phyletic relationship in aromatic grasses. *Biochemical Systematics and Ecology* 26 (3): 347-356.
- Kellogg, E. A. (2000) Molecular and morphological evolution in the Andropogoneae. In: *Grasses: Systematics and Evolution* (eds. Jacobs, S. W. L. and Everett, J.) 149-158. Commonwealth Scientific and Industrial Research Organisation (CSIRO), Melbourne.

- Kellogg, E. A. (2006) Beyond taxonomy: prospects for understanding morphological diversity in the grasses (Poaceae). *Darwiniana* 44: 7-17.
- Khanuja, S. P. S., Shasany, A. K., Pawar, A., Lal, R. K., Darokar, M. P., Naqvi, A. A., Rajumar, S., Sundarsan, V., Lal, N. and Kumar, S. (2005) Essential oil constituent and RAPD markers to establish species relationship in *Cymbopogon* Spreng. (Poaceae). *Biochemical Systematics and Ecology* 33: 171-186.
- Kim, H. J., Chen, F., Wang, X., Chung, H. Y. and Jin, Z. (2005) Evaluation of antioxidant activity of Vetiver (*Vetiveria zizanioides* L.) oil and identification of its antioxidant constituents. *Journal of Agricultural and Food Chemistry* 53: 7691-7695.
- Kiranraj, M. S. (2008) A new species of *Dimeria* (Poaceae, Panicoideae: Andropogoneae) from Goa, India. *Novon* 18: 183-186.
- Kumar, M., Seqiera, S. and Ramesh, M. (2008) The position of Indian Bamboos in Bamboo Systematics. *Rheedia* 10: 33-48.
- Linder, H. P. and Rudall, P. J. (2005) Evolutionary history of Poales. *Annual Review Ecology. Evolution Systematics* 36:107-124
- Molina, A. and De Agrasar, Z. E. R. (2004) Taxonomic revision of the species of the genus *Chloris* (Poaceae; Chloridoideae) in South America. *Candollea* 59: 347-427.
- Nair, N. C. (1978) Flora of the Punjab plains. Records of the Botanical Survey of India. Indian Botanical Garden, Howrah.
- Prat, H. (1960) Vers une classification naturelle des graminees. *Bulletin of Society of Botany, France* 107: 32-79.
- Ravi, N. Mohanan, N., Shaju, T., Kiranraj, M. S. and Rajesh, R. (2001) Three more species of *Ischaemum* L. (Poaceae) from Kerala, India. *Botanical Bulletin of Academy of Singapore* 42: 223-230.
- Saarela, J. M., Peterson, P. M., Soreng, R. J. and Chapman, R. E. (2003) A taxonomic revision of eastern north America and eastern Asian disjunct genus *Brachyelytrum* (Poaceae): evidence from morphology, phytogeography and AFLPs. *Systematic Botany* 28: 674-692.
- Sabnis, T. S. (1940) A contribution to the flora of Punjab plains and the associated Hill regions. *Journal of Bombay Natural History Society* 42(1): 125-149.
- Sahu, S. C., Rout, N. C. and Dha, N. K. (2010) Ethnobotany of *Eulaliopsis binata* (Retz.) Hubbard-Poaceae, in Odissa, Eastern India. *Journal of Advances in Developmental Research* 1(2): 155-160.
- Salamin, N., Hodgkinson, T. R. and Savolainen, V. (2002) Building supertrees: an empirical assessment using the grass family (Poaceae). *Systematic Biology* 51(1): 136-150
- Saxena, R. and Chandra, A. (2006) RAPD and cytological analysis and histological changes caused by moisture stress in *Dichanthium annulatum* accessions. *Cytologia* 71:197-204.
- Sharma, M. (1990) Punjab plants check list. Bishen singh Mahendra pal Singh, Dehradun, India.
- Sharma, M. and Bir, S. S. (1978) Flora of Patiala. Punjab University, Patiala Publications , Patiala, India.
- Sharma, M. L. and Khosla, P. K. (1989) Grasses of Punjab and Chandigarh. Publication Bureau, Punjab University, Lahore.
- Sharma, S., Kaur, R. and Soodan, A. S. (2010) Morphology of diaspores of some range grasses of Punjab. *Journal of Range Management and Agroforestry* 31(1):52-58.
- Soodan, A. S., Kumar, R. and Singh, B. P. (2009) Species association in the vascular plant community of the tropical upper Bari Doab region of the Punjab. *Indian Journal of Ecology*

- 36(2): 111-117.
- Soreng, R. J., Davis, J. I. and Voionmaa, M. A. (2007) A phylogenetic analysis of Poaceae tribe Poeae based on morphological characters and sequence data from three chloroplast-encoded genes: Evidence for reticulation and a new classification for the tribe. *Kew Bulletin* 62: 425-454.
- Spangler, R. E. (2003) Taxonomy of *Sarga*, *Sorghum* and *Vacoparis* (Poaceae: andropogoneae). *Australian Systematic Botany* 16: 279-299.
- Stewart, J. L. (1869) Punjab plants. Government Press, Lahore. (Reprint 1977, Bishen Singh, Dehradun).
- Sujatha, S. (2010) Essential oil and its insecticidal activity of medicinal aromatic plant *Vetiveria zizanioides* (L.) against the Red Flour Beetle *Tribolium castaneum* (Herbst). *Asian Journal of Agricultural Sciences* 2(3): 84-88.
- Sur, P. R. (2001) Taxonomic review of the genus *Eulaliopsis* Honda (Poaceae) in India. *Journal of Economic and Taxonomic Botany* 25(2): 374-378.
- Takhar, H. K. and Katewa, S. S. (2008) Systematic and floristic study of some new records of grasses of Indian Desert Rajasthan. *The Indian Forester* 134(8): 1017-1036.
- Ture, C. and Bell, R. W. (2004) Plant distribution and its relationship to extractable boron in naturally occurring High Boron Soils Turkey. *Israel Journal of Plant Science* (52): 125-132.
- Ture, C. and Bocuk, H. (2007) An investigation of diversity, distribution and conservation on Poaceae growing naturally in Eskişehir Province (Central Anatolia-Turkey). *Pakistan Journal of Botany* 39(4): 1055-1070.
- Veldkamp, J. F. and Salunkhe, C. B. (2000) *Chrysopogon castaneus* (Poaceae- Andropogoneae), a new species from Maharashtra, India. *Rheedea* 10(1): 59-61.
- Watson, L. and Dallwitz, M. J. (1992 onwards) The families of flowering plants: descriptions, illustrations, identification and information retrieval. Retreived from <http://delta-intkey.com/angio>. On 5 May 2012.
- Watson, L. and Dallwitz, M. J. (1999) Grasses genera of the world: descriptions, illustrations, identification and information retrieval, including synonyms, morphology, anatomy, physiology, phytochemistry, cytology, classification, pathogens, world and local distribution and references. retreived from: <http://biodiversity unoedu/delta/> On: 18 August 1999.
- Yuan, W., Zhou, G., Wang, Z., Han, X. and Wang, X. (2007) Stimulating phenological characteristics of two dominant grass species in a semi arid steppe ecosystem. *Ecological Research* 22: 784-791.
- Zuloaga, F. O. and Morrone, O. (2005) Revisión de las especies de *Paspalum* para América del Sur Austral (Argentina, Bolivia, Sur del Brasil, Chile, Paraguay y Uruguay). *Monographs in Systematic Botany from the Missouri Botanical Garden* 102: 1-297.

گزارش به روز شده‌ای از گندمیان پنجاب (هندوستان)

آمارجیت سینگ سودان*

گروه گیاه‌شناسی و علوم محیطی، دانشگاه گورو ناناک، پنجاب، هندوستان

چکیده

تیره Poaceae چهارمین تیره بزرگ گیاهان گل‌دار است. این تیره در بردارنده ۸۰۰–۷۰۰ جنس و ۱۱۰۰۰–۱۳۰۰۰ گونه با پراکنش جهانی است. این خانواده، از اهمیت بوم‌شناسی و اقتصادی بی‌نظیری برخوردار است. پیدایش این تیره به دوره کرتاسه نخستین و گونه‌گونی عمدۀ آن به دوره سنوزوئیک میانی باز می‌گردد. در حال حاضر، این تیره نزدیک به یک پنجم سطح خشکی‌ها را پوشش می‌دهد و تقریباً در تمامی زیستگاه‌ها در سطح جهان وجود دارد. با وجود شرایط بوم‌شناسی نیمه گرمسیری و موقعیت اقتصادی منطقه پنجاب، گندمیان بخش قابل ملاحظه‌ای را در منطقه تشکیل می‌دهند. با وجود اهمیت قریب به اتفاق، مطالعات تاکسونومیک در مورد گندمیان در این منطقه از توجه کافی برخوردار نبوده است. تنها در مطالعات Sharma و Khosla (۱۹۸۹) است که گونه‌های گندمیان به زیرتیره‌ها و طایفه‌ها طبقه‌بندی شده‌اند. با این حال، پس از تأسیس کارگروه تبارشناسی گندمیان (GPWG) جهان تحولی نوین در زمینه طبقه‌بندی گندمیان داشته است. اما کشور هند تا آنجا که به تنوع گندمیان مربوط می‌شود هنوز به طور کامل مورد توجه قرار نگرفته است. تحقیقات ما برای اکتشاف و رده‌بندی فلور تیره گندمیان منطقه، تلاشی برای تقویت و به روزرسانی اطلاعات درباره تنوع آن در منطقه مورد مطالعه است. مطالعات حاضر، جمع‌بندی ۱۹۲ مجموعه گونه از جمله ۷ گزارش جدید را ارائه کرده است. نمونه‌های گونه‌ای از از زیرتیره‌ها عبارتند از: (۱) Chloridoideae، (۲) Centothecoideae، (۳) Erhartoideae، (۴) Panicoideae، (۵) Arundoideae، (۶) Aristidoideae، (۷) Bambooidae، (۸) Pooideae، (۹) و (۱۰) Assoidae.

واژه‌های کلیدی: هندوستان، تیره گندمیان، پنجاب، رده‌بندی، تاکسونومی، گرمسیری

* assoodan@gmail.com