Determination of Genetic Variation in Indian Sesame (Sesamum indicum) Genotypes for Agro-Morphological Traits

Spandana Bandila^{1*}, Anuradha Ghanta¹, Sivaraj Natarajan², Sivaramakrishnan Subramoniam¹

 1- Institute of Biotechnology (IBT), Acharya N.G. Ranga Agricultural University (ANGRAU), Hyderabad-500030, India
2- National Bureau of Plant Genetic Resources (NBPGR), Hyderabad, India

Received: 5 August 2011

Accepted: 23 November 2011

*Corresponding author: Email: spandanabandila@gmail.com

ABSTRACT

Sesame (*Sesamum indicum* L.) is one of the oldest oil crops and is widely cultivated in Asia and Africa. Characterization and quantification of genetic diversity has long been a major goal in evolutionary biology. Information on the genetic diversity within and among closely related crop varieties is essential for a rational use of genetic resources. The analysis of genetic variation within and among elite breeding materials is of fundamental interest to plant breeders. It contributes to monitoring of germplasm and can also be used to predict potential genetic gains .To determine the level of diversity in relation to geographical origins and morphological characteristics, a total of 60 accessions have been collected from different parts of the India were analyzed using statistical techniques for thirteen quantitative and four qualitative parameters using D^2 analysis. Number of seeds per capsule contributed highest towards the divergence. The distribution pattern of genotypes in different clusters indicated that genetic divergence was not related to geographical differentiation.

Keywords: D² analysis; Euclidean distance; genetic diversity; morphological traits; *Sesamum indicum* L.

INTRODUCTION

Sesame (Sesamum indicum L., Pedaliaceae) is the most commonly cultivated edible oil crop species out of over 30 species in the genus Sesamum (Nayar and Mehra, 1970; Kobayashi et al., 1990). The crop has been cultivated in various ecological regions of Vietnam and Cambodia hundreds for of years. Information on genetic diversity and relationships among populations is important for plant breeding programs as it helps to select the right genetic material to be used (Ganesh and Thangavelu, 1995).

Genetic diversity in crop species can be determined bv using the agromorphological as well as biochemical and molecular markers (Liu, 1997; Geleta et al., 2007, 2008). Studies on sesame genetic diversity and divergence have been mainly on agro-morphological based traits. Several of these agro-morphological trait based studies have found a high genetic diversity in sesame populations (Bisht et al. 1998; Arriel et al., 2007). There is an improving ample scope for the productivity of this important oil seed crop through varietal improvement and hybrid cultivar development.

MATERIALS AND METHODS

Plant material

Germplasm of Sesamum indicum L. collected and stored by National Bureau of Plant Genetic Resources (N.B.P.G.R) Rajendranagar, Hyderabad from eight different states of India, were selected for the present study. Details of the germplasm accessions are furnished in Table 1. The experiment was carried out during late Kharif 2008-09. The experimental materials were sown in simple Randomised Block Design with 60 ×10 cm spacing in three replications at College Farm, College of Agriculture, Acharya N.G. Ranga Agricultural University, Rajendranagar, Hyderabad. Recommended agronomic practices and prophylactic measures were adopted for raising a good crop for field observation.

-	Table 1. Details of the experimental material included for the study										
S.No	Accession No.	State	S.No	Accession No.	State						
1	IC751		31	IC14329							
2	IC16225		32	IC21705							
3	IC16236		33	IC23233							
4	IC16238		34	IC23271)						
5	IC16243	Maharastra(13)	35	IC23321							
6	IC16248		36	IC23325							
7	IC16249		37	IC23327							
8	IC16250		38	IC23332							
9	IC41906		39	IC23335							
10	IC41910		40	IC23341	Madhyapradesh(24)						
11	IC41911		41	IC23346							
12	IC41912		42	IC41932							
13	IC41978		43	IC41948							
14	IC14080		44	IC41953							
15	IC14106		45	IC41962							
16	IC14135	Rajasthan(8)	46	IC41964							
17	IC14155	(U)	47	IC41966							
18	IC14174		48	IC42200							
19	IC26303		49	IC52585							
20	IC42965		50	IC52586							
21	IC42987		51	IC52592							
22	IC14163		52	IC52593							
23	IC43169		53	IC52599							
24	IC43171	Guiarat(8)	54	IC52600							
25	IC43177	Sujuu(0)	55	IC96098							
26	IC43179		56	IC96109	Uttar Pradesh(3)						
27	IC43181		57	IC96113							
28	IC43185		58	IC16832							
29	IC43217		59	IC31379	Punjab(2)						
30	IC20156	Nagaland(1)	60	IC96079	Himachal Pradesh(1)						

. 1

STATISTICAL ANALYSIS

The data recorded were subjected to the following statistical analysis:

ANOVA

Difference between genotypes for various characters was tested for significance by using analysis of variance technique as suggested by Panse and Sukhatme (1957).

Variance

The genotypic and phenotypic variances were calculated as per the formula suggested by Burton and Devane (1953).

Genetic diversity

The genetic diversity in 60 genotypes for characters was estimated using Mahalanobis's (1936) D^2 statistic technique.

Contribution of individual characters towards divergence

In all the combinations, each character was ranked on the basis of its contribution towards divergence between two entries ($d_i = y_{it} - y_{jt}$). Rank- I was given to the highest mean difference and rank-P to the lowest

differences. 'P' is the total number of characters considered.

Heirarchial method of clustering of genotypes into various clusters

Clustering of genotypes into different clusters was done by using Euclidean method.

Intra and inter cluster distances

Based on D^2 values, average intra and inter cluster distances were calculated as per Euclidean method.

RESULTS

Sixty sesame germplasm accessions were characterized for the morphological characters using descriptor guidelines developed by IPGRI. The descriptors were unambiguous and easily identifiable. Characterization was done for each genotype to establish their diagnostic features. The experimental material exhibited large variability for all of the morphological characters.

Mean performance

Mean performance for the thirteen quantitative characters are presented in Table 2.

SPANDANA BANDILA ET AL. / JRAS Vol. 7, No. 2 (2011) 88-99

Trait	Plant height(cm)	Branches/ plant	No. of leaves/ plant	leaf length(cm)	leaf width(cm)	No. of nodes/ main stem	Internode length(cm)	No. of flowers/ plant	No. of flowers/ Axil	No. of capsules/ plant	No. of seeds/ capsule	seed weight(gr)	Seed yield/plant(gr)
Mean	25.6297	4.0628	72.3097	3.2953	1.0820	3.6783	3.7453	43.8588	1.0333	43.4770	50.8995	2.7243	5.6336
C.V.	15.0773	43.5200	21.7463	20.9874	28.6289	45.4369	18.7708	19.9311	0.0000	17.2046	11.3103	5.7380	25.2867
F ratio	1.7242	1.7007	3.2847	1.6387	1.1001	1.8242	1.1609	4.1567	0.0000	6.2651	11.9878	2.6299	4.9843
F Prob.	0.0063	0.0075	0.0000	0.0119	0.3266	0.0029	0.2450	0.0000	1.0000	0.0000	0.0000	0.0000	0.0000
S.E.	2.2310	1.0208	9.0787	0.3993	0.1788	0.9649	0.4059	5.0469	0.0000	4.3186	3.3237	0.0903	0.8225
C.D. 5%	6.2481	2.8589	25.4252	1.1183	0.5008	2.7023	1.1367	14.1341	0.0000	12.0944	9.3083	0.2528	2.3034
C.D. 1%	8.2607	3.7797	33.6149	1.4785	0.6622	3.5728	1.5029	18.6869	0.0000	15.9902	12.3066	0.3342	3.0453
Min	15.6500	1.9000	44.7000	1.3000	0.7000	1.8000	3.0000	22.4000	1.0000	18.5000	22.6000	2.3667	2.3433
Max	32.8333	6.9333	119.2667	4.6000	1.8000	7.4000	5.3333	68.8000	2.0000	67.8667	72.4000	2.9800	10.8600

Table 2. Mean performance for the thirteen quantitative characters

Qualitative characters

Two types of Leaf shapes were observed; entire and lobed. Only one accession showed the lobed type of shape whereas the remaining accessions showed entire type. Flower colour for all the accessions was found to be white with purple shading.

Six colours (white, black, brown, grey, light brown, and reddish brown) of seed were observed. Among the sixty accessions studied, forty were white, one brown, seven reddish brown, two light brown, four grey, and six were black

Capsule beak was long in all the accessions studied.

Analysis of variance

ANOVA showed significant differences for all the traits evaluated. The results of ANOVA are presented in Table 3.

Variability

Among the thirteen characters studied, seed yield (29.14%) recorded higher GCV, followed by nodes/main stem (23.82%), while least GCV was recorded for seed weight (4.23%). Highest PCV was recorded by number of branches/plant (48.34%) and the least PCV by seed weight (7.13%) (Table 4).

The difference between the PCV and GCV values for nodes/main stem was high indicating the influence of environment on these traits. However, the difference between the PCV and GCV values for other characters was low indicating minimum effect of environment.

Sl. No	Character	PCV (%)	GCV (%)	ECV (%)
1	Plant height	16.7	7.4	15.02
2	No. of branches/plant	48.3	21.03	43.52
3	No. of leaves/plant	28.6	18.97	21.7
4	Leaf length	23.11	9.68	20.9
5	Leaf width	29.10	5.2	28.6
6	No. of nodes/main stem	51.3	23.8	45.4
7	Internode length	19.3	4.3	18.8
8	No. of flowers/plant	28.55	20.44	19.93
9	No. of flowers/axil	17.51	17.51	0.0
10	No. of capsules/plant	28.55	22.79	17.20
11	No. of seeds/capsule	24.42	21.64	11.3
12	1000 seed weight	7.12	4.22	5.73
13	Seed yield	38.6	29.14	25.28

Table 3. Coefficient of variability for thirteen characters in 60 sesame genotypes

SPANDANA BANDILA ET AL. / JRAS Vol. 7, No. 2 (2011) 88-99

	df	C/P		LW	РН	B/P	IL	N/M	L/P	F/P	SW	SY	N/A	Ns/c
Replicate	2	246.4*	13.12**	0.1615	172.24**	81.39**	3.65**	110.00**	22.3048	168.0608	0.0382	23.79**	0.0000	27.1854
Treatments	59	350.53**	0.7838*	0.1056	25.7461**	5.3167**	0.5738	5.0956**	812.18**	317.63**	0.06**	10.11**	0.0983**	397.9**
Error	118	55.9511	0.4783	0.0959	14.9326	3.1262	0.4942	2.7933	247.2663	76.4144	0.0244	2.0294	0.0000	33.1417

Table 4. Analysis of variance for thirteen characters of 60 sesame genotypes Mean of Squares

C/P-capsules per plant, B/P-branches per plant, F/P-flowers per plant, Ns/C-number of seeds/capsule, LL-leaf length, IL- internode length, SW-seed weight LW-Leaf width, N/M-nodes per mainstem, SY-seed yield, PH-plant height, L/P-leaves per plant, N/A-number of flowers per axil

**Significant at 1% level

* Significant at 5 % level

Genetic divergence

The quantitative assessment of genetic divergence was made by adopting Mahalanobis D^2 statistics for yield and its contributing characters. D^2 statistic was carried out following the procedure of Rao (1952).

The distribution of 60 genotypes of sesame into different clusters is presented in Table 4.

Average intra and inter cluster distances

The average intra and inter cluster D^2 values are presented in Table 7. Intra cluster values ranged from 10.31 (Cluster II) to 17.32 (Cluster VII) (Figure 2 and 3). From the inter-cluster distances, it can be inferred that highest divergence occurred between Cluster I and VII (64.2), while it was least between Cluster II and Cluster III (16.88) (Table 6).

Tab	Table 5. Distribution of 60 sesame genotypes into different clusters									
Cluster No.	No. of Accessions	Accessions								
Ι	4	IC14080, IC 52600, IC 43185, IC 96098								
II	7	IC 16243, IC 20156, IC 16248, IC43217, IC 52593, IC 23335, IC 14329								
III	7	IC 23271, IC 52592, IC 43169, IC 23332,IC 41964, IC 14174, IC 96079								
IV	5	IC 16236, IC 41948, IC 41932, IC751, IC52599								
V	10	IC 16238, IC 42987, IC 42965, IC 16250, IC 52586, IC 41910, IC 41911, IC 16249, IC 42200, IC 96109								
VI	13	IC 14106, IC43177, IC 14135,IC 41966,IC 21705,IC 23321,IC 23341, IC 16225,96113,IC 14155, IC 41978, IC 41953, IC 26303								
VII	7	IC 16832, IC 23346, IC 41912, IC 23233, IC 41906, IC31379, IC 43181								
VIII	7	IC 23325,IC 52585, IC14163,IC 23327, IC43179, IC41962, IC43179								

	Plant	Branches/	Leaves	Leaf	Leaf	Nodes/ main	Internode	Flowers/	Capsules	No. of seeds/	seed	Seed
	height(cm)	plt	/ Plt	length(cm)	width(cm)	stem	length(cm)	plt	/ plt	capsule	weight(gm)	yield(gm)
Cluster I	25.9	4.90	60.95	3.28	0.988	4.55	3.52	39.96	39.43	70.48	2.59	4.80
ClusterII	26.8	4.62	67.47	3.78	1.104	4.66	3.55	39.67	38.89	55.51	2.64	5.17
ClusterIII	23.2	2.45	61.91	2.60	0.930	2.38	3.92	31.63	30.87	60.95	2.81	4.54
ClusterIV	24.0	4.83	80.17	3.57	1.469	4.11	3.78	29.20	27.83	46.43	2.63	3.49
ClusterV	26.5	4.43	79.80	3.48	1.053	3.91	3.56	51.93	51.46	61.29	2.83	8.41
ClusterVI	25.5	3.49	69.66	3.14	1.079	3.00	3.98	41.88	40.77	43.11	2.73	4.84
ClusterVII	24.9	4.47	68.09	3.28	1.067	4.04	4.03	54.10	53.75	36.97	2.67	5.91
ClusterVIII	27.6	4.20	86.85	3.35	1.053	3.74	3.39	54.88	57.54	41.80	2.76	6.43
Mean	25.6	4.06	72.31	3.29	1.082	3.68	3.74	43.86	43.48	50.90	2.72	5.63
TreatMSS	14.7	4.72	571.6	0.87	0.138	4.26	0.46	655.89	772.24	899.52	0.052	17.98
ErrMSS	7.76	1.38	230.22	0.18	0.021	1.35	0.16	31.83	28.61	29.17	0.017	1.40
F Ratio	1.88	3.43	2.48	4.81	6.46	3.14	2.91	20.60	26.98	30.83	3.01	12.8
Probability	0.09	0.00	0.02	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.01	0.00

Table 6. Cluster means for thirteen characters



Figure 2. Cluster diagram representing diversity for 60 Sesamum genotypes



Figure 3. Ward's minimum variance dendrogram

	Cluster I	Cluster II	Cluster III	Cluster IV	Cluster V	Cluster VI	Cluster VII	Cluster VIII
Cluster I	12.540	20.131	23.357	45.941	30.589	44.831	64.196	54.660
Cluster II		10.318	16.875	22.901	21.350	18.615	33.374	29.877
Cluster III			12.521	27.734	24.222	23.623	45.509	39.550
Cluster IV				13.473	45.598	22.603	45.176	45.437
Cluster V					13.940	28.391	36.924	27.629
Cluster VI						10.816	19.537	21.628
Cluster VII							17.315	21.796
Cluster VIII								17.304

Table 7. Average intra (bold) and inter Euclidean cluster distances

DISCUSSION

The diversity analysis of 60 accessions of *S. indicum* L. carried out by using morphological and the data was analyzed by Cluster analysis.

The Ward's minimum variance dendrogram based on morphological characters showed that accessions of same origin did not group in same cluster. The first cluster showed more subclusters than the second cluster thus showing considerable phenotypic variation of morphological characters indicum. The phenotypic of S. differences may be due to genetic diversity which may in turn be due to allelic diversity. Genetic variability among Indian sesame accessions is very high as shown both for the morphological (Bisht et al., 1998; and Baneriee Kole. 2006) and molecular markers (Bhat et al., 1999; Laurentin and Karlovsky, 2006). The aim of this study was to bring results that could help sesame breeders to select suitable parental material for crossing and increase the efficiency of selection in combination with other diversity data. Number of flowers per leaf axil is one of the mportant characters for plant breeding programs. Most of the varieties had one flower per axil (96.6%). Of the sixty sesame

accessions, only two varieties, IC31379 (Punjab), IC21705 (M.P) bear two flowers per axil as there is not much variation this character is not included for relative contribution of characters towards divergence study. As the number of capsules per plant is one of important contributing character to seed yield of sesame (Ibrahim *et al.*, 1983; Osman, 1989), plants with two flowers per leaf axil are important resources for plant breeding programs.

Variation was least for qualitative characters like leaf shape, flower colour, seed colour and type of capsule beak. There was only one flower colour *i.e* white with purple shading. This type of flower colour was quite similar in Indian sesame collection. The analysis of variance revealed significant difference among the each genotypes for character. indicating the existence of variability among the genotypes for the character studied. The cluster analysis based on agro-morphological traits assigned the sixty sesame germplasm accessions into eight main clusters. A dendogram grouped the sesame accessions into individual groups. The cluster analysisdid not separate the germplasm based on their geographical origins. This result was in agreement with findings of Dixit and Swain (2000) and Gupta et al. (2001). This might be

migration of the sesame materials from one region to another in collection sites through farmer to farmer exchange of seeds. Although sesame has been described as an autogamous plant, recent evidence raises the possibility of natural outcrossing in sesame (Pathirana, 1994, Baydar and Gurel, 1999). Some ecological conditions could also lead to gene flow between from populations different geographical origins.

Sixty genotypes were grouped into eight clusters based on D² values. The pattern of group constellations proved that significant amount of variability existed. It is interesting to note that genotypes representing some differences in their origin were grouped in the same cluster. This was an indication for the absence of relationship between genetic diversity geographic diversity. Similar and results have been reported by Ganesh and Thangavelu (1996), Manivannan and Nadarajan (1996), Swain and Dikshit (1997), Johnjoel *et al.* (1998) and Gupta *et al.* (2001).

The number of seeds per capsule contributed highest towards the divergence followed by number of capsules per plant. The remaining characters showed negligible contribution. Similar observations have been recorded by Alarmelu and Ramanathan (1998) and Sudhakar et al. (2006). In sesame, Solanki and Gupta (2002) reported that seed yield, number of capsules per plant, plant height and 1000 seed weight are the important contributing factors.

There was a wide range of variation in the cluster mean values for most of the characters under study. Therefore a crossing program should be initiated between the genotypes belonging different clusters. The greater the distance between two clusters, the wider the genetic diversity among the parents to be included in hybridization program.

REFERENCES

- Arriel N.H.C., Mauro A.O.D., Arriel E.F., Trevisoli S. H.U., Costa M.M., Bárbaro I.M. and F.R.S. Muniz. 2007. Genetic divergence in sesame based on morphological and agronomic traits. Crop Breeding and Applied Biotechnology, 7: 253-261.
- Alarmelu S. and T. Ramanathan 1998. Genetic divergence and heterosis in sesame (*Sesamum indicum* L,). Journal of Oil Seeds Research, 15: 25-31.
- Banerjee P.P. and P.C. Kole. 2006. Genetic variability for some physiological characters in sesame (*Sesamun indicum* L.) Sesame and Safflower Newsletter, 21: 20–24.
- Baydar H. and F. Gurel.1999. The effects of honey bees on cross-pollination and hybrid seed production in sesame (Sesamum indicum L.). Turkish Journal of Field Crops, 4:21-24.
- Bhat V., P. Babrekar and S. Lakhanpaul. 1999. Study of genetic diversity in Indian and exotic sesame (*Sesamum indicum*) germplasm using random amplified polymorphic DNA (RAPD) markers. Euphytica, 110: 21-33.
- Bisht I., R. Mahajan, T. Loknathan and R. Agrawal. 1998. Diversity in Indian sesame collection and stratification of germplasm accessions in different diversity groups. Genetic Resources and Crop Evolution, 45: 325-335.
- Burton G. W. and E.H. Devane. 1953. Estimating heritability in tall fescue (*Festuca arundinacea*) from replicated clonal material. Agronomy Journal, 45: 478-481.
- Dixit U.N. and D. Swain. 2000. Genetic divergence and heterosis in sesame. Indian Journal of Genetic, 60:213-219.
- Ganesh S. K. and S. Thangavelu. 1996. Genetic divergence in sesame. Madras Agricultural Journal, 82: 263-265.

- Geleta M., T. Bryngelsson and E. Bekele. 2007. Genetic diversity of *Guizotia abyssinica* (L. f.) Cass. (Asteraceae) from Ethiopia as revealed by random amplified polymorphic DNA (RAPD). Genetic Resources Crop Evolution, 54: 601–614.
- Geleta M., T. Bryngelsson and E. Bekele. 2008. Assessment of genetic diversity of *Guizotia abyssinica* (L.f.) Cass. (Asteraceae) from Ethiopia using amplified fragment length polymorphism. Plant Genetic Resources Characterization Utilization, 6: 41–51.
- Gupta R. R., B.M.S. Parihar and P.K. Gupta. 2001. Genetic diversity for some metric characters in sesame(*Sesamum indicum L.*). Crop Research, 21:350-354.
- Ibrahim A.F., D.A.E. Ahmed and S.A. Sharief. 1983. Interrelationships and path-coefficient analysis for some characters in sesame (*Sesamum indicum* L.). Journal of Agronomy Crop Science, 152: 454-459.
- Johnjoel A., S. Alarmelu and S. Thangavelku. 1998. Genetic diversity in sesame (*Sesamum indicum* L.). Journal of Oilseeds Research, 15: 71-75.
- Kobayashi T. 1981. The wild and cultivated species in the genus *Sesamum*. In: Sesame: Status and Improvement. Proceedings of Expert Consultation, Rome, Italy, 8-12 December, 1980. FAO, Rome, Italy. pp: 157-163.
- Laurentin H.E. and P. Karlovsky. 2006. Genetic relationship and diversity in sesame (*Sesamum Indicum* L.) germplasm collection using amplified fragment length polymorphism (AFLP). BMC Genetics, 7:10.
- Liu C.J. 1997 Geographical distribution of genetic variation in *Stylosanthes scabra* revealed by RAPD analysis. Euphytica, 98: 21–27.
- Mahalanobis P.C. 1936. On the generalized distances in statistics. Proc. Nation. Inst. Sci., India, 2:49-55.
- Manivannan N. and N. Nadarajan 1996, Genetic divergence in sesame. Madras Agricultural Journal, 83: 789-790.
- Nayar N.M. and K.L. Mehra. 1970. Sesame its uses, botany, cytogenetic and origin. Economic Botany, 24:20-31.
- Osman H.E. 1989. Heterosis and path coefficient analysis in sesame (*Sesamum indicum L.*). Acta Agronomy Hungry, 38:105-112.
- Panse V.G. and P.V. Sukhatme. 1957. Statistical methods for agricultural workers, Indian Council of Agricultural Research Publication, New Delhi, pp: 87-89.
- Pathirana R .1994. Natural cross-pollination in sesame (Sesamum indicum L.). Plant Breeding, 112:167-170.
- Rao C.R. 1952. Advanced Statistical Methods In Biometrical Research. John Wiley and Sons, New York, U.S.A.
- Solanki Z.S. and D. Gupta. 2002. Genetic divergence for seed yield and other characters in sesame (*Sesamum indicum* L.). Journal of Oilseeds Research, 19: 35-37.
- Sudhakar N., O. Sridevi and P.M. Salimath. 2006. Genetic divergence in sesame (Sesamum indicum L.). Journal of Oilseeds Research, 23:295-296.
- Swain D. and U.N. Dikshit. 1997 Genetic divergence in rabi sesame (Sesamum indicum L.).Indian Journal of Genetic, 57: 296-300.