RESEARCH

Open Access

Oblique propagating whistler mode wave with parallel AC electric field at magnetosphere of **Uranus**

Rama Shankar Pandey^{1*}, Shailendra Kumar² and Mukesh Kumar²

Abstract

Oblique propagating electromagnetic whistler mode instability has been studied having a k vector at angle to ambient magnetic field for bi-Maxwellian distribution function with parallel AC electric field in the magnetosphere of Uranus. The dispersion relation and growth rate have been calculated for plasma parameters suited to the magnetosphere of Uranus by using the method of characteristic solutions and kinetic approach. The effect of AC electric field and temperature anisotropy for obligue propagating electromagnetic whistler mode instability has been reported and discussed in the 'results and discussion section. The comparisons have been made with observations made by Voyager 2. It is found that whistler mode emissions observed at lower radial distances cannot reproduce in our calculations. The present studies should be helpful in making estimates on the scattering properties of whistler mode waves and, thus, contribute to a better understanding of auroral activity in the planetary atmosphere.

Keywords: Whistler mode wave, Magnetosphere, Uranus

Introduction

The Voyager 2 encounter of Uranus in January 1986 provided the opportunity to observe yet another planetary magnetosphere and compare the plasma physical processes taking place there to those occurring in the magnetosphere of the Earth, Jupiter, and Saturn. In particular, in this paper, we survey the electromagnetic emissions observed at Uranus and apply our knowledge of similar emissions at the Earth to establish similarities and differences in the signatures. In this way, we can begin to understand how variations in the configuration of the Uranian magnetosphere might affect changes in the emissions. We can also use the known behavior of the electromagnetic waves as diagnostic tools to lend insight into such questions as the variation of electron density within the magnetosphere and possibly some information about the configuration of the magnetic field itself.

Full list of author information is available at the end of the article



Voyager 2 carried a plasma wave receiver designed to measure the amplitude of the electric fields associated with waves in the frequency range of 10 to 56 Hz [1]. The instrument utilizes a single electric dipole antenna for a sensor. Since no wave magnetic field measurements were made, information concerning the mode of propagation of the detected waves must be deduced from the spectral and temporal characteristics of the emissions, the relationship of the frequency of the emission to characteristic frequencies of the plasma, and comparisons with similar emissions studied at the Earth. Fortunately, most of the waves observed in the magnetospheres of the outer planets thus far are easily associated with terrestrial analogs; hence, much progress can be made even without more sophisticated instrumental and additional sensors. Whistler mode emissions are also triggered by lightning-generated whistler Plasma waves such as lightning-generated whistler provide also important diagnostics tools, from which fundamental plasma parameters such as the electron density can be computed. As is well known, whistler mode waves are electromagnetic waves in a magnetized plasma that propagate through the magnetosphere at frequencies below the

© 2012 Pandey et al.; licensee Springer. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/2.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

^{*} Correspondence: rspandey@amity.edu

¹Department of Applied Physics, Amity Institute of Applied Science, Amity University, Noida, Sector-125, Uttar Pradesh 201303, India

electron-cyclotron frequency and electron plasma frequency. Because magnetic field introduces anisotropy in the plasma, whistler waves tend to be guided along the ambient magnetic field [2,3]. Recently, up-to-date views of plasma waves at each outer planets have been reviewed by Zarka [4].

An initial overview of plasma waves observed in the Uranian magnetoshere was given by Gurnett et al. [5,6] and a general comparison with plasma waves observed at Jupiter and Saturn is included [7]. The observations of electromagnetic waves in the magnetosphere of Uranus have been studied, focusing primarily on the Bernstein or $(n + 1/2) f_c$ emissions, where f_c is the electron cyclotron frequency f_c (Hz) = 28 |B| (nT). Bernstein waves were first observed at the Earth by Kennel et al. [8] and have been studied extensively both experimentally and theoretically by a large number of investigators [9-11] Survey of electrostatic waves at Jupiter and Saturn were given by Kurth et al. [12,13]. In recent past, Kurth [14,15] and Kurth and Gurnett [16] have provided for the first time, overviews of comparative study of plasma wave spectra at the outer planetary magnetosphere. They have also reported various modes of wave. Gurnett et al. [17] and Kurth [15] have reported that significant whistler mode emissions could exist near the magnetic equator in the intermediate radial distance ranges say $3 < R < 8 R_N$ as in the other planetary magnetospheres. The intensity of these emissions is quite weak from 10 to $30 \mu V/m$. The variation of field intensity, magnetic field strength, and electron cyclotron frequency with radial distance has similar characteristics as occurred in Uranus.

An analysis of the Voyager 2 plasma science experiment electron measurements made during the Uranus encounter of a thermal (cold) plasma ($K_{\rm B}T$ is approximately 10 eV) and non-Maxwellian suprathermal (hot) plasma ($K_{\rm B}T$ is approximately 1 KeV) component [18] could not be explained in terms of an internal source provided by the Uranian satellites. The source of the thermal ion component could be either the Uranian ionosphere or the 'Uranus' hydrogen corona observed by the ultraviolet spectrometer on Voyager 2 [19] if some acceleration mechanism which heats the newly picked up ions from 1 to 10 eV is present. It has also been suggested by Brice and Lucas [20] that cold plasma plays an important role in controlling the wave particle interactions in the radiation belts within the plasma sphere because of the possibility of excitation of whistler mode instability by injection of cold plasma in the magnetosphere. The modifications in the whistler mode instability produced by cold plasma injection to an infinite uniform anisotropic distribution function plasma in the presence of a perpendicular AC electric field have been studied by [21] in the Uranus.

Pandey et al. [22] have studied oblique whistler mode instabilities for a generalized drifted distribution function in the presence of perpendicular AC electric field by the method of characteristic solution. The growth rate was evaluated for the plasma parameters suited to the magnetosphere of Uranus. It was found that only AC frequencies significantly affect the growth characteristic of the whistler waves, in addition to the external triggering AC field [23,24] studied, by taking into account the finite size of the cold plasma enhancement region, the temporal evolution of whistler growth in a time-dependent cold plasma injection experiment and made semi-quantitative estimates of the magnitude and duration of enhancement of very low frequency (VLF) noise and electron precipitation effects. The results of these studies have shown that the injected cold plasma modifies the propagation characteristics of whistlers, and there is a possibility to have stabilization through finite electron density.

A theoretical study of relativistic whistler mode with perpendicular AC electric field has been studied for the generalized distribution function at the Uranian bow shock by Pandey et al. [25]. The effect of cold plasma injection and temporal evolution of whistler mode instability have been studied in the presence of perpendicular AC electric field for the generalized distribution function in the magnetosphere of Uranus [21,24,25].

In this paper, oblique propagating electromagnetic whistler mode wave has been studied having a \mathbf{k} vector at an angle to ambient magnetic field for bi-Maxwellian distribution function with parallel AC electric field. The dispersion relation and growth rate have been calculated for plasma parameters suited to the magnetosphere of Uranus.

Dispersion relation

The plasma under consideration is of infinite extent embedded with a uniform magnetic field \mathbf{B}_0 \hat{e}_z and an electric field $E_0 \sin vt$ \hat{e}_z and whose distribution function $f(\mathbf{r}, \mathbf{v}, t)$ is governed by the Boltzmann equation. In the collisionless case, the self-consistent Vlasov-Maxwell equations to be used are

$$\frac{\partial f_s}{\partial t} + \nu \cdot \frac{\partial f_s}{\partial r} + \frac{e_s}{m_s} \left[E + \frac{\nu \times \mathbf{B}}{c} \right] \frac{\partial f_s}{\partial \nu} = 0 \tag{1}$$

and

$$\nabla \cdot \mathbf{E} = \sum_{s} 4\pi e_{s} \int d^{3} v f_{s}$$

$$\nabla \times \mathbf{B} = \sum_{s} \frac{4\pi e_{s}}{c} \int d^{3} v v f_{s} + \frac{1\partial \mathbf{E}}{c\partial t}$$

$$\nabla \cdot \mathbf{B} = 0$$

$$\nabla \times \mathbf{E} = -\frac{1}{c} \frac{\partial B}{\partial t}$$
(2)

Here, the subscript *s* denotes the particle species, and *i*, *e*, for ions and electrons, respectively. The electric field

www.SID.ir

 \mathbf{E}_0 and the magnetic field \mathbf{B}_0 represent the resultants of the external AC field and magnetic field together with the self-consistent characteristic oscillations of \mathbf{E} and \mathbf{B} , and all other symbols have their usual meanings.

In this analysis, AC frequency has been assumed much smaller than the plasma and gyrofrequency and the collision frequency negligibly small and independent of particle velocity, where as the thermal velocity of the background electrons are much larger than the induced velocity due to AC field. The directed motion of the electrons due to AC field along the magnetic field may lead to the appearance of an electromagnetic field whose effect is neglected.

In order to find the dispersion relation, first, the motion of the particles (trajectories) in the given fields are computed, and the perturbed distribution function, which depends on the integrals of motion, is then determined following a method similar to that adopted by Harris [26]. Following the methods and techniques of Misra and Pandey [27] and Misra and Haile [28], the unperturbed particle trajectories are obtained as

$$x_{0} = x + \frac{v_{y}}{\omega_{cs}} - \frac{1}{\omega_{cs}} \left(v_{x} \sin \omega_{cs} t' + v_{y} \cos \omega_{cs} t' \right)$$

$$y_{0} = y + \frac{v_{x}}{\omega_{cs}} - \frac{1}{\omega_{cs}} \left(v_{x} \cos \omega_{cs} t' + v_{y} \sin \omega_{cs} t' \right)$$

$$z_{0} = z - v_{z} t' + \frac{\Gamma_{z}}{v^{2}} \sin v t' - \frac{\Gamma_{z}}{v} t'$$

and the velocities as

$$\nu_{x_0} = \nu_x \cos\omega_{cs} t' - \nu_y \sin\omega_{cs} t'$$

$$\nu_{y_0} = \nu_x \sin\omega_{cs} t' - \nu_y \cos\omega_{cs} t'$$

$$\nu_{z_0} = \nu_z + \frac{\Gamma_z}{v} (\cos v t' - 1)$$
(4)

where $\omega_{cs} = e_s B_0/m_s$ is the cyclotron frequency and $\Gamma_z = e_s E_0/m_s$. From Maxwell's curl **E** equation, for the perturbed quantities, one gets the following:

$$\mathbf{B}_{1} = \frac{c}{\omega} (\mathbf{k} \times \mathbf{E}_{1})$$

$$S(\mathbf{r}, \mathbf{v}, t) = \frac{-e_{s}}{m_{s}\omega} e^{i(k.r-\omega t)} [(\omega - \mathbf{k} \cdot \mathbf{v})\mathbf{E}_{1} + (\nu \cdot \mathbf{E}_{1})\mathbf{k}] \cdot \frac{\partial f_{0}}{\partial \nu},$$
(5)

where $E_1 = E_1 e^{i(k. r - \omega t')}$ has been used.

Inverting Equation 5, one can get the following:

$$S(\mathbf{r_0}, \mathbf{v_0}, t - t') = \frac{e_s}{m_s \omega} e^{i \left(\mathbf{k} \cdot \mathbf{r}(\mathbf{r}, \mathbf{v}, t') - \omega(t - t') \times \left[(\omega - \mathbf{k} \cdot \mathbf{v_0})\mathbf{E}_1 + (\mathbf{v_0} \cdot \mathbf{E}_1)\mathbf{k}\right] \frac{\partial f_0}{\partial v}},$$
(6)

where Bessel identity $e^{i\lambda\sin\alpha} = \sum_{k=-\infty}^{\infty} j_k(\lambda) e^{ik\alpha}$ has been used.

The result of Equation 6 is used to obtain the first order perturbed distribution function $f_1(r, \mathbf{v}, t)$.

$$f_{1}(r, \mathbf{v}, t) = \frac{-ie}{m\omega} \sum_{m,n,p} \frac{J_{m}(\lambda_{1})J_{p}(\lambda_{2})e^{i(m-n)\alpha}}{\left(\omega - k_{||}v_{||} - \frac{k_{||}\Gamma_{z}}{v} - n\omega_{c} + pv\right)} \left[U^{*}\mathbf{E}_{1x}\frac{\mathbf{n}}{\lambda_{1}}J_{n}(\lambda_{1}) - iU^{*}\mathbf{E}_{1y}J_{n}'(\lambda_{1}) + W^{*}\mathbf{E}_{1z}J_{n}(\lambda_{1}) \right],$$
(7)

where α is the angle between the vectors **k** and **B**₀.

$$U^{*} = \frac{1}{\nu_{\perp}} \frac{\partial f_{0}}{\partial \nu_{\perp}} \left[\left(\omega - k_{\parallel} \nu_{\parallel} \right) \nu_{\perp} + \frac{k_{\parallel} \Gamma_{z}}{v} \nu_{\parallel} \left(1 - \frac{p}{\lambda_{2}} \right) \right] \\ + k_{\parallel} \nu_{\perp} \frac{\partial f_{0}}{\partial \nu_{\parallel}} W^{*} = \frac{1}{\nu_{\perp}} \frac{\partial f_{0}}{\partial \nu_{\perp}} \left[\nu_{\parallel} n \omega_{c} - \frac{\Gamma_{z}}{v} n \omega_{c} \left(1 - \frac{p}{\lambda_{2}} \right) \right] \\ \lambda_{1} = \frac{k_{\perp} \nu_{\perp}}{\omega}, \ \lambda_{2} = \frac{k_{\parallel} \Gamma_{z}}{v^{2}} \text{ and } J_{n}'(\lambda_{1}) = \frac{d J_{n}(\lambda_{1})}{d \lambda_{1}}.$$
(8)

The current density is calculated using the relation

$$=\sum_{s}e_{s}\int d^{3}\nu f_{1}(\nu)\nu \tag{9}$$

or in terms of the conductivity $||\sigma(\mathbf{k},\omega)||$ by

$$I = \|\sigma(\mathbf{k}, \omega)\| \cdot \mathbf{E}_1.$$
(10)

Using Equations 7, 9, and 10, the conductivity is written in matrix form as

$$\sigma(k,\omega) = -i\sum_{np} \frac{e^2}{m\omega} J_p(\lambda_2) \frac{S_{ij}}{\omega - k_{||}v_{||} - \frac{k_{||}\Gamma_z}{v} + pv - n\omega_c},$$
(11)

where

$$S_{ij} = \begin{bmatrix} \nu_{\perp} U^* \left(\frac{n}{\lambda_1} J_n\right)^2 & i \nu_{\perp} U^* \frac{n}{\lambda_1} J_n J'_n & \nu_{\perp} W^* \frac{n}{\lambda_1} J_n^2 \\ i \nu_{\perp} U^* \frac{n}{\lambda_1} J_n J'_n & \nu_{\perp} U^* J'_n 2 & i \nu_{\perp} W^* J_n J'_n \\ \nu_{||} U^* \frac{n}{\lambda_1} J_n^2 & -i \nu_{||} U^* J_n J'_n & \nu_{||} W^* J_n^2 \end{bmatrix}.$$

Now, from Maxwell's curl equations, the dielectric tensor is given as

$$\varepsilon_{ij} = 1 + \sum_{np} \frac{4 \pi e^2}{m\omega^2} J_p(\lambda_2) \frac{S_{ij} d^3 \nu}{\left(\omega - k_{||}\nu_{||} - \frac{k_{||}\Gamma_z}{\nu} + p\nu - n\omega_c\right)}.$$
(12)
(12)

www.SID.ir

The generalized dielectric tensor may be written as

$$\begin{bmatrix} N^2 \cos^2 \theta_1 + \varepsilon_{11} & \varepsilon_{12} & N^2 \cos \theta_1 \sin \theta_1 + \varepsilon_{13} \\ \varepsilon_{21} & N^2 \varepsilon_{22} & \varepsilon_{23} \\ N^2 \cos \theta_1 \sin \theta_1 + \varepsilon_{31} & \varepsilon_{33} & N^2 \sin^2 \theta_1 + \varepsilon_{33} \end{bmatrix}.$$

If we remove the contribution of AC electric field above, the dielectric tensor is similar to Sazhin [29]. After using the limits $k_{\perp} = ksin\theta_1 \rightarrow 0$ and $k_{\parallel} = kcos\theta_1$, the generalized dielectric tensor becomes a simplified tensor and the dispersion relation is reduced as

$$\begin{bmatrix} -N^2 + \varepsilon_{11} & \varepsilon_{12} & 0 \\ -\varepsilon_{21} & -N^2 + \varepsilon_{22} & 0 \\ 0 & 0 & \varepsilon_{33} \end{bmatrix} .$$

This is rewritten in a more convenient form. Now, for whistler wave

$$-N^4 - 2\epsilon_{11}N^2 + \epsilon_{11}^2 + \epsilon_{12}^2 = 0$$
 (13)

for electrostatic wave, $\varepsilon_{33} = 0$.

Neglecting the higher power of *N*, the resulting relation becomes as follows:

$$\varepsilon_{11} \pm \varepsilon_{12} = N^2 \tag{14}$$

$$f_0 = \frac{n_0}{\pi^2 \alpha_\perp^2 \alpha_\parallel} \exp\left[-\left(\frac{\nu_\perp}{\alpha_\perp}\right)^2 - \left(\frac{\nu_\parallel}{\alpha_\parallel}\right)^2\right],\tag{15}$$

where the parallel and perpendicular thermal velocities are

$$\alpha_{\perp} = \left(\frac{K_{\rm B}T_{\perp}}{m}\right)^{\frac{1}{2}}, \ \alpha_{||} = \left(\frac{K_{B}T_{||}}{m}\right)^{\frac{1}{2}}, \ A_{\rm T} = \frac{\alpha_{\perp}^{2}}{\alpha_{||}^{2}} - 1.$$

Using Equations 14 and 15 and doing lengthy velocity integrations, now the dispersion relation of oblique whistler wave is obtained from above for n = 1, p = 1 and $J_p = 1$, $J_q = 1$:

$$D(k,\omega) = 1 + \frac{\omega_p^2}{\omega^2} \left\{ \frac{\omega + \omega^*}{k_{||}\alpha_{||}} Z(\xi) + A_T (1 + \xi Z(\xi)) \right\},$$
(16)

where $Z(\xi)$ is the plasma dispersion function

 $k_{||}\alpha_{||}$

$$Z(\xi) = i\sqrt{\pi} \exp(\xi)^2 - \frac{1}{\xi} \left(1 + \frac{1}{2\xi^2}\right)$$
(17)
$$c \qquad \omega - n\omega_c + pv - \frac{k_{\parallel}\Gamma_z}{v}$$

Now, using the asymptotic expansion of plasma dispersion function, the dispersion relation may be separated in real and imaginary form of dispersion relation for limit $k^2c^2 / \omega^2 >> 1$:

$$D_{I} = \left[\frac{\omega + \omega^{*}}{k_{||}\alpha_{||}}\xi + A_{T}\xi\right]\sqrt{\pi}exp\left(-(\xi)^{2}\right)$$
(18)

$$D_{R} = -\frac{k^{2}c^{2}}{\omega^{2}} + \left\{ \frac{\omega + \omega^{*}}{k_{||}\alpha_{||}} \left(-\frac{1}{\xi} \left(1 + \frac{1}{2\xi^{2}} \right) + A_{T} \left(-\left(1 + \frac{1}{2\xi^{2}} \right) \right) \right) \right\}.$$
(19)

Using the standard definition of growth rate

$$\gamma/\omega_{\rm c} = \frac{-\mathrm{Im}D(k,\omega)}{\omega_{\rm c}\frac{\partial\mathrm{R}D(k,\omega)}{\partial\omega}}\bigg|_{\omega=\omega_{\rm r}}$$
(20)

after using Equations 17 to 20, the required growth rate and real frequency dimensionless form is written as follows:

$$\gamma/\omega_{c} = \frac{\frac{\sqrt{\pi}}{K_{1}} \left(A_{T} - \frac{X_{3} + X_{4}}{K_{3}}\right) K_{3}^{3} \left(-\frac{K_{3}}{K_{1}}\right)^{2}}{1 + \frac{K_{1}^{2}}{2K_{3}^{2}} - \frac{K_{1}^{2}}{K_{3}} \left(A_{T} - \frac{X_{3} + X_{4}}{K_{3}}\right)},$$
(21)

where

$$X_{3} = \frac{\omega_{r}}{\omega_{c}} = \frac{K_{1}^{2}}{\beta} \left[1 + X_{4} + \frac{A_{T}\beta}{1 + X_{4}} \right] - X_{4}$$

$$X_{4} = \frac{\omega^{*}}{\omega_{c}}, \quad \omega^{*} = pv - \frac{k_{||} \Gamma_{z} - k_{||} \nu_{d}}{v}$$

$$\overline{k} = \frac{k_{||}\alpha_{||}}{\omega_{c}}, \quad K_{3} = 1 - X_{3} + X_{4}$$

$$\beta = \frac{K_{B}T_{||} \mu_{0}n_{0}}{B_{0}^{2}}, \quad K_{1} = \overline{k}\cos\theta_{1}.$$
(22)

Results and discussion

Numerical calculations for the growth rate of the whistler instabilities with plasma parameters suited to the Uranian magnetosphere [21,24,30] $B_0 = 5 \times 10^{-10}$ T, $K_{\rm B}T_{||} = 1$ KeV, $n_0 = 5 \times 10^4$ m⁻³, $A_{\rm T} = .25$ and external field parameters E_0 from 10 MV/m to 30 mV/m with frequency v from 400 to 800 Hz have been carried out. The growth rate of whistler waves in the presence of a parallel AC electric field have been displayed in a series of curves from (Figures 1,2,3,4,5).

Figure 1 shows the dependence of growth rate of whistler waves with respect to a dimensionless parameter ($\overline{k} = k_{||}\alpha_{||}/\omega_c$) for various values of AC field frequency v. The introduction of the AC signal for fixed values of other plasma parameters increases the magnitude of the growth rate while increasing the possible range of wave propagation. The contribution of AC frequency reduces the upper limit of wave number \overline{k} and normalized real frequency $X_3 = \omega_r / \omega_{ci}$ and increases the growth rate. The AC frequency appears only through modification of resonant instability. This shows that the **www.SID.ir**

presence of a minimum value of parallel AC field frequency is enough to trigger the whistler emission [27,28]. This result is different from the magnetosphere; it is only due to the low values of the Earth's magnetic field. In Figure 2, the dependence of growth rate on various values of thermal anisotropy has been plotted. It is seen that for other fixed plasma parameters, an increase in thermal anisotropy increases the growth rate. For a fixed value of temperature anisotropy increases the growth rate initially increases rapidly, reaches a maxima and then levels off with a small decrease from the maxima. Temperature anisotropy is the prime source of instability. Thermal electrons in bi-Maxwellian distribution are supposed to modify the intensity and frequency of electron plasma [3]. Figure 3 shows the variation of growth rate with \overline{k} for different values of the number density n_0 . The increase in the number density for other fixed plasma parameters increases the

0.1 0.01 ν/ως A, =0.75 A₁=0.5 1.000E-03 1.000E-04 0.2 0.4 0.6 0.8 k Figure 2 Variation of growth rate with respect to $\overline{\mathbf{k}}$ for various values of temperature anisotropy. At other fixed plasma parameters.

growth, and the maxima shifted toward the higher values of \overline{k} . The effect of the number density is an important role because it changes the plasma frequency, and modification has been made in the value of β . It means that the value of β is lower in comparison with the magnetosphere.

Figure 4 shows the dependence of growth rate with \overline{k} on various values of external electric field. Increase of the magnitude of AC field increases the growth rate and widens the bandwidth significantly. Figure 5 shows the variation of growth rate with \overline{k} for various values of angle of propagation θ_1 at other fixed plasma parameters. In this case, increasing the angle of propagation increases the growth rate. After 40°, the electrostatic conditions start dominating and non-resonant instability





0.2

0.4

Figure 1 Variation of growth rate with respect to \overline{k} for various

values of AC field frequency. At other fixed plasma parameters.

k

v=6KHz

0.6

=4KH2

0.8

0.1

0.01

1.000E-03

1.000E-04

٥



0.8

starts contributing or even change in polarization may take place making it as a right-handed mode.

Conclusion

The results are compared with the magnetospheric study which has been done by Misra and Pandey [27] and Misra and Haile [28]. The perpendicular electric field is modifying the perpendicular velocity and contributing to the energy exchange significantly to the emission of VLF signals and can explain the low frequency side of the spectrum. In this case, only AC frequency is effective for growth rate, but in the case of parallel AC electric field, the Doppler shift in frequency is affected by both magnitude and AC frequency. In the case of Uranus, the growth rate is affected by AC frequency as well as the magnitude of parallel AC electric field. However, in the case of Uranus, low value of AC frequency is enough to trigger the whistler mode instability. The growth rate is also affected by the magnitude of AC electric field due to the low value of gyrofrequency in Uranus with respect to the magnetosphere. In Uranus, the density of plasma particle is less than the magnetosphere. Hence, the low value of β is enough to the generation and propagation of whistler mode emission. The effect of temperature anisotropy is the same for Uranus and the magnetosphere. From all figures, it is clear that our results are very much in order of the observed normalized frequency range of whistler emission at Uranus as reported by Coroniti et al. [31]. It is clear that whistler mode propagates even at the low magnitude of magnetic field and other Uranus condition. In the case of Misra and Haile [28], the propagation of wave along the direction of magnetic field is considered, but in this paper, oblique incidence. The magnitude of growth rate has been changed, and band increases with respect to wave number.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

In the present work, RSP participated in the sequence alignment and drafted the manuscript. SK helped draft the manuscript. MK participated in the computer programming and mathematical analysis. All authors read and approved the final manuscript.

Acknowledgments

The excellent expert comments of the referees and editor are gratefully acknowledged.

Author details

¹Department of Applied Physics, Amity Institute of Applied Science, Amity University, Noida, Sector-125, Uttar Pradesh 201303, India. ²Department of Physics, Nalanda College-Biharsharif, Magadh University, Bodh Gaya, Bihar 824234, India.

Received: 30 March 2012 Accepted: 13 October 2012 Published: 22 November 2012

References

- Scarf, FL, Gurnett, DA: A plasma wave investigation for the Voyager mission. Space Sci. Rev. 21, 289 (1977)
- Gurnett, DA, Kurth, WS, Cairns, IH, Granroth, LJ: Whistlers in Neptune's magnetosphere: evidence of atmospheric lightning. J. Geophys. Res. 95, 20967 (1990)
- Stenzel, RL: Whistler waves in space and laboratory plasmas. J. Geophys. Res. 104(A7), 14379 (1999)
- Zarka, P: Radio and plasma waves at the outer planets. Adv. Space Res. 33, 2045 (2004)
- Gurnett, DA, Kurth, WS, Scarf, FL, Poynter, RL: First plasma wave observations at Uranus. Science 233, 106 (1986)
- Gurnett, DA, Kurth, WS, Scarf, FL, Poynter, RL, Granroth, LJ, Cairns, IH, Macek, WM, Moses, SL, Coroniti, FV, Kennel, CF, Barbosa, DD: First plasma wave observations at Neptune. Science 246, 1494 (1989)
- Scarf, FL, Gurnett, DA, Kurth, WS, Coroniti, FV, Kennel, CF, Poynter, RL: Plasma wave measurements in the magnetosphere of Uranus. J. Geophys. Res. 92, 15217 (1987)
- Kennel, CF, Scarf, FL, Fredricks, RW, McGehee, JH, Coroniti, FV: VLF electric field observations in the magnetosphere. J. Geophys. Res. 75, 6136 (1970)
- Christiansen, PJ, Gough, MP, Martelli, G, Bloch, JJ, Cornilleau, N, Etcheto, J: GEOS-1 observations of electrostatic waves, and their relationship with plasma parameters. Space Sci. Rev. 22, 383 (1978)
- Ashour-Abdalla, M, Kennel, CF: Nonconvective and convective electron cyclotron harmonic instabilities. J. Geophys. Res. 83, 1531 (1978)
- Kennel, CF, Ashour-Abdalla, M: Electrostatic waves and the strong diffusion of magnetospheric electrons. In: Nishida (ed.) Magnetospheric Plasma Physics, p. 245. Center for Academic Publications, Tokyo (1982)
- 12. Kurth, WS, Barbosa, DD, Gurnett, DA, Scarf, FL: Electrostatic waves in the Jovian magnetosphere. Geophys. Res. Lett. **7**, 57 (1980)
- Kurth, WS, Scarf, FL, Gurnett, DA, Barbosa, DD: A survey of electrostatic waves in Saturn's magnetosphere. J. Geophys. Res. 88, 8959 (1983)
- Kurth, WS: Voyager plasma wave observations near the outer planets. Adv. Space Res. 11, 59 (1991)
- Kurth, WS: Comparative observations of plasma waves at the outer planets. Adv. Space Res. 12, 83 (1992)
- Kurth, WS, Gurnett, DA: Plasma waves in planetary magnetospheres. J. Geophys. Res. 96, 18977 (1991)
- Gurnett, DA, Kurth, WS, Granroth, LJ, Cairns, IH, Macek, WM, Poynter, RL, Moses, SL, Coroniti, FV, Kennel, CF, Barbosa, DD: Plasma wave observations at Neptune. Adv. Space Res. 12(11), 47 (1992)
- Bridge, HS, Belcher, JW, Coppi, B, Lazarus, AJ, McNutt Jr, RL, Olbert, S, Richardson, JD, Sands, MR, Selesnick, RS, Sullivan, JD, Hartle, RE, Ogilvie, KW, Sittler Jr, EC, Bagenal, F, Wolff, RS, Vasyliunas, VM, Siscoe, GL, Goertz, CK, Eviatar, A: Plasma observations near Uranus: initial results from Voyager 2. Science 233, 89 (1986)
- Broadfoot, AL, Herbert, F, Holberg, JB, Hunten, DM, Kumar, S, Sandel, BR, Shemansky, DE, Smith, GR, Yelle, RV, Strobel, DF, Moos, HW, Donahue, TM, Atreya, SK, Bertaux, JL, Blamont, JE, McConnell, JC, Dessler, AJ, Linick, S,

www.SID.ir



Springer, R: Ultraviolet spectrometer observations of Uranus. Science 233, 74 (1986)

- Brice, NM, Lucas, C: Influence of magnetospheric convection and polar wind loss of electrons from the outer radiation belt. J. Geophys. Res. 76, 900 (1971)
- Pandey, RP, Md Karim, S, Singh, KM, Pandey, RS: Effect of cold plasma injection on whistler mode instability triggered by perpendicular Ac electric field at Uranus. Earth, Moon, and Planets 91(4), 195 (2002)
- Pandey, RP, Singh, KK, Singh, KM, Pandey, RS: Current driven oblique whistler wave in the magnetosphere of Uranus. Indian Journal of Phys. 73B(4), 593 (1999)
- 23. Ganguli, G, Palmadesso, P, Feeder, J: Temporal evolution of whistler growth in a cold plasma injection experiment. J. Geophys. Res. **78**, 8150 (1984)
- Pandey, RP, Pandey, RS, Misra, KD: Temporal evolution of Whistler instability due to cold plasma injection in the presence of perpendicular AC electric field in the magnetosphere of Uranus. Earth, Moon, and Planets 91(4), 209 (2002)
- Pandey, RP, Singh, KM, Pandey, RS: A theoretical study of the whistler mode instability at the Uranian bow-shock. Earth Moon and Planets 87, 59 (2001)
- Harris, EG: Plasma instability, Chapter 4. Physics of Hot Plasmas, p. 158. Oliver and Boyd, Edinburgh (1970)
- Misra, KD, Pandey, RS: Generation of whistler emissions by injection of hot electrons in the presence of a perpendicular AC electric field. J. Geophys. Res. 100, 19405 (1995)
- Misra, KD, Haile, T: Effect of AC electric field on the whistler mode instability in the magnetosphere. J. Geophys. Res. 98(A6), 9297 (1993)
- Sazhin, SS: Oblique whistler mode growth and damping in a hot anisotropic plasma. Planets. Space Sci. 36, 663 (1988)
- Smith, CW, Wong, HK, Goldstein, ML: Whistler waves associated with the Uranian bow shock: outbound observations. J. Geophys. Res. 96, 15841 (1991)
- Coroniti, FV, Kurth, WS, Scarf, FL, Krimigis, SM, Kennel, CF, Gurnett, DA: Whistler mode emissions in the Uranian radiation belts. J. Geophys. Res. 92, 15234 (1987)

doi:10.1186/2251-7235-6-31

Cite this article as: Pandey *et al.*: **Oblique propagating whistler mode wave with parallel AC electric field at magnetosphere of Uranus**. *Journal of Theoretical and Applied Physics* 2012 **6**:31.

www.SID.ir