

# Current State of Art Management for Vascular Complications after Liver Transplantation

S. Kamran Hejazi Kenari<sup>1</sup>, Asha Zimmerman<sup>1</sup>, Mohammad Eslami<sup>2</sup>, Reza F. Saidi<sup>1,\*</sup>

1. Division of Organ Transplantation, Department of Surgery, Alpert Medical School of Brown University, Providence, RI, USA
2. Division of Vascular Surgery, Department of Surgery, Boston University School of Medicine, Boston, MA, USA

## ABSTRACT

Vascular complications by compromising the blood flow to the allograft can have significant and sometimes life-threatening consequences for the patient. High level of suspicion and aggressive utilization of diagnostic modalities can lead to early diagnosis and salvage of the allograft. This review will summarize the current trends in the management of vascular complications after liver transplantation. Current trends show an increase in the utilization of endovascular interventions initially to address vascular complications after liver transplantation. Operative repair still has its major role, especially if endovascular procedures fail.

## KEYWORDS

Liver Transplantation; Vascular Complications; Endovascular interventions; Outcomes

Please cite this paper as:

S. Hejazi Kenari K, Zimmerman A, Eslami M, F. Saidi R. Current State of Art Management for Vascular Complications after Liver Transplantation. *Middle East J Dig Dis* 2014;6:121-30.

## INTRODUCTION

Vascular complications are rare but serious causes of morbidity and mortality after liver transplantation (LT). Bleeding, stenosis, thrombosis, and aneurysm can arise at any of the vascular anastomoses. The incidence is generally about 8-15%.<sup>1</sup> However this rate can be as high as 20%,<sup>2</sup> especially in cases such as split liver transplantation or LDLT.<sup>2-8</sup> Hepatic artery thrombosis (HAT) and portal vein thrombosis (PVT) interrupt the allograft's blood supply and produce early allograft loss, longterm dysfunction, or even patient's death.

Arterial complications are the most common (5-10%) vascular complication after LT. Early HAT generally needs re-transplantation while venous complications including portal and caval problems are less frequent (each about 2%) and can usually be treated by surgical or endovascular intervention.<sup>1</sup> Vascular problems must be treated aggressively, particularly HAT and PVT which can interrupt the liver's blood supply and cause dysfunction and even graft loss or death.<sup>9</sup>

Many factors have been described as the cause of these vascular complications such as technical problems in anastomosis, problems in allograft anatomy which may result in vascular kinking, and differences in the size of donor's and recipient's vessels.<sup>3,10,11</sup> Similar to other arterial anastomoses, the process of intimal hyperplasia may affect the patency of the arterial anastomosis in long term.

### \* Corresponding Author:

Reza F. Saidi, MD, FICS, FACS  
 Assistant Professor of Surgery, Division of Organ Transplantation, Department of Surgery, Alpert Medical School of Brown University, 593 Eddy Street, Providence, RI 02903  
 Tel: (401) 444 4861  
 Fax: (401) 444 3283  
 Email: RSaidi@lifespan.org  
 Received: 28 Apr. 2014  
 Accepted: 30 May 2014

### Arterial Complications

Arterial problems are the most common vascular complications after LT. These problems include HAT, hepatic artery stenosis (HAS), and hepatic artery kinks (HAK).<sup>12</sup> HAS and HAT can lead to allograft ischemia, which carries a high mortality and morbidity rate. Arterial complications are often diagnosed firstly by Duplex followed by CT angiography. Role of Doppler ultrasonography in the early diagnosis of HAS has been evaluated in several reports. Doppler ultrasonography showed a sensitivity of 100%, a specificity of 99.5%, a PPV of 95% and NPV of 100%, and overall accuracy of 99.5% in early diagnosis of HAS.<sup>13</sup> Many protocols also use angiography to confirm the diagnosis.<sup>14</sup>

Many patients with HAS are asymptomatic and most commonly present only with abnormal liver function tests (LFT). Compared with HAST (hepatic artery stenosis), risk of development of biliary complications are less with HAS. Balloon angioplasty can be an effective treatment option in these cases.<sup>15</sup> Anastomotic stenosis is the most common place for development of hepatic artery stenosis within 3 months after LT.<sup>13</sup>

Hepatic Artery thrombosis represents more than 50% of all arterial complications following LT.<sup>9,12,14,16-20</sup> HAT presentations may be a continuum from only abnormal transaminase, to fever and sepsis due to biliary dysfunction and graft failure.<sup>9</sup> Based on the interval between LT and development of thrombosis, HAT can be divided into early HAT (within 4 weeks) and late HAT. The rate of re-transplantation in untreated HAT is 25-83% while it is 28-35% in patients who underwent re-vascularization.<sup>21-29</sup> Early HAT may be the result of technical problems and can have dramatic presentation.<sup>15</sup> Because early HAT has a higher mortality compared with late HAT, related to the ischemia/necrosis of bile duct and subsequent sepsis, an emergent intervention is required.<sup>12</sup> Earlier studies showed a higher rate of early HAT after LT probably due to limited initial experience and also the insidious nature of late HAT.<sup>12</sup> Up to 50% of patients with late HAT can be asymptomatic with only elevated LFT.<sup>21,30</sup> Symptomatic patients often present with biliary

complications with recurrent cholangitis, abscess and biliary leakage or stricture, and the presentation may be insidious. Late HAT is usually due to ischemic or immunologic injuries.<sup>15</sup> Patients with late symptomatic HAT can be initially treated with biliary stent placement and/or endovascular interventions.<sup>15</sup>

Duffy and colleagues<sup>9</sup> evaluated 4,234 LT from 1984 to 2007. Of them 203 (5%) developed HAT including 133 early and 70 late HAT. Occurrence of HAT was 3.9% in adults. Overall 90 patients were treated with surgical exploration, thrombectomy, or anastomotic revision. Nine patients were treated with catheter-based thrombolysis and 13 patients received anticoagulation. Of the patients with early HAT who underwent thrombectomy and anastomotic revision, only 9 (10.5%) had graft salvage and the remaining patients needed re-transplantation. Overall, re-transplantation was necessary in 153 (75%) patients with HAT. However re-transplantation after HAT had a better survival rate compared with revision or thrombolysis.<sup>9</sup>

Factors associated with HAT can be divided into several categories (table 1). Technical problems are mainly associated with early HAT. Positive CMV antibody in donor and negative CMV in recipients have been shown to be associated with late HAT.<sup>30-33</sup> On the other hand, while some authors believe that HAS and HAK are the initiating factor,<sup>34</sup> others suggest perioperative hypercoagulable state as a possible underlying cause.<sup>9,12,16,20</sup> Other factors leading to HAT include longer cold ischemia, longer operative time, shorter warm ischemia, transfusion of more than 6 units of whole blood and/or  $\geq 15$  units of fresh frozen plasma (FFP).

Use of aortic conduit may also be a risk factor for development of HAT. While some earlier studies indicate that there is no difference in the results when aortic conduits are used,<sup>35</sup> more recent studies emphasize a higher risk of HAT in those with aortic conduits.<sup>36</sup> Duffy and co-workers also reported a higher incidence of HAT with aorto-hepatic grafting compared with standard reconstructions from the celiac trunk.<sup>9</sup> However, other studies reported no difference between these two approaches. Ni-

**Table 1: Risk factors associated with development of hepatic artery thrombosis after liver transplantation.**<sup>6,9,16,30,34,38-41</sup>

Donor factors	<ul style="list-style-type: none"> <li>Abnormal donor arterial anatomy</li> <li>CMV (+) donor while recipient is CMV(-)</li> </ul>
Recipient factors	<ul style="list-style-type: none"> <li>Abnormal recipient arterial anatomy</li> <li>Immunological or genetic factors</li> <li>Infection</li> <li>Celiac stenosis or compression by median arcuate ligament</li> <li>Multiple rejections</li> </ul>
Peri-operative factors	<ul style="list-style-type: none"> <li>Cold ischemia time</li> <li>Warm ischemia time</li> <li>Ischemia-reperfusion injury</li> </ul>
Operational factors	<ul style="list-style-type: none"> <li>Technical problems in creating an anastomosis</li> <li>Performing complex back-table arterial reconstruction</li> <li>Increased number of FFP and/or whole blood transfusion</li> <li>Aortic conduit for arterial re-construction</li> <li>Roux-en-Y biliary reconstruction</li> <li>Roux-en-Y biliary reconstruction</li> </ul>
Post-operative factors	<ul style="list-style-type: none"> <li>Bile leak</li> <li>Cholangitis</li> <li>Re-laparotomy</li> </ul>

kitin and colleagues,<sup>37</sup> for example, reported graft survival rate of 59% with arterial conduit and 67% without the conduit in 5-year follow-up and 33% and 35%, respectively, in 15 years of follow-up. With 20 years of follow-up, there was no significant difference in graft or patients' survival or incidence of hepatic artery or biliary complications.<sup>37</sup>

Other factors unrelated to operation or operative technique may also play a role in developing HAT. For example, use of transarterial chemoembolization (TACE) as a "bridging" or "down staging" therapy before transplantation for treatment of hepatocellular carcinoma (HCC) can induce periarterial inflammation which in turn can lead to HAT.<sup>9,42</sup>

Although urgent re-transplantation is considered the main treatment for early HAT, endovascular interventions including PTA, intra-arterial thrombolysis (IAT) in selective cases, and stent placement may be alternative treatments. Currently, in many centers, percutaneous transluminal angioplasty (PTA) and stent placement are tried first to resolve the problem.<sup>43,44</sup>

Studies have shown that both PTA and stent placement have comparable results for treatment of HAS (table 2).<sup>45</sup> According to some reports use of PTA for management of HAS can reduce the rate of HAT more than threefold.<sup>46</sup> While solitary short

stenotic lesions are usually most amenable to PTA, angioplasty has a high success rate in tandem lesions.<sup>46,47</sup> It is also important to consider that while recent reports have shown encouraging results of endovascular interventions,<sup>12</sup> the efficacy and risk of complications of these procedures particularly potential risk of hemorrhage are still controversial and in some cases these endovascular interventions are ineffective and surgical intervention (including anastomotic reconstruction or even re-transplantation) must be applied.<sup>1</sup> The complications of PTA include thrombosis, vascular dissection or even rupture.<sup>3</sup>

Intra-arterial infusion of thrombolytic agents can also be used in selective cases of HAT. Intra-arterial thrombolysis (IAT) is contraindicated in patients who has had recent (<2 weeks) abdominal operations. It is necessary to carefully monitor the use of thrombolytic agents by evaluating either fibrinogen level, prothrombin time (PT), or activated partial thromboplastin time (aPTT).<sup>12</sup> Hemorrhage is the most common complication of IAT. Singhal and co-workers<sup>12</sup> performed a systematic literature search on endovascular treatment of HAT, which had been done before June 2009. In their review the success rate of thrombolysis was 68% (47 out of 69) and 29 out of the 47 patients underwent further interven-

**Table 2: Summary of studies with utilization of endovascular interventions for arterial complications after liver transplantation**

Author/ref.	Year published	Number of cases (EP)	Type of vascular problem	Endovascular intervention(s)	Success rate
Orons <sup>59</sup>	1995	21	HAS	PTA	81% (17/21)
Abbasoglu <sup>60</sup>	1997	6	HAS	PTA	100%
Saad <sup>46</sup>	2005	42	HAS	PTA	81% (34/42)
Jeon <sup>61</sup>	2008	4	HAT	Catheter-directed thrombolysis	50%
Jeon <sup>61</sup>	2008	19	Peritoneal hemorrhage	Trans-catheter arterial embolization	84% (16/19)
Jiang <sup>49</sup>	2008	3	HAT, HAS and HAP	Thrombectomy, Balloon angioplasty/ stent placement and embolization (respectively)	100%
Chen <sup>62</sup>	2009	20	HAS	PTA + Stent placement	100%
Maruzzelli <sup>64</sup>	2010	25	HAS	PTA or stenting	96% (24/25)
Pérez-Saborido <sup>1</sup>	2011	4	HAS	Angioplasty and stent placement	25%
Sabri <sup>65</sup>	2011	35	HAS	For HAS: balloon angioplasty +/- stent placement	91%
Sabri <sup>65</sup>	2011	6	HAT	Catheter-directed thrombolysis	17% (1/6)
Kim <sup>58</sup>	2011	6	HAT, HAS	3 IAT (for HAT) 3 PTA (for HAS)	100%
Abdelaziz <sup>56</sup>	2012	11	HAT	IAT or PTA	82% (9/11)
Hamby <sup>66</sup>	2013	35	HAS	PTA alone or primary stent placement	97%
Sommacale <sup>67</sup>	2013	37	HAS	Stent placement	100%

tion including PTA or stent placement or both. Additionally, percutaneous mechanical thrombectomy of HAT can be an alternative when IAT is contraindicated. In these cases, the thrombus is dissolved and removed by intraluminal suction device.

Vascular complications in patients receiving live donor liver transplantation (LDLT) are unique to these complicated reconstructions. These patients have a higher rate of vascular complications,<sup>48</sup> because of the complexity of vascular reconstruction of smaller and shorter vessels.<sup>49</sup> In recipients of LDLT, the hepatic artery is short and surgical field is deep and the graft artery's anatomical arrangement may be different from the recipient's hepatic artery.<sup>50</sup> These vascular complications expose the recipients of LDLT to a higher risk of developing biliary leakage, resistant infection, and other morbidities.<sup>51</sup> HAT can be developed in few hours after the LDLT.<sup>52</sup> There are also reports of simultaneous HAT and PVT following LDLT, which is an ultimately fatal complication.<sup>53</sup> Careful perioperative evaluation and intraoperative microsurgical technique for vascular reconstructions can reduce

the chance of vascular complications following LDLT.<sup>49</sup> Yan and co-workers<sup>54</sup> reported their experience in 101 HA reconstruction in LDLT using microsurgical techniques. Only 2 patients (2%) in that group developed HAT. There are also several recent studies that showed excellent results after endovascular interventions for treatment of arterial complications following LDLT.<sup>48,49,51-66</sup> Furuta and colleagues also reported using of microsurgical techniques to create 44 artery anastomoses in 40 LDLT patients. They dissected the recipient arteries that were slightly smaller than the graft's artery and created an end-to-end anastomosis. No decreases in the arterial blood flow or HAT were reported.<sup>50</sup>

### Venous Complications

Venous complications after LT include caval/hepatic vein and portal problems. In clinical settings, portal vein stenosis leads to portal vein hypertension, which manifests with ascites, anemia, splenomegaly, hypersplenism, and GI bleeding.<sup>44,68-70</sup> Hepatic vein obstruction is clinically similar to Budd-Chiari syndrome.<sup>3</sup> Hepatic vein outflow ob-

struction (HVOO) is a general term reflecting any obstruction of the HV caused by either compression and twisting of the anastomosis resulted from graft regeneration or by intimal hyperplasia and fibrosis at the anastomotic sites.<sup>71</sup>

There are many reports regarding the incidence and treatment of venous complications after LDLT.<sup>48,53,57,71-75</sup> Although the incidence of venous complications can be even more than 2 times higher in recipients of LDLT,<sup>48</sup> acceptable long-term graft and patient's survival can be achieved after endovascular or operational treatments. In a study by Kyoden and co-workers 5-year survival rate after portal vein complications were 77%.<sup>73</sup> The success rate of endovascular interventions for the treatment of venous complications varies from 60% to 100%.<sup>71-73</sup>

Factors associated with PVT include technical problems, small diameter of the portal vein (< 5 mm), previous splenectomy, simultaneous thrombectomy for pre-existing PVT and use of venous conduits for portal vein reconstruction.<sup>9,76-79</sup> Additionally, longer cold ischemia time (>12 h) can be a risk factor for developing venous problems. This can be due to difficulties in venoplasty (and more manipulation) before anastomosis.<sup>71</sup> Kyoden and colleagues analyzed the outcome of 310 LDLT in their institution.<sup>73</sup> Portal vein complications were identified in 28 recipients. Presence of PVT at the time of LDLT, smaller graft size, and use of vein grafts were the risk factors for PV complications. Additionally they found out that the use of jump or interposition cryo-preserved vein grafts predisposed to the occurrence of late, but not early, PV complications.<sup>73</sup>

Use of cryopreserved vein for portal conduits has been shown to be associated with portal vein stenosis.<sup>80</sup> In a study by Buell and co-workers,<sup>80</sup> use of cryopreserved vein for portal conduit led to 6.3% PVS, which decreased to 1% when cryopreserved vein was not used. They showed that the use of segmental grafts without venous conduits did not cause a significant increase in the rate of long term venous complication.

Pre-existing venous complications can also

cause post-operative PVS and PVT. Lerut and colleagues<sup>10</sup> reported their experience of 393 LT from 1980 to 1984. Nine out of 313 patients (2.9%) had pre-existing vena cava abnormalities, which needed adjustment during operation. These included absent retro-hepatic IVC (n=2), extra-hepatic IVC (n=2), absent SVC (n=1), thrombosis of IVC (n=2), and aneurysm of IVC (n=2). Additionally, 51 patients (16.3%) had pre-existing portal vein abnormalities, which included thrombosis (n=22), hypoplasia (n=20), phlebosclerosis (n=8), and absent portal vein (n=1). These preoperative conditions may necessitate adjunct procedures at the time of transplantation such as thrombectomy and retrograde dissection of the abnormal portal vein to the confluence of the splenic and superior mesenteric veins. The graft portal vein is then anastomosed to this confluence directly or with a graft of the donor's iliac vein, pulmonary artery, or IVC. Seven patients developed post-operational portal vein thrombosis. Four of these patients had an early convalescence. Portal flow was restored in only one of them. These 4 patients had a pre-existing and undiagnosed thrombosis of the portal vein, splenic vein, or superior mesenteric vein. Overall the incidence of thrombosis of the reconstructed portal vein was only 1.8%. Only 3 patients (0.8%) developed post-operative IVC thrombosis, all at the lower IVC anastomosis.<sup>10</sup>

Compared with arterial problems, venous complications usually have a better response to endovascular interventions such as angioplasty or stent placement (table 3). Transcatheter endovascular procedures are now considered as the first line of treatments for post-transplant portal vein complications and many studies showed the high successful results.<sup>68,81-85</sup> The initial success rate of hepatic vein PTA is higher than 80%.<sup>3</sup> The standard approach for treatment of portal vein complications is via transhepatic access of intrahepatic branch of portal vein, however in case of concurrent portal vein thrombosis this approach is challenging and percutaneous trans-splenic access can be substituted, particularly in patients with short occluded portal vein branches.<sup>16,77,78,81-92</sup>

**Table 3:** Summary of studies with utilization of endovascular interventions for venous complications after liver transplantation

Author/ref.	Year published	Number of cases	Type of vascular problem	Endovascular intervention(s)	Success rate
Shin <sup>74</sup>	2006	11	Interposition vein graft (IVG) stenosis	Stent placement	10/11 (91%)
Yang <sup>72</sup>	2009	6	Intra-operative HV tension, HVS, PVS	Balloon dilation and stent placement	100%
Cheng <sup>78</sup>	2010	16	PVS and PV occlusion	Stent placement	68.8% (11/16)
Pérez-Saborido <sup>1</sup>	2011	6	PV, PVS and Caval complications	Angioplasty and stent placement	83% (5/6)
Umehara <sup>71</sup> (A)	2012	5	HV outflow obstruction	Balloon angioplasty and stent placement	100%

The use of anticoagulants after endovascular procedures for venous complications is controversial. While some surgeons use heparin for 1-3 days postoperation the others use it just at the time of surgery.<sup>3</sup> Aspirin is usually used continuously afterward. It is also important to know that re-transplantation after PVT particularly an extensive one, is more challenging than re-transplantation after HAT. The reason is that re-transplantation after PVT requires patent mesenteric graft inflow to provide hepatotrophic factors which is not always easy in such patients.<sup>9</sup>

#### Aneurysmal complications

Aneurysm and pseudoaneurysm of the hepatic artery (HA) are very rare complications after LT. The rate of post-operational aneurysm or pseudoaneurysm of is much less than other post-transplant vascular complications but they are associated with high mortality (>50%).<sup>15</sup> These complications generally require emergency surgical or angiographic interventions. The diagnosis is usually made by splanchnic angiography or perfusion angiosonography.<sup>93</sup> Aneurysmal complications can be divided to extra- and intrahepatic complications. The most common presentation of extrahepatic aneurysm or pseudoaneurysm is hypotension following its rupture. Bleeding can happen into gastrointestinal system or into peritoneal cavity. They can also be presented by the signs of biliary compression and obstruction.<sup>15</sup> Intrahepatic arterial aneurysms are most frequently iatrogenic and happen following

liver biopsy or stent placement.<sup>15</sup> While many of these patients can be treated by coil embolization in case of bleeding and hemodynamic shock, urgent laparotomy is required.<sup>15</sup>

Aneurysmal complications can be treated by either surgical or endovascular procedures.<sup>58,94-96</sup> Both surgery and angioembolization are equally effective for hepatic artery pseudoaneurysm. However, patients who undergo angioembolization have more rapid bleeding control, shorter hospital stay, and less transfusion requirements.

In summary, the incidence of vascular complications reported in the literature varies widely among centers. Despite technical progress, vascular complications are still a significant determinant of allograft loss, increasing postoperative morbidity and mortality. Arterial complications are more common, occur early in the postoperative period, and are associated with high rates of graft loss and patient mortality. Conversely, venous complications are less frequent, occur late in the postoperative period, and have no significant effect on graft loss or mortality rates. Strategies for identification and mitigation of risk factors, prevention of technical complications, and protocols for early detection of vascular complications may reduce mortality, morbidity, and the need for re-transplantation. Current trends showed an increase utilization of endovascular interventions initially to address vascular complications after LT with good success. Operative repair still has its role, especially if endovascular procedures fail.

## CONFLICT OF INTEREST

The authors declare no conflict of interest related to this work.

## REFERENCES

- Pérez-Saborido B, Pacheco-Sánchez D, Barrera-Rebollo A, Asensio-Díaz E, Pinto-Fuentes P, Sarmentero-Prieto JC, et al. Incidence, management, and results of vascular complications after liver transplantation. *Transplant Proc* 2011;**43**:749-50.
- Sieders E, Peeters PM, TenVergert EM, de Jong KP, Porte RJ, Zwaveling JH, et al. Early vascular complications after pediatric liver transplantation. *Liver Transpl* 2000;**6**:326-32.
- Carnevale FC, Borges MV, Pinto RA de P, Oliva JL, Andrade W de C, Maksoud JG. Endovascular treatment of stenosis between hepatic vein and inferior vena cava following liver transplantation in a child: a case report. *Pediatr Transplant* 2004;**8**:576-80.
- Emond JC, Whittington PF, Thistlethwaite JR, Alonso EM, Broelsch CE. Reduced-size orthotopic liver transplantation: use in the management of children with chronic liver disease. *Hepatol Baltim Md* 1989;**10**:867-72.
- Sindhi R, Rosendale J, Mundy D, Taranto S, Baliga P, Reuben A, et al. Impact of segmental grafts on pediatric liver transplantation--a review of the United Network for Organ Sharing Scientific Registry data (1990-1996). *J Pediatr Surg* 1999;**34**:107-110; discussion 110-111.
- Wozney P, Zajko AB, Bron KM, Point S, Starzl TE. Vascular complications after liver transplantation: a 5-year experience. *AJR Am J Roentgenol* 1986;**147**:657-63.
- Zajko AB, Claus D, Clapuyt P, Esquivel CO, Moulin D, Starzl TE, et al. Obstruction to hepatic venous drainage after liver transplantation: treatment with balloon angioplasty. *Radiology* 1989;**170**:763-5.
- Goss JA, Shackleton CR, McDiarmid SV, Maggard M, Swenson K, Seu P, et al. Long-term results of pediatric liver transplantation: an analysis of 569 transplants. *Ann Surg* 1998;**228**:411-20.
- Duffy JP, Hong JC, Farmer DG, Ghobrial RM, Yersiz H, Hiatt JR, et al. Vascular complications of orthotopic liver transplantation: experience in more than 4,200 patients. *J Am Coll Surg* 2009;**208**:896-903; discussion 903-5.
- Lerut J, Tzakis AG, Bron K, Gordon RD, Iwatsuki S, Esquivel CO, et al. Complications of venous reconstruction in human orthotopic liver transplantation. *Ann Surg* 1987;**205**:404-14.
- Mazariegos GV, Garrido V, Jaskowski-Phillips S, Towbin R, Pigula F, Reyes J. Management of hepatic venous obstruction after split-liver transplantation. *Pediatr Transplant* 2000;**4**:322-7.
- Singhal A, Stokes K, Sebastian A, Wright HI, Kohli V. Endovascular treatment of hepatic artery thrombosis following liver transplantation. *Transpl Int* 2010;**23**:245-56.
- Uller W, Knoppke B, Schreyer AG, Heiss P, Schlitt HJ, Melter M, et al. Interventional Radiological Treatment of Perihepatic Vascular Stenosis or Occlusion in Pediatric Patients After Liver Transplantation. *Cardiovasc Intervent Radiol* 2013;**36**:1562-71.
- Margarit C, Hidalgo E, Lázaro JL, Murio E, Charco R, Balsells J. Biliary complications secondary to late hepatic artery thrombosis in adult liver transplant patients. *Transpl Int* 1998;**11 Suppl 1**:S251-4.
- Porrett PM, Hsu J, Shaked A. Late surgical complications following liver transplantation. *Liver Transpl* 2009;**15 Suppl 2**:S12-8.
- Tzakis AG, Gordon RD, Shaw BW Jr, Iwatsuki S, Starzl TE. Clinical presentation of hepatic artery thrombosis after liver transplantation in the cyclosporine era. *Transplantation* 1985;**40**:667-71.
- Langnas AN, Marujo W, Stratta RJ, Wood RP, Shaw BW Jr. Vascular complications after orthotopic liver transplantation. *Am J Surg* 1991;**161**:76-82; discussion 82-3.
- Drazan K, Shaked A, Olthoff KM, Imagawa D, Jurim O, Kiai K, et al. Etiology and management of symptomatic adult hepatic artery thrombosis after orthotopic liver transplantation (OLT). *Am Surg* 1996;**62**:237-40.
- Saad WEA, Davies MG, Saad NEA, Westesson KE, Patel NC, Sahler LG, et al. Catheter thrombolysis of thrombosed hepatic arteries in liver transplant recipients: predictors of success and role of thrombolysis. *Vasc Endovascular Surg* 2007;**41**:19-26.
- Bekker J, Ploem S, de Jong KP. Early hepatic artery thrombosis after liver transplantation: a systematic review of the incidence, outcome and risk factors. *Am J Transplant* 2009;**9**:746-57.
- Bhattacharjya S, Gunson BK, Mirza DF, Mayer DA, Buckels JA, McMaster P, et al. Delayed hepatic artery thrombosis in adult orthotopic liver transplantation-a 12-year experience. *Transplantation* 2001;**71**:1592-6.
- Langnas AN, Marujo W, Stratta RJ, Wood RP, Li SJ, Shaw BW. Hepatic allograft rescue following arterial thrombosis. Role of urgent revascularization. *Transplantation* 1991;**51**:86-90.
- Lerut J, Gordon RD, Iwatsuki S, Starzl TE. Surgical complications in human orthotopic liver transplantation. *Acta Chir Belg* 1987;**87**:193-204.
- Stange BJ, Glanemann M, Nuessler NC, Settmacher U, Steinmüller T, Neuhaus P. Hepatic artery thrombosis after adult liver transplantation. *Liver Transpl* 2003;**9**:612-20.
- Sheiner PA, Varma CV, Guarrera JV, Cooper J, Garatti M, Emre S, et al. Selective revascularization of hepatic artery thromboses after liver transplantation improves patient and graft survival. *Transplantation* 1997;**64**:1295-9.
- Pinna AD, Smith CV, Furukawa H, Starzl TE, Fung JJ. Urgent revascularization of liver allografts after early hepatic artery thrombosis. *Transplantation* 1996;**62**:1584-7.

27. Cavallari A, Vivarelli M, Bellusci R, Jovine E, Mazziotti A, Rossi C. Treatment of vascular complications following liver transplantation: multidisciplinary approach. *Hepato-gastroenterology* 2001;**48**:179-83.
28. Pawlak J, Wróblewski T, Małkowski P, Nyckowski P, Zieniewicz K, Grzelak I, et al. Vascular complications related to liver transplantation. *Transplant Proc* 2000;**32**:1426-8.
29. Karatzas T, Lykaki-Karatzas E, Webb M, Nery J, Tsaroucha A, Demirbas A, et al. Vascular complications, treatment, and outcome following orthotopic liver transplantation. *Transplant Proc* 1997;**29**:2853-5.
30. Silva MA, Jambulingam PS, Gunson BK, Mayer D, Buckels JA, Mirza DF, et al. Hepatic artery thrombosis following orthotopic liver transplantation: a 10-year experience from a single centre in the United Kingdom. *Liver Transpl* 2006;**12**:146-51.
31. Gunsar F, Rolando N, Pastacaldi S, Patch D, Raimondo ML, Davidson B et al. Late hepatic artery thrombosis after orthotopic liver transplantation. *Liver Transpl* 2003;**9**:605-11.
32. Valente JF, Alonso MH, Weber FL, Hanto DW. Late hepatic artery thrombosis in liver allograft recipients is associated with intrahepatic biliary necrosis. *Transplantation* 1996;**61**:61-5.
33. Oh CK, Pelletier SJ, Sawyer RG, Dacus AR, McCullough CS, Pruett TL, et al. Uni- and multi-variate analysis of risk factors for early and late hepatic artery thrombosis after liver transplantation. *Transplantation* 2001;**71**:767-72.
34. Segel MC, Zajko AB, Bowen A, Bron KM, Skolnick ML, Penkrot RJ, et al. Hepatic artery thrombosis after liver transplantation: radiologic evaluation. *AJR Am J Roentgenol* 1986;**146**:137-41.
35. Goldstein RM, Secrest CL, Klintmalm GB, Husberg BS. Problematic vascular reconstruction in liver transplantation. Part I. Arterial. *Surgery* 1990;**107**:540-3.
36. Del Gaudio M, Grazi GL, Ercolani G, Ravaioli M, Varotti G, Cescon M, et al. Outcome of hepatic artery reconstruction in liver transplantation with an iliac arterial interposition graft. *Clin Transplant* 2005;**19**:399-405.
37. Nikitin D, Jennings LW, Khan T, Sanchez EQ, Chinnakotla S, Randall HB, et al. Twenty years of follow-up of aortohepatic conduits in liver transplantation. *Liver Transpl* 2008;**14**:1486-90.
38. Brems JJ, Millis JM, Hiatt JR, Klein AS, Quinones-Baldrich WJ, Ramming KP, et al. Hepatic artery reconstruction during liver transplantation. *Transplantation* 1989;**47**:403-6.
39. Mas VR, Fisher RA, Maluf DG, Wilkinson DS, Garrett CT, Ferreira-Gonzalez A. Hepatic artery thrombosis after liver transplantation and genetic factors: prothrombin G20210A polymorphism. *Transplantation* 2003;**76**:247-9.
40. Mazzaferro V, Esquivel CO, Makowka L, Belle S, Kahn D, Koneru B, et al. Hepatic artery thrombosis after pediatric liver transplantation--a medical or surgical event? *Transplantation* 1989;**47**:971-7.
41. Tzakis AG. The dearterialized liver graft. *Semin Liver Dis* 1985;**5**:375-6.
42. Yao FY, Kinkhabwala M, LaBerge JM, Bass NM, Brown R Jr, Kerlan R, et al. The impact of pre-operative loco-regional therapy on outcome after liver transplantation for hepatocellular carcinoma. *Am J Transplant* 2005;**5**:795-804.
43. Zajko AB, Sheng R, Bron K, Reyes J, Nour B, Tzakis A. Percutaneous transluminal angioplasty of venous anastomotic stenoses complicating liver transplantation: intermediate-term results. *J Vasc Interv Radiol JVIR* 1994;**5**:121-6.
44. Carnevale FC, de Tarso Machado A, Moreira AM, Dos Santos AC, da Motta-Leal-Filho JM, Suzuki L, et al. Long-term results of the percutaneous transhepatic venoplasty of portal vein stenoses after pediatric liver transplantation. *Pediatr Transplant* 2011;**15**:476-81.
45. Rostambeigi N, Hunter D, Duval S, Chinnakotla S, Golzarian J. Stent placement versus angioplasty for hepatic artery stenosis after liver transplant: a meta-analysis of case series. *Eur Radiol* 2013;**23**:1323-34.
46. Saad WEA, Davies MG, Sahler L, Lee DE, Patel NC, Kitanosono T, et al. Hepatic artery stenosis in liver transplant recipients: primary treatment with percutaneous transluminal angioplasty. *J Vasc Interv Radiol* 2005;**16**:795-805.
47. Harman A, Boyvat F, Hasdogan B, Aytakin C, Karakayali H, Haberal M. Endovascular treatment of active bleeding after liver transplant. *Exp Clin Transplant* 2007;**5**:596-600.
48. Khalaf H. Vascular complications after deceased and living donor liver transplantation: a single-center experience. *Transplant Proc* 2010;**42**:865-70.
49. Jiang X-Z, Yan L-N, Li B, Zhao JC, Wang WT, Li FG, et al. Arterial complications after living-related liver transplantation: single-center experience from West China. *Transplant Proc* 2008;**40**:1525-8.
50. Furuta S, Ikegami T, Nakazawa Y, Hashikura Y, Matsunami H, Kawasaki S, et al. Hepatic artery reconstruction in living donor liver transplantation from the microsurgeon's point of view. *Liver Transpl Surg* 1997;**3**:388-93.
51. Unal B, Gonultas F, Aydin C, Otan E, Kayaalp C, Yilmaz S. Hepatic artery thrombosis-related risk factors after living donor liver transplantation: single-center experience from Turkey. *Transplant Proc* 2013;**45**:974-7.
52. Kim BW, Won JH, Lee BM, Ko BH, Wang HJ, Kim MW. Intraarterial thrombolytic treatment for hepatic artery thrombosis immediately after living donor liver transplantation. *Transplant Proc* 2006;**38**:3128-31.
53. Kaneko J, Sugawara Y, Togashi J, Kishi Y, Akamatsu N, Makuuchi M. Simultaneous hepatic artery and portal vein thrombosis after living donor liver transplantation. *Transplant Proc* 2004;**36**:3087-9.
54. Yan S, Zhang Q-Y, Yu Y-S, He JJ, Wang WL, Zhang M, et al. Microsurgical reconstruction of hepatic artery in living donor liver transplantation: experiences and lessons. *Hepatobiliary Pancreat Dis Int* 2009;**8**:575-80.



55. Rodriguez-Davalos MI, Arvelakis A, Umman V, Tanjavur V, Yoo PS, Kulkarni S, et al. Segmental Grafts in Adult and Pediatric Liver Transplantation: Improving Outcomes by Minimizing Vascular Complications. *JAMA Surg* 2014;**149**:63-70.
56. Abdelaziz O, Hosny K, Amin A, Emadeldin S, Uemoto S, Mostafa M. Endovascular management of early hepatic artery thrombosis after living donor liver transplantation. *Transpl Int* 2012;**25**:847-56.
57. Azzam AZ, Tanaka K. Management of vascular complications after living donor liver transplantation. *Hepatogastroenterology* 2012;**59**:182-6.
58. Kim SJ, Yoon YC, Park JH, Oh DY, Yoo YK, Kim DG. Hepatic artery reconstruction and successful management of its complications in living donor liver transplantation using a right lobe. *Clin Transplant* 2011;**25**:929-38.
59. Orons PD, Zajko AB, Bron KM, Trecha GT, Selby RR, Fung JJ. Hepatic artery angioplasty after liver transplantation: experience in 21 allografts. *J Vasc Interv Radiol JVIR* 1995;**6**:523-9.
60. Abbasoglu O, Levy MF, Vodapally MS, Goldstein RM, Husberg BS, Gonwa TA, et al. Hepatic artery stenosis after liver transplantation--incidence, presentation, treatment, and long term outcome. *Transplantation* 1997;**63**:250-5.
61. Jeon GS, Won JH, Wang HJ, Kim BW, Lee BM. Endovascular treatment of acute arterial complications after living-donor liver transplantation. *Clin Radiol* 2008;**63**:1099-1105.
62. Chen GH, Wang GY, Yang Y, Li H, Lu MQ, Cai CJ, et al. Single-center experience of therapeutic management of hepatic artery stenosis after orthotopic liver transplantation. Report of 20 cases. *Eur Surg Res* 2009;**42**:21-7.
63. Singhal A, Mukherjee I, Stokes K, Wright HI, Sebastian A, Kohli V. Continuous intraarterial thrombolysis for early hepatic artery thrombosis following liver transplantation: case report. *Vasc Endovascular Surg* 2010;**44**:134-8.
64. Maruzzelli L, Miraglia R, Caruso S, Milazzo M, Mamone G, Gruttadauria S, et al. Percutaneous endovascular treatment of hepatic artery stenosis in adult and pediatric patients after liver transplantation. *Cardiovasc Intervent Radiol* 2010;**33**:1111-9.
65. Sabri SS, Saad WEA, Schmitt TM, Turba UC, Kumer SC, Park AW, et al. Endovascular therapy for hepatic artery stenosis and thrombosis following liver transplantation. *Vasc Endovascular Surg* 2011;**45**:447-52.
66. Hamby BA, Ramirez DE, Loss GE, Bazan HA, Smith TA, Bluth E, et al. Endovascular treatment of hepatic artery stenosis after liver transplantation. *J Vasc Surg* 2013;**57**:1067-72.
67. Sommacale D, Aoyagi T, Dondero F, Sibert A, Bruno O, Fteriche S, et al. Repeat endovascular treatment of recurring hepatic artery stenoses in orthotopic liver transplantation. *Transpl Int* 2013;**26**:608-15.
68. Carnevale FC, Borges MV, Moreira AM, Cerri GG, Maksud JG. Endovascular treatment of acute portal vein thrombosis after liver transplantation in a child. *Cardiovasc Intervent Radiol* 2006;**29**:457-61.
69. Funaki B, Rosenblum JD, Leef JA, Hackworth CA, Szymiski GX, Alonso EM, et al. Portal vein stenosis in children with segmental liver transplants: treatment with percutaneous transhepatic venoplasty. *AJR Am J Roentgenol* 1995;**165**:161-5.
70. Funaki B, Rosenblum JD, Leef JA, Hackworth CA, Szymiski GX, Alonso EM. Angioplasty treatment of portal vein stenosis in children with segmental liver transplants: mid-term results. *AJR Am J Roentgenol* 1997;**169**:551-4.
71. Umehara M, Narumi S, Sugai M, Toyoki Y, Ishido K, Kudo D, et al. Hepatic venous outflow obstruction in living donor liver transplantation: balloon angioplasty or stent placement? *Transplant Proc* 2012;**44**:769-71.
72. Yang J, Xu M-Q, Yan L-N, Lu WS, Li X, Shi ZR, et al. Management of venous stenosis in living donor liver transplant recipients. *World J Gastroenterol* 2009;**15**:4969-73.
73. Kyoden Y, Tamura S, Sugawara Y, Matsui Y, Togashi J, Kaneko J, et al. Portal vein complications after adult-to-adult living donor liver transplantation. *Transpl Int* 2008;**21**:1136-44.
74. Shin JH, Sung K-B, Yoon H-K, Ko GY, Kim KW, Lee SG, et al. Endovascular stent placement for interposed middle hepatic vein graft occlusion after living-donor liver transplantation using right-lobe graft. *Liver Transpl* 2006;**12**:269-76.
75. Cheng Y-F, Ou H-Y, Tsang LL-C, Yu CY, Huang TL, Chen TY, et al. Vascular stents in the management of portal venous complications in living donor liver transplantation. *Am J Transplant* 2010;**10**:1276-83.
76. Charco R, Fuster J, Fondevila C, Ferrer J, Mans E, Garcia-Valdecasas JC. Portal vein thrombosis in liver transplantation. *Transplant Proc* 2005;**37**:3904-5.
77. Kishi Y, Sugawara Y, Matsui Y, Akamatsu N, Makuuchi M. Late onset portal vein thrombosis and its risk factors. *Hepatogastroenterology* 2008;**55**:1008-9.
78. Lladó L, Fabregat J, Castellote J, Ramos E, Torras J, Jorba R, et al. Management of portal vein thrombosis in liver transplantation: influence on morbidity and mortality. *Clin Transplant* 2007;**21**:716-21.
79. Seu P, Shackleton CR, Shaked A, Imagawa DK, Olthoff KM, Rudich SR, et al. Improved results of liver transplantation in patients with portal vein thrombosis. *Arch Surg* 1996;**131**:840-4; discussion 844-5.
80. Buell JF, Funaki B, Cronin DC, Yoshida A, Perlman MK, Lorenz J, et al. Long-term venous complications after full-size and segmental pediatric liver transplantation. *Ann Surg* 2002;**236**:658-66.
81. Yilmaz A, Arıkan C, Tumor G, Kilic M, Aydogdu S. Vascular complications in living-related and deceased donation pediatric liver transplantation: single center's experience from Turkey. *Pediatr Transplant* 2007;**11**:160-4.

82. Bertram H, Pfister E-D, Becker T, Schoof S. Transsplenic endovascular therapy of portal vein stenosis and subsequent complete portal vein thrombosis in a 2-year-old child. *J Vasc Interv Radiol JVIR* 2010;**21**:1760-4.
83. Funaki B, Rosenblum JD, Leef JA, Zaleski GX, Farrell T, Lorenz J, et al. Percutaneous treatment of portal venous stenosis in children and adolescents with segmental hepatic transplants: long-term results. *Radiology* 2000;**215**:147-51.
84. Ko GY, Sung KB, Yoon HK, Kim JH, Song HY, Seo TS, et al. Endovascular treatment of hepatic venous outflow obstruction after living-donor liver transplantation. *J Vasc Interv Radiol* 2002;**13**:591-9.
85. Ko GY, Sung KB, Lee S, Yoon HK, Kim KR, Kim KM, et al. Stent placement for the treatment of portal vein stenosis or occlusion in pediatric liver transplant recipients. *J Vasc Interv Radiol* 2007;**18**:1215-21.
86. Liang HL, Yang CF, Pan HB, Chen CK, Chang JM. Percutaneous transsplenic catheterization of the portal venous system. *Acta Radiol* 1997;**38**:292-5.
87. Tuite DJ, Rehman J, Davies MH, Patel JV, Nicholson AA, Kessel DO. Percutaneous transsplenic access in the management of bleeding varices from chronic portal vein thrombosis. *J Vasc Interv Radiol JVIR* 2007;**18**:1571-5.
88. Orons PD, Hari AK, Zajko AB, Marsh JW. Thrombolysis and endovascular stent placement for inferior vena caval thrombosis in a liver transplant recipient. *Transplantation* 1997;**64**:1357-61.
89. Sze DY, Semba CP, Razavi MK, Kee ST, Dake MD. Endovascular treatment of hepatic venous outflow obstruction after piggyback technique liver transplantation. *Transplantation* 1999;**68**:446-9.
90. Koo BY, Yu HC, So MC, Jin GY, Kwak HS, Cho BH. Endovascular management of early portal vein thrombosis caused by coronary vein steal after liver transplantation and its outcome. *Clin Transplant* 2008;**22**:668-71.
91. Alonso L, Alvite M, Seoane M, Berdeal M, Pérez J, Marini M, et al. Interventional radiologist treatment for inferior vena caval thrombosis in a liver transplant recipient. *Transplant Proc* 2009;**41**:1060-1.
92. Nosaka S, Isobe Y, Kasahara M, Miyazaki O, Sakamoto S, Uchida H, et al. Recanalization of post-transplant late-onset long segmental portal vein thrombosis with bidirectional transhepatic and transmesenteric approach. *Pediatr Transplant* 2013;**17**:E71-75.
93. Piscaglia F, Cecilioni L, Gaiani S, Rossi C, Losinno F, Cescon M, et al. Use of perfusional angiosonography in liver transplantation and conservative management of post-transplant intra-hepatic pseudo-aneurysm. *Transpl Int* 2004;**17**:634-8.
94. Nagaraja R, Govindasamy M, Varma V, Yadav A, Mehta N, Kumaran V, et al. Hepatic artery pseudoaneurysms: a single-center experience. *Ann Vasc Surg* 2013;**27**:743-9.
95. Ginat DT, Saad WEA, Waldman DL, Davies MG. Stent-graft placement for management of iatrogenic hepatic artery branch pseudoaneurysm after liver transplantation. *Vasc Endovascular Surg* 2009;**43**:513-7.
96. Saad WEA, Dasgupta N, Lippert AJ, Turba UC, Davies MG, Kumer S, et al. Extrahepatic pseudoaneurysms and ruptures of the hepatic artery in liver transplant recipients: endovascular management and a new iatrogenic etiology. *Cardiovasc Intervent Radiol* 2013;**36**:118-27.