





Ascertaining the Effects of Grass and Leaf Meals on the Gut Health and Blood Indices of Broiler Chickens – A Systematic Review

Yee Lyn Ong¹ , Eric Lim Teik Chung^{1,2*} , Nazri Nayan^{1,2} , Ngai Paing Tan³ , Faez Firdaus Abdullah Jesse⁴  & Awis Qurni Sazili^{1,2} 

¹ Institute of Tropical Agriculture and Food Security, Universiti Putra Malaysia, Selangor, Malaysia

² Department of Animal Science, Faculty of Agriculture, Universiti Putra Malaysia, Selangor, Malaysia

³ Department of Land Management, Faculty of Agriculture, Universiti Putra Malaysia, Selangor, Malaysia

⁴ Department of Veterinary Clinical Studies, Faculty of Veterinary Medicine, Universiti Putra Malaysia, Selangor, Malaysia

Poultry Science Journal 2024, 12(1): 1-17

Abstract

As the broiler chicken industry continues to develop, antibiotic growth promoters in poultry feed are being phased out because of increasing country restrictions and consumer concerns about food safety. As a result of these bans and efforts to prevent antimicrobial resistance, research into antibiotic alternatives is accelerated to preserve or improve broilers' production performance. Due to the presence of beneficial compounds like tannins, saponins, flavonoids, and various others found in grass and leaf meals, they possess the potential to substitute antibiotics. This is because secondary metabolites in plant-derived phytobiotics have useful pharmacological qualities that may benefit broilers' overall production and health. Previous studies, including grass or leaf meals at a dosage of 0.025-20%, have shown several positive effects on gut histomorphology, gut microflora, and blood biochemistry. For instance, lipid profile and liver functions of broiler chickens improved through hypo-cholesterolaemic and hepatoprotective functions of phytocompounds. However, there are also contradictory data and a lack of information on the effect of these plant-based meals on broilers' blood biomarkers, such as acute phase proteins and heat shock proteins. Therefore, this review provides insight into the potential of grass or leaf meals and their effects on gut health, blood biochemistry, and biomarkers of broiler chickens.

Keywords

Gut microflora
Phytocompounds
Heat shock proteins
Gut histomorphology

Corresponding author

Eric Lim Teik Chung
ericlim@upm.edu.my

Article history

Received: September 05, 2023

Revised: December 12, 2023

Accepted: January 07, 2024

Introduction

Antibiotics usage in the poultry industry

Presently, the broiler industry is the most intensive area of animal production. Through the years, the broiler industry has progressed from a backyard subsistence level to a highly commercialized and efficient production system with 'integrators' who manage all activities. These activities which include feed manufacturing, breeding, hatchery, broiler farming and meat processing are all integrated into a system known as the vertical integration system (Ferlito, 2020). During this transition to highly commercialized systems, antibiotic growth promoters (AGP) have been incorporated in birds' feed to enhance broiler traits such as daily weight gains and

feed efficiency since the favorable benefits satisfy the demands of meat production (Hao *et al.*, 2014; Alghirani *et al.*, 2021a). This antibiotic use has been linked to higher rates of antimicrobial resistance (AMR) among bacteria isolated from animals, demonstrating that antibiotic resistance genes may be transmitted to human microbiota via the food chain (Dibner and Richards, 2005). Consequences of AMR may range from prolonged illness and side effects due to the use of alternative treatments to death following complete treatment failure (Hughes and Heritage, 2004). For that reason, improved control measures, public health interventions, and a monitoring system should be implemented to minimize the spread of antibiotic resistance in local chicken farms. More

Please cite this article as Yee Lyn Ong, Eric Lim Teik Chung, Nazri Nayan, Ngai Paing Tan, Faez Firdaus Abdullah Jesse & Awis Qurni Sazili. 2024. Ascertaining the Effects of Grass and Leaf Meals on the Gut Health and Blood Indices of Broiler Chickens – A Systematic Review. *Poult. Sci. J.* 12(1): 1-17.

© 2024 PSJ. All Rights Reserved

importantly, alternatives to antibiotics and their effects on broilers should be researched further to be effectively utilized in the poultry sector.

Grasses and leaves as phytobiotic in poultry

Because of prohibitions on the use of AGP in meat production in various countries, phytobiotics as an alternative has gained traction in the past years (EP, 2003; FAO, 2004). With increased knowledge and consumer awareness, intensified rejection of synthetic antibiotics makes way for consumers to accept the use of phytobiotic in animal feed since humans have taken herbal remedies for ages (Surai, 2014). Common natural feed additives that could be used as an antibiotic alternative are phytochemical groups which include essential oils (Reis *et al.*, 2018; Coles *et al.*, 2023), enzymes (Rizwanuddin *et al.*, 2023), prebiotics and probiotics (Fathima *et al.*, 2023), herbal extracts (Galli *et al.*, 2020; Erwan *et al.*, 2021 a,b) and others (Reis *et al.*, 2019; Reda *et al.*, 2020; Bajagai *et al.*, 2022). Another form of phytobiotics such as grass or leaf meals, could also replace AGP as they contain a variety of bioactive molecules or phytochemicals such as flavonoids, saponins, tannins, alkaloids, and many more. These compounds are responsible for their antioxidant and antibacterial activities (Jimenez-Garcia *et al.*, 2018; Kuralkar and Kuralkar, 2021; Chung *et al.*, 2018; Ojo *et al.*, 2022). Certain phytochemicals such as tannins have a reputation for being an anti-nutritional factor when present in high doses, but supplementing broiler chicks with tannin diets at low doses has been shown to improve and promote immunological competence, intestinal microbial ecology, and gut health (Huang *et al.*, 2018). Similarly, multiple studies have demonstrated that appropriate levels of saponins in poultry diets may produce note-worthy results despite reports of affecting palatability due to bitter taste (Alghirani *et al.*, 2021b; Alghirani *et al.*, 2023). As a result, they might be utilized to enhance broiler diets to produce a profitable and sustainable production cycle, although additional research is needed. Therefore, this review aims to discuss the application of grass or leaf meals as a feed additive in poultry diets and the effects on gut histomorphology, gut microflora, serum lipid profile, blood biomarkers and biochemistry of broilers. This systematic research was finalized according to the guidelines of "Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement" (PRISMA). A systematic literature search was conducted via the electronic database of Scopus, where articles were identified using keywords related to production. Keywords included were 'broiler', 'grass meal', 'leaf meal', 'gut histomorphology', 'gut microflora', 'lipid profile', 'liver functions', 'acute phase proteins', and 'heat shock proteins'. Articles were then refined by restricting the publication year

to 2012 to 2023, a range of 11 years, and restricted to English-only publications. Only full-text papers that featured at least one of the gut health or blood-related features were chosen to ensure that the information acquired was sufficient. A total of 31 full-text papers were reviewed for eligibility, and nine were eliminated for failing to match the appropriate criteria, resulting in 22 articles being included in this review.

PRISMA results

A comprehensive search of the Scopus database yielded 31 papers. There are no duplicates since they were removed during the article scanning process. Publications were filtered for irrelevant components before being examined for eligibility. A total of 22 articles were included in the information analysis. The publishing dates were restricted to 11 years, from 2012 to 2023, with the first in 2012 and the most recent in 2023. The year with the most articles published was 2020, followed by four in 2021 and 2022. Figure 1 shows that Asia accounted for the majority of the studies (55%), followed by Africa (32%), Oceania (9%), and Europe (4%). Furthermore, the plants researched in the selected studies can be classified as grasses, legumes, or non-legumes. Only one of the papers (4%) studied supplemental grass feed to broilers. Non-legume papers comprised 73% of those evaluated, whereas legumes comprised 23% (Figure 2). This review will go through the usage of grass or leaf meals as a feed supplement in broiler diets, as well as the effects on gut histomorphology, gut microbiota, serum lipid profile, blood biomarkers, and biochemistry.

The effect of grass or leaf meals on gut histomorphology of broilers

One of the hypothesized ways phytobiotic may stimulate development is through modifications in the gastrointestinal system (Valenzuela-Grijalva *et al.*, 2017). Though the method by which phytochemicals alter gut structure is not fully known, there have been multiple theories on how changes in intestinal morphology affect chicken production performance. A greater villi height: crypt depth ratio (VH:CD) may indicate that the broiler's small intestine has superior digesting and absorption capabilities due to a larger absorptive surface area, allowing for increased nutrient uptake. Besides enhancing VH: CD, changes in gut pH, epithelial cell proliferation, acid secretion, gastrin synthesis, and dietary modifications may alter gut histomorphology and health by decreasing microbial pressure, stabilizing bowel health, and preventing intestinal illnesses (Chrubasik *et al.*, 2005; Pohl *et al.*, 2012). The increase in villus height and VH: CD was also found to have a direct correlation with the increase in epithelial cell turnover, thus activating cell mitosis

(Khan et al., 2017). Alghirani et al. (2021) further explained that the improved growth performance due to decreased turnover rate in intestinal mucosal cells, results in a shorter crypt depth and lower maintenance needs, thus allowing more energy to be dedicated to development. In addition to changes in gut morphology, a rise in the production of enzymes such as proteases and amylases improves growth performance in the gastrointestinal tract by increasing

absorptive cell development (Jamroz et al., 2006). Several studies have found that incorporating phytobiotic in poultry feed can improve the morphological features of a chicken's intestine, hence enhancing its production ability and promoting food safety (Jamroz et al., 2006; Ferdous et al., 2021; Gilani et al., 2021). As a result, including phytobiotic in chicken feed has been proven to increase VH: CD ratio and promote digestive enzyme secretions.




Figure 1. Number of included publications on grass or leaf meals as a feed additive in broiler diets and their effects on gut health and blood indices based on the world region.




Figure 2. Number of included publications on grass or leaf meals as a feed additive in broiler diets and their effects on gut health and blood indices based on plant classification.

Supplementation of 25 mg/kg of *Brachiaria decumbens* powdered meal was observed to show an increase ($P < 0.05$) in villus height while simultaneously decreasing ($P < 0.05$) crypt depth in the duodenum, jejunum, and ileum of Cobb broilers on at both starter and finisher phase (Alghirani et al., 2022). This might be explained by saponins' antioxidant properties, which may have changed intestinal morphology, enhancing lumen conditions such as number of pathogens, intestinal thickness,

and mucus secretion (Tavangar et al., 2021). Antioxidant properties may also promote digestive tract growth and development by increasing villus height, which enhances absorptive surface area, enzyme secretions, and nutrition transfer (Olukosi and Dono, 2014). In the same line, the administration of 15 or 30 mL/L nanoencapsulated *Terminalia catappa* leaf extract, which is rich in various phytochemicals into broilers' drinking water did not affect the crypt depth but improved the villus height

($P < 0.01$) and thus, a higher VH:CD ratio ($P < 0.05$). Results showed that the extract capsulated or not, produced higher ($P < 0.01$) villus height in comparison to the usage of tetracycline (1521.01-1723.00 μm vs 1407.15 μm) as antibiotics. There were no apparent differences ($P > 0.05$) between the villus width and crypt depth throughout all treatments supplemented with *Terminalia catappa* leaf extract and the antibiotic group. Therefore, VH:CD ratio of nanocapsulated (9.20) or non-nanocapsulated treatments (10.61) produced better results than the control (5.68) and antibiotic (6.15) group (Hidayati *et al.*, 2022). Similar results were discovered by Saharan *et al.* (2022) which studied the influence of supplementing 15, 25 and 35g of *Psidium guajava* leaf meal (PLGM) per kg of feed found that VH:CD ratio was significantly improved ($P < 0.05$). On top of that, the incorporation of *Azolla* leaf meal (ALM) into broiler diets at 50g and 100g per kg of baseline food resulted in an increase in villi length in the duodenum ($P=0.09$). Meanwhile, in the jejunum, treatment groups given 50 g of ALM per kg of basal diet increased villi length by 29.8% compared to the control group. However, compared to the control group, villi length fell by 8% in the ileum ($P=0.03$). Because the jejunum is the primary location of nutrient absorption in chickens, an increase in the villi length of the jejunum with the addition of 50 g ALM improved the intestine's absorptive capacity, which would be beneficial to the poultry (Abdelatty *et al.*, 2021). These positive results could be explained by adding dietary phytobiotic with antibacterial action. These compounds may diminish pathogenic microbe growth and toxic chemicals. These chemicals inhibit intestinal mucous production by Goblet cells, minimizing intestinal damage and lumen repair (Hidayati *et al.*, 2022). This theory is further supported by Ogwuegbu *et al.* (2021) which studied the effects of sodium butyrate and rosemary leaf meal on the gut histology and microflora. They discovered that supplementing birds with 5.0 g of rosemary leaf meal per kg of basal diet improved ($P<0.05$) gut histological traits which include villus length, crypt depth, epithelial thickness, muscularis thickness in the duodenum, jejunum, and ileum. Rosemary leaf meal is said to have antioxidant, antimicrobial, and antifungal properties derived from bioactive compounds such as carvacrol, thymol, capsaicin, cineole, rosmannol, carnosol, and their acid forms, or flavonoids (Ahsan *et al.*, 2018). Basit *et al.* (2020b) also reported that with the supplementation of phytobiotics (4 g/kg of *Piper betle* and 8g/kg *Persicaria odorata* leaf meal) improved ($P < 0.0001$) the villus height in the jejunum and duodenum up to

8.6% similarly to the addition of tetracycline as antibiotics and 0.03g of halquinol/kg of basal diet in comparison to the negative control. For the CD, no change ($P > 0.05$) was seen in jejunum but improvements were significant ($P < 0.05$) in the duodenum and ileum of groups supplemented with *Piper betle*, *Persicaria odorata*, halquinol and antibiotics. Therefore, significant ($P < 0.05$) positive results were seen in the VH: CD ratio of the duodenum, jejunum and ileum of broilers with treatment diets in comparison to the negative control (basal diet only).

On the other hand, Sugiharto *et al.* (2020) compared the effects of feeding fermented mixture of cassava pulp and *Moringa oleifera* leaf meal (FCPMO) through studying four diet groups: corn-soybean-based feed with no additive (CONT), corn-soybean-based diet supplemented with 0.1% zinc bacitracin (BACI), diet containing 20% FCPMO (FERM), and diet containing 20% FCPMO and added with 0.1% *Bacillus subtilis* (FERB). Although FERM enhanced other criteria such as immunological responses, antioxidative state, and physiological circumstances, the findings for intestinal morphology were less than favorable. While the villi height was not significant across treatment groups ($P > 0.05$), the CD in FERM was larger ($P < 0.05$). Thus, the VH:CD ratio in FERM (4.54) was lower ($P < 0.05$) than in CONT (6.60) and FERB (6.09). The precise explanation of the lower VH:CD ratio is obscure; however, it is thought to be owing to the high fiber content in FERM, since Saadatmand *et al.* (2019) showed that dietary fiber (rice hull) would result in a lower VH: CD. This negative effect would decrease absorptive capacity, resulting in poor development performance in grill chickens.

A summary of the effects of grass or leaf meals on the gut histomorphology of chickens can be found in Table 1 below. Based on previous experiments, there were more positive or non-significant results produced with the addition of plant-based phytobiotic in the diets of broilers. Negative findings could be explained by the high fiber content which is why it is crucial for thorough studies to be conducted on grass and leaf meals. This is necessary to determine the optimum inclusion level of feed additive that will produce beneficial results in the gut morphology without compromising the growth performance of the bird. Future research might benefit from a deeper examination of the relationship between fiber content and villus height, as well as the significance of intestinal mucus and the microbial community to gain the chicken industry's trust and acceptance of phytobiotic as an antibiotic substitute.

Table 1. The different effects of various leaf meal supplementation on poultry's gut histomorphology.

Source	Inclusion levels	Effect on gut histomorphology	References
<i>Brachiaria decumbens</i> ground leaf powder	25mg/kg of basal diet without antibiotics	Increased villus height and decrease in crypt depth.	Alghirani et al. (2022)
<i>Terminalia catappa</i> leaf extract	15 or 30 mL of <i>Terminalia catappa</i> leaf extract (nanoencapsulated or not) per L of drinking water	Improved villus height. No effect in crypt depth. Higher VH: CD ratio.	Hidayati et al. (2022)
Guava (<i>Psidium guajava</i>) leaf meal	15, 25 and 35g of PLGM	Improved VH:CD.	Saharan et al. (2022)
<i>Azolla</i> leaf meal	50 g and 100 g of <i>Azolla</i> leaf meal per kg of baseline food	Improved villi length in duodenum and jejunum. Depressed villi length in the ileum.	Abdelatty et al. (2021)
Rosemary leaf meal	5.0 g of rosemary leaf meal per kg of basal diet	Enhanced villus length, crypt depth, epithelial thickness, muscularis thickness in the duodenum, jejunum, and ileum	Ogwuegbu et al. (2021)
<i>Piper betle</i> and <i>Persicaria odorata</i> leaf meal	4g/kg of <i>Piper betle</i> and 8g/kg <i>Persicaria odorata</i> leaf meal	Higher villus height in the jejunum and duodenum. Improved crypt depth of jejunum and ileum. No change in crypt depth of jejunum. Better VH:CD ratio of duodenum, jejunum and ileum.	Basit et al. (2020a)
Cassava pulp and <i>Moringa oleifera</i> .	20% fermented mixture of cassava pulp and <i>Moringa oleifera</i>	No significant differences in crypt depth. Lower VH:CD ratio	Sugiharto et al. (2020)

The effect of grass or leaf meals on gut microflora of broilers

The chicken gastrointestinal system is home to diverse microorganisms living in symbiotic partnerships, ultimately impacting the host's nutrition, metabolism, and immunity. The balance of pathogenic and non-pathogenic bacteria in the gut could disrupt the microflora equilibrium and compromise the intestinal barrier, allowing toxic compounds or pathogenic bacteria to enter the intestinal lumen (Gilani *et al.*, 2021). As such, another way to enhance the general health and performance of broiler chickens is to improve gut health conditions, particularly by lowering the potentially harmful microbe population in the gut. Previous research indicates that plant-based feed additives positively impact the gut bacteria community when incorporated into the chicken diet. This is because bioactive substances present can break down the cellular membrane and interfere with the cytoplasm ecology of infections, thus acting as phytobiotic (Mohammadi Gheisar and Kim, 2018). When phytobiotics come into contact with the microbial cell membrane, they modify permeability for H⁺ and K⁺ cations, reducing the population of enteropathogens (Gilani *et al.*, 2021). For example, because microorganisms require iron for growth, tannins' antibacterial method of action is hypothesized to limit metal ions, impair microbe cell membrane permeability through complexes, and modify the morphology of cell walls (Suresh *et al.*, 2017; Basit *et al.*, 2020b). The tannin process of iron chelation reduces metal ion availability, limiting bacteria growth except for *Lactobacillus*, which does not require iron for development (Yilmaz and Li, 2018). Whereas saponins form interactions with membrane sterols, causing membrane damage, while alkaloids block topoisomerase and disrupt DNA synthesis (Suresh *et al.*, 2017). Moreover, with the incorporation of plant-origin active substances in the broiler diet, increased mucus production on the jejunum wall supports villi-related protective characteristics of phytochemicals, which might explain the lowered quantity of detrimental microbiomes to the epithelial wall (Jamroz *et al.*, 2006). These phytobiotic qualities will result in bacteriostatic and bacteriocidal actions, reducing pathogen populations and increasing the production and numbers of beneficial bacteria in the broiler's gut.

Alghirani *et al.* (2022) discovered that supplementing Ross308 broiler chicks with *B. decumbens* ground leaf meal suppressed the development of *Enterococcus faecalis* during the starter phase, similar to the antibiotic-fed group. The presence of saponins, tannins, flavonoids, and alkaloids, all of which regulate the cecal microbiota and have antibacterial properties, implies that

B. decumbens has antimicrobial properties that protect against harmful bacteria such as *E. faecalis* (Zdunczyk *et al.*, 2010; Viveros *et al.*, 2011; Low, 2015; Hidayat *et al.*, 2021). Furthermore, compared to the other treatments, the group that received 25 mg/kg of *B. decumbens* had the highest standard plate count and coliform count throughout the experiment. These findings could be attributed to increased fecal coliforms and/or decreased lactobacilli, anaerobes, aerobes, and enterococci groups contributing to increased feed utilization. In another experiment, Mandal *et al.* (2014) discovered that adding MOLM, which also contains a wide variety of phytochemicals to day-old broiler chicks reduced microbial load compared to the other groups, including the control and antibiotic-supplemented groups. Incorporation of ALM into broiler diets was also demonstrated to linearly enhance *Lactobacilli* species and reduce *Bacilli* as ALM inclusion levels rose, but decreased *Enterobacteriaceae* independent of ALM inclusion level. Abdelatty *et al.* (2021) could not determine if the alteration in intestinal mucus was driven by the change in microbiota with the addition of ALM or vice versa since ALM contains a large amount of silica, which might have affected the mucous contents. *Lactobacillus* count in the ileum and caecum was also higher ($P < 0.05$) in birds treated with rosemary leaf meal regardless of inclusion level in comparison to those fed with the negative control diet. Conversely, those fed the negative control diet had the highest ($P < 0.05$) count of *Escherichia coli* and *Salmonella*. Beneficial bacteria in the gut, such as *Lactobacillus*, contribute to the production of antibiotic agents, bile salt hydrolase chemicals, and the preservation of gut integrity via their probiotic properties (Ogwuegbu *et al.*, 2021). According to Saeed *et al.* (2019), increased *Lactobacillus* and decreased *E. coli* and *Salmonella* improve the intestinal microflora balance by reducing the number of hazardous microbes. This promotes the development of intestinal absorptive cells, which in turn encourages the growth of birds. Furthermore, it was reported that the supplementation of phytobiotics (*P. betle* and *P. odorata*) significantly increased ($P < 0.001$) the *Lactobacillus* population count compared to other groups (halquinol, antibiotics and negative control). Phytobiotics treated group also significantly decreased ($P < 0.05$) population count of *E. coli*, *Staphylococcus aureus*, *Clostridium*, and *Salmonella* (Basit *et al.*, 2020b). Thus, the inclusion of phytobiotic might result in a healthy gut microbiota population since they contain beneficial secondary metabolites such as flavonoids and phenolic compounds that favorably influence the microbiota population (Liaqat *et al.*, 2016; Rahman and Yang, 2018; Mustafa, 2019).

Table 2: The different effects of various leaf meal supplementation on poultry's gut microbial population.

Source	Inclusion levels	Effect on gut microbial	References
<i>Brachitaria decumbens</i> ground leaf powder	25 mg/kg of basal diet without antibiotics	Inhibited the development of <i>E. faecalis</i> during the starter phase.	Alghirani et al. (2022)
<i>Moringa oleifera</i> leaf meal	5g, 10g, 15g and 20g of <i>Moringa oleifera</i> leaf meal	Reduced microbial load and coliform count in all treatment levels.	Mandal et al. (2014)
<i>Azolla</i> leaf meal	50g and 100g of <i>Azolla</i> leaf meal per kg of baseline food	Increased <i>Lactobacilli</i> sp. Reduced <i>Bacilli</i> and <i>Enterobacteriaceae</i> .	Abdelatty et al. (2021)
Rosemary leaf meal	2.5g and 5.0g of rosemary leaf meal	Higher <i>Lactobacillus</i> count in the ileum and caecum and lower <i>E. coli</i> and <i>Salmonella</i> count.	Ogwuegbu et al. (2021)
Guava (<i>Psidium guajava</i>) leaf meal	15, 25 and 35g of PLGM	Increased ($P < 0.001$) <i>Bifidobacteria</i> and <i>Lactobacilli</i> Reduced <i>E. coli</i> and <i>Clostridia</i> .	Saharan et al. (2022)
Copra meal or cassava leaf meal	100g and 200g/kg of feed	No effect on <i>E. coli</i> population count.	Diarra and Anand (2020)
Copra meal, palm kernel meal and cassava leaf meal	100g and 200g/kg of feed	No effect on <i>E. coli</i> population count.	Diarra et al. (2023)

Conversely, adding copra meal or cassava leaf meal to broiler diets had no effect ($P > 0.05$) on *E. coli* counts, even though the fermentation of dietary fibre in the ceca is thought to suppress pathogenic bacteria by lowering gut pH through the production of short-chain fatty acids (Khan and Iqbal, 2016; Jha and Mukku, 2019; Diarra and Anand, 2020). Diarra *et al.* (2023) found similar results ($P > 0.05$) on the *E. coli* count with the supplementation of copra meal, palm kernel meal and cassava leaf meal. They deduced that the supplementation doses did not provide substantial fibre content for relevant bacterial fermentation in the ceca. This theory is supported by Saharan *et al.* (2022), who found that PGLM supplementation significantly reduced ($P < 0.001$) *E. coli*, *Clostridia* and increased ($P < 0.001$) *Bifidobacteria* and *Lactobacilli* counts in the ceca as the intake of crude fibre was significantly ($P < 0.05$) higher with the increase of PGLM supplementation rate.

Despite some insignificant results with adding these leaf meals on the gut microbial population, it can also be said that including plant-based products has no deleterious effects on the gut microbial population in broiler birds. Table 2 elaborates on the different effects of various leaf meal supplementation on poultry's gut microbial population.

The effect of grass or leaf meals on serum lipid profile, liver functions, and blood biomarkers of broilers

Tropical environments substantially influence animal well-being and impact animal output globally by encouraging animals to restrict heat generation by lowering feed intake. This leads to a detrimental impact on growth performance (Hansen, 2009). Besides, environmental, pathogenic, and nutritional issues, for example, can induce a state of stress, prompting a series of behavioral and physiological reactions that would reduce poultry performance (Tamzil *et al.*, 2013). Because of their antioxidant activity, phytobiotics could be utilized during heat stress and other stressor conditions to influence poultry serum biochemistry by lowering serum concentrations of cholesterol, triglycerides (TG), and low-density lipoproteins (LDL) while increasing high-density lipoproteins (HDL) levels in broilers (Gilani *et al.*, 2018). This may be due to the distinct features of phytobiotic, which have been linked to hypocholesterolaemic effects as volatile oils found in plants can impede the function of 3-hydroxy-3-methylglutaryl-coenzyme, a reductase (HMGCoA reductase) liver enzyme that controls cholesterol production, thus lowering blood cholesterol levels (Fujioka *et al.*, 2003). On top of that, most therapeutic plants have been proven to be hepatotoxic, either owing to phytochemical ingredients or dose (Addy *et al.*, 2013) and blood

parameters provide valuable insight for the health status of broiler birds (Rehman *et al.*, 2017). Plant secondary metabolites have also been examined for their immunomodulatory effects, which are assumed to be linked to the stimulation of heat shock proteins, which improve protein translation efficiency (Suresh *et al.*, 2017). Furthermore, the study on hepatic function indicators such as aspartate aminotransferase (AST), alanine transaminase (ALT), and alkaline phosphatase (ALP) levels would offer information on liver function and the occurrence of liver damage caused by toxins (Muazu and Aliyu-Paiko, 2020).

Additionally, blood biomarkers play a vital role in the poultry industry to estimate the broilers' welfare and health status (Attia *et al.*, 2011). Acute phase proteins (APPs) are thought to have diagnostic and prognostic potential due to the link between blood biomarker levels and host response to inflammation and/or infection (Ceron *et al.*, 2005). Whereas heat shock proteins (HSPs) are synthesized and accumulated as a result of cellular stress reaction, therefore may be used to identify stressful periods for the organism (Balakrishnan *et al.*, 2023). Both biomarkers are significant components and indications of the immunological response and well-being of broiler chickens, notably in infections and non-infectious illnesses, as well as in nutrition studies. APPs are a class of blood proteins primarily synthesized by hepatocytes, though they can be produced in other tissues and organs, whose concentrations either increase (positive APP) or decrease (negative APP) in order to maintain homeostasis balance as part of the acute phase response (APR). APR is a part of the early-defense mechanism that serves to trigger the acquired immune response (Janeway *et al.*, 2001). Therefore, in veterinary clinical pathology, APPs are helpful indicators for identifying infection, inflammation, trauma, and overall animal health (Gruys *et al.*, 2005). On the other hand, heat shock proteins (HSPs) are a group of proteins that are produced in response to various stressors such as physical, chemical, or biological stimuli such as heat and pathogenic infection (Ganter *et al.*, 2006; Staib *et al.*, 2007). Depending on their molecular weight, HSPs can be classified into several families (Basu *et al.*, 2002). While each HSP family has a unique purpose and function inside cells, the key role of HSPs is to protect and repair cells by promoting proper protein folding, which can occur under stressful situations (Sahin *et al.*, 2009). Research shows a substantial link between oxidation and HSP70 production (Ming *et al.*, 2010). Therefore, understanding blood biomarkers and biochemistry functions and reactions can aid in the development of better diagnostic procedures, treatment approaches, and diet formulations in the broiler industry.

Serum cholesterol reduced ($P < 0.05$) in birds supplemented with 10 and 15 g of mucuna leaf meal compared to birds in other treatments. Additionally, the supplementation of mucuna leaf meal had no effects ($P > 0.05$) on the AST levels of broilers across all treatments, signifying that there were no negative impacts on the liver's normal physiological and anatomical function (Oloruntola et al. 2022). Widiastuti et al. (2021) found that the addition of acidified papaya leaf and seed meal (APLS) at 10, 25, and 50 g per kg of broiler diet linearly increased ($P < 0.05$) HDL to LDL ratio and linearly decreased ($P=0.06$) cholesterol to HDL ratio with higher inclusion levels of APLS. These favorable results would prevent unwanted physiological conditions in broilers such as cardiovascular problems (Bueno et al., 2017). Similarly, broilers supplemented with 10, 20 and 30 g of *Vernonia amygdalina* leaf meal per kg diet had lower total cholesterol (107.3-118.7 mg/dL) and LDL values (93.9-113.7 mg/dL) than those in the control group (139.3 mg/dL and 122.3 mg/dL respectively) suggesting the reduction in triglyceride synthesis and improved redistribution of cholesterol between lipoprotein molecules (Tokofai et al., 2020).

Another experiment that included *Moringa oleifera* leaf meal at 0, 25, 50, and 100 g/kg DM of diet of broiler birds discovered that the diet had a significant effect ($P < 0.05$) on AST, ALT, and ALP. Higher levels of MOLM resulted in lower AST and ALP levels, indicating no toxic effects in the liver (Sebola and Mokoboki., 2019). In the same way, dietary supplementation of *Persicaria odorata* leaf meal at 2, 4, and 8g/kg diet showed a linear decrease ($P < 0.05$) of AST, ALT, and ALP on day 21 and linear decrease of AST, ALT, cholesterol, and triglycerides ($P < 0.05$) on day 42 with higher inclusion levels of *P. odorata* leaf meal. Reduced ALT and AST levels suggest hepatoprotective action, which is thought to be associated with flavonoids and secondary metabolites in the leaf meal. Likewise, the improved serum levels of triglycerides and cholesterol are thought to be from the phytobiotic effect of *P. odorata* leaf meal (Basit et al., 2020a).

Compared to the negative control group, broiler chicks given 1.25, 2.5, and 5.0g of *Azadirachta indica* leaf meal per kg of feed and antibiotics exhibited reduced ($P < 0.05$) ALP values. Furthermore, when leaf meal levels increased, blood and tissue cholesterol levels reduced dramatically ($P < 0.05$), suggesting a decrease in lipid mobilization (Ansari et al., 2012). Dietary inclusion of 0.1 and 0.2g/kg of *Morinda lucida* leaf meal were found to have lower ($P < 0.05$) levels of AST and ALT, with or without routine medication included. However, all treatments did not affect serum cholesterol and triglyceride results ($P > 0.05$). The authors ascribed the reduced AST levels to the presence of tannins and saponins, which have hepatoprotective properties (Lala et al., 2018). On the other hand, the addition of *Accacia angustissima* leaf meal at 50 and 100g/kg

diet had no effect ($P > 0.05$) on the serum AST, ALP, TG, HDL, and LDL of broiler birds (Ncube et al., 2018). But at 4 and 6 weeks, ALT levels were elevated with increased levels of *A. angustissima* leaf meal after the starter phase. As ALT is the most sensitive hepatocyte damage marker, this may indicate liver damage when the liver attempts to metabolize the antinutritional substances (Giannini et al. 2005; Ojo et al. 2013; Hassan et al. 2016).

An experiment by Song et al. (2017) which studied the effects of enzymatically treated *Artemisia annua* on the intestinal inflammatory response of heat-stressed Arbor Acres broilers aged 21 days' old found that supplementing *A.annua* at 1g/kg of feed significantly decreased ($P < 0.05$) the mRNA expression of HSP70. *A. annua* is an annual weedy herb that has been demonstrated to boost antioxidant capacity and immunological function due to its high antioxidant content, which includes flavonoids and phenolics (Brisibe et al., 2009; Cherian et al. 2013; Gholamrezaie Sani et al., 2013). The presence of higher levels of HSP70 mRNA in the heat stress model group in the current study shows that heat stress was induced, and the results indicate that supplementation of enzymatically treated *A. annua* could be linked to the modification of pro-inflammatory cytokine production. The research also examined plasma diamine oxidase (DAO), which is prominent in the top section of the intestinal mucosa and revealed a significant increase ($P < 0.05$) due to heat stress (Li et al., 2015). Elevated DAO levels suggest necrosis of the intestinal mucosal cells, which ultimately impair intestinal function and permeability (Li et al., 2002). More importantly, enzymatically treated *A. annua* supplementation was observed to significantly ($P < 0.05$) lower plasma DAO levels, indicating that enzymatically treated *A. annua* may enhance intestinal barrier function. These findings could be supported by another study that looked at the effects of pure rosemary extracts on heat stress in broilers and discovered that rosemary created a higher antioxidative condition (Tang et al., 2018). Antioxidant activity of rosemary is due to the leaves containing antioxidant compounds such as carnosol, carnosic acid, rosmaridiphenol, rosmanol, isorosmanol, epirosmanol, rosmariquinone, and rosmarinic acid which could be comparable to butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA), which are synthetic antioxidants (Zhang et al., 2010). The results showed that HSP70 levels in broiler birds' hearts were raised ($P < 0.0001$) before and during heat stress by introducing purified rosemary extracts at a rate of 3%, providing a protective effect to the chicken heart during heat stress. According to the authors, the addition of rosemary stimulated the development of HSP70, which adheres to and inhibits apoptotic proteins while also repairing unfolded or misfolded proteins in response to diverse environmental stresses.

Table 3: The different effects of various leaf or grass meal supplementation on serum lipid profile, liver functions, and blood biomarkers of poultry.

Source	Inclusion levels	Effect on serum lipid profile, blood biomarkers and biochemistry	References
Mucuna leaf meal	10 and 15g per kg diet	Reduced serum cholesterol. No effect on AST.	Oloruntola <i>et al.</i> (2022)
Acidified papaya leaf and seed meal	10, 25, and 50g per kg diet	Increased HDL: LDL with higher inclusion levels. Linear decrease of cholesterol: HDL with higher inclusion levels.	Widiastuti <i>et al.</i> (2021)
<i>Vernonia amygdalina</i> leaf meal	10, 20 and 30g per kg diet	Lower total cholesterol and LDL	Tokofai <i>et al.</i> (2020)
<i>Moringa oleifera</i> leaf meal	25,50, and 100g/kg DM of diet	Higher inclusion levels resulted in lower AST and ALP levels.	Sebola and Mokoboki (2019)
<i>Persicaria odorata</i> leaf meal	2,4, and 8g/kg diet	Decrease of AST, ALT, and ALP with higher inclusion levels.	Basit <i>et al.</i> (2020a)
<i>Azadirachta indica</i> leaf meal	1.25, 2.5, and 5.0g/kg diet	Reduced ALP, blood and tissue cholesterol.	Ansari <i>et al.</i> (2012)
<i>Morinda lucida</i> leaf meal	0.1 and 0.2g/kg diet	Lower AST and ALT. No effect on serum cholesterol and triglyceride.	Lala <i>et al.</i> (2018)
<i>Accacia angustissima</i> leaf meal	50 and 100g/kg diet	No effect on serum AST, ALP, TC, HDL, and LDL.	Neube <i>et al.</i> (2018)
Enzymatically treated <i>Artemisia annua</i>	1g/kg of feed	Lowered mRNA expression of HSP70.	Song <i>et al.</i> (2017)
Pure rosemary extracts	3% rosemary nanoemulsion liquid	Higher HSP70 levels in the heart.	Tang <i>et al.</i> (2018)
Powdered <i>Yucca schidigera</i> saponins	100 mg/kg of feed	Lowered levels of cholesterol, TG, LDL, SAA, AGP, corticosterone and serum HSP70. Higher levels of HDL.	Alghirani <i>et al.</i> (2023)

While there were no known researches conducted that studied the effect of grass or leaf meals on the blood biomarkers such as APP, corticosterone and HSP, Alghirani *et al.* (2023) found that supplementing powdered *Yucca schidigera* saponins to broiler chickens at 25, 50, 75, and 100mg per kg of basal feed resulted in significant differences ($P < 0.05$) in serum lipid profile, liver function, acute phase protein, hormone, and heat shock protein analyses. Compared to the other treatment groups, broilers given 100mg/kg of *Y. schidigera* saponins had the lowest cholesterol, TG, and LDL, as well as the greatest concentration of HDL. On top of that, 100 mg/kg *Y. schidigera* saponins supplemented broilers revealed the lowest levels of serum amyloid A (SAA), alpha-1-acid glycoprotein (AGP) and corticosterone which are stress biomarkers, indicating a reduced inflammatory reaction and stress response, possibly leading to enhanced growth performance. Conversely, no significant variations in ceruloplasmin (CP) concentrations were seen between treatments ($P > 0.05$). Saponins' cholesterol-lowering impact, which inhibits pancreatic cholesterol esterase bile acid binding, and decreases cholesterol solubility in micelles, potentially delaying cholesterol absorption in the gut, was linked to the positive effects of those lipid profiles in this experiment (Ngamukote *et al.*, 2011). SAA concentration, which is the most sensitive protein of APP was found to be lower in the group supplemented with 100 mg/kg of *Y. schidigera*. Similarly, AGP which is a major and more sensitive APP in poultry (Nazifi *et al.*, 2010) also showed a decrease in concentration on day 42. The reduced APP levels could be explained by saponins' anti-inflammatory and anti-microbial effects. Additionally, corticosterone levels were lower at 100 mg/kg of *Y. schidigera* powder because saponins have a hypocholesterolemic effect, and since serum cholesterol is a precursor to serum corticosterone, lower serum corticosterone levels could be linked to lower serum cholesterol levels. Since this experiment was conducted in an open-sided house in a tropical environment, serum HSP70 was measured and found to be lower in the highest concentration of *Y. schidigera* supplementation (100 mg/kg) as saponins present could scavenge radicals, metal chelate, and synergize with other antioxidants, thus lowering HSP70 levels.

References

Abdelatty AM, Mandouh MI, Mohamed SA, Busato S, Badr OAM, Bionaz M, Elolimy AA, Moustafa MMA, Farid OAA & Al-Mokaddem AK. 2021. *Azolla* leaf meal at 5% of the diet improves growth performance, intestinal morphology and P70S6K1 activation, and affects cecal microbiota

Table 3 summarizes the effect of leaf meals on serum lipid profile, blood biomarkers and biochemistry of broilers. While a majority of the previous experiments indicated positive or neutral effects with the inclusion of leaf meals, further studies should be carried out to determine the phytocompound or nutritional compositions responsible for these effects as well as the dosage required for that could potentially allow the replacement of antibiotics in feed. Moreover, no previous research studied the effects of grass or leaf meals on blood biomarkers such as APP, corticosterone and HSP in broiler birds. Therefore, with the plethora of available plants, more viable options should be discovered and their effects studied to ensure a continuous supply should phytobiotics be adopted in the commercial poultry industry.

Summary

To summarize, based on the information provided and the conclusions of prior research, adequate inclusion levels of grass or leaf meals may positively influence overall gut health, serum lipid profile, blood biomarkers, and biochemistry. However, because phytocompound profiles and levels differ amongst plant species, additional study is needed to find the optimal amount in broiler feed to utilize them effectively. Furthermore, additional aspects like management, harvesting age, preparation procedures, and so on will all impact the phytocompound profiles of each plant meal and must be addressed. Additional research is also required to investigate the economic benefits of employing phytobiotics in broiler production, and the effects on end products such as meat quality. More data and information from studies will aid in filling research gaps on the use of grass and leaf meals and their effects on broilers' physiology. As a result, both the broiler industry and consumers would profit from using phytobiotics as a viable, natural alternative to synthetic antibiotics, thereby improving food safety.

Acknowledgement

The authors would like to thank the Ministry of Higher Education, Malaysia, for providing funds for some of the studies conducted in this review. The project was funded by the Fundamental Research Grant Scheme (FRGS), (Grant no: FRGS/1/2020/WAB04/UPM/02/1).

in broiler chicken. *Animals*, 15(10): 100362. DOI: 10.1016/j.animal.2021.100362
Addy BS, Owodo HT, Gyapong RN, Umeji CO & Mintah DN. 2013. Phytochemical screening and antimicrobial study on the leaves of *Morinda lucida* (Rubiaceae), *Journal of Natural Sciences*, 3(14): 131–136.

- Ahsan U, Kuter E, Raza I, Köksal BH, Cengiz Ö, Yıldız M, Kızanlık PK, Kaya M, Tatlı O, & Sevim O. 2018. Dietary supplementation of different levels of phytogetic feed additive in broiler diets: the dynamics of growth performance, caecal microbiota, and intestinal morphometry. *Brazilian Journal of Poultry Science*, 20(4): 737-746. DOI: 10.1590/1806-9061-2017-0698
- Alghirani, MM, Chung, ELT, Jesse, FFA, Sazili, AQ, & Loh, TC. 2021. Could Phytobiotics replace Antibiotics as Feed Additives to Stimulate Production Performance and Health Status in Poultry? An Overview. *Journal of Advanced Veterinary Research*, 11(4), 254-265.
- Alghirani MM, Chung ELT, Sabri DSM, Tahir MNJM, Kassim NA, Kamalludin MH, Nayan N, Jesse FFA, Sazili AQ & Loh TC. 2021b. Can *Yucca schidigera* be used to enhance the growth performance, nutrient digestibility, gut histomorphology, cecal microflora, carcass characteristic, and meat quality of commercial broilers raised under tropical conditions? *Animals*, 11(8): 2276. DOI: 10.3390/ani11082276
- Alghirani MM, Chung ELT, Kassim NA, Ong YL, Jesse FFA, Sazili AQ & Loh TC. 2023. Blood biochemistry and stress biomarkers of broiler chickens supplemented with different levels of *Yucca schidigera* saponins reared under tropical conditions, *Veterinary Intergrative Sciences*, 21(1): 1-15. DOI: 10.12982/VIS.2023.001
- Alghirani MM, Chung ELT, Kassim NA, Ong YL, Jesse FFA, Sazili AQ & Loh TC. 2022. Effect of *Brachiaria decumbens* as a novel supplementation on the production performance of broiler chickens. *Tropical Animal and Health Production*, 54: 386.
- Ansari J, Khan S, Ul-Haq A & Yousaf M. 2012. Effect of the levels of *Azadirachta indica* dried leaf meal as phytogetic feed additive on the growth performance and haemato-biochemical parameters in broiler chicks. *Journal of Applied Animal Research*, 40(4): 336-345. DOI: 10.1080/09712119.2012.692329
- Attia YA, Hassan RA, Tag El-Din AE & Abou-Shehema BM. 2011. Effect of ascorbic acid or increasing metabolizable energy level with or without supplementation of some essential amino acids on productive and physiological traits of slow-growing chicks exposed to chronic heat stress. *Journal of Animal Physiology and Animal Nutrition*, 95(6): 744-755. DOI: 10.1111/j.1439-0396.2010.01104.x
- Bajagai YS, Petranyi F, J Yu S, Lobo E, Batacan Jr R, Kayal A, Horyanto D, Ren X, Whitton MM & Stanley D. 2022. Phytogetic supplement containing menthol, carvacrol and carvone ameliorates gut microbiota and production performance of commercial layers. *Scientific Reports*, 12(1): 11033. DOI: 10.1038/s41598-022-14925-0
- Balakrishnan KN, Ramiah SK & Zulkifli I. 2023. Heat shock protein response to stress in poultry: a review. *Animals*, 13(2): 317. DOI: 10.3390/ani13020317
- Basit MA, Arifah AK, Loh TC, Saleha AA, Salleh A, Kaka U & Idris SB. 2020a. Effects of inclusion of different doses of *Persicaria odorata* leaf meal (POLM) in broiler chicken feed on biochemical and haematological blood indicators and liver histomorphological changes. *Animals*, 10(7): 1209. DOI: 10.3390/ani10071209
- Basit MA, Kadir AA, Loh TC, Abdul Aziz S, Salleh A, Zakaria ZA & Banke Idris S. 2020b. Comparative efficacy of selected phytobiotics with halquinol and tetracycline on gut morphology, ileal digestibility, cecal microbiota composition and growth performance in broiler chickens. *Animals*, 10(11): 2150. DOI: 10.3390/Ani10112150
- Basu N, Todgham AE, Ackerman PA, Bibeau MR, Nakano K, Schulte PM & Iwama GK. 2002. Heat shock protein genes and their functional significance in fish. *Gene*, 295(2): 173-183. DOI: 10.1016/S0378-1119(02)00687-X
- Brisibe EA, Umoren UE, Brisibe F, Magalhães PM, Ferreira JF, Luthria D, Wu X & Prior RL. 2009. Nutritional characterisation and antioxidant capacity of different tissues of *Artemisia annua* L. *Food Chemistry*, 115(4): 1240-1246. DOI: 10.1016/j.foodchem.2009.01.033
- Bueno JPR, de Mattos Nascimento MRB, da Silva Martins JM, Marchini CFP, Gotardo LRM, de Sousa GMR, Mundim AV, Guimarães EC & Rinaldi FP. 2017. Effect of age and cyclical heat stress on the serum biochemical profile of broiler chickens. *Semina-Ciencias Agrarias*, 38(3): 1383-1392. DOI: 10.5433/1679-0359.2017v38n3p1383
- Ceron JJ, Eckersall PD & Martynez-Subiela S. 2005. Acute phase proteins in dogs and cats: current knowledge and future perspectives. *Veterinary Clinical Pathology*, 34: 85-9. DOI: 10.1111/j.1939-165X.2005.tb00019.x
- Cherian G, Orr A, Burke IC & Pan W. 2013. Feeding *Artemisia annua* alters digesta pH and muscle lipid oxidation products in broiler chickens. *Poultry Science*, 92(4): 1085-1090. DOI: 10.3382/ps.2012-02752
- Chrubasik S, Pittler MH & Roufogalis BD. 2005. *Zingiberis rhizoma*: a comprehensive review on the ginger effect and efficacy profiles. *Phytomedicine*, 12(9): 684-701. DOI: 10.1016/j.phymed.2004.07.009
- Chung, ELT, Predith, M, Nobilly, F, Samsudin, AA, Jesse, FFA & Loh TC. 2018. Can treatment of *Brachiaria decumbens* (Signal grass) improve

- its utilisation in the diet in small ruminants? — a review. *Tropical Animal Health and Production*, 50, 1727–1732. DOI: 10.1007/s11250-018-1641-4
- Coles M, Graham B, Latorre J, Petrone-Garcia V, Hernandez-Velasco X, Castellanos-Huerta I, Sun X, Hargis B, El-Ashram S, Shehata A & Tellez-Isaias G. 2023. Essential oils as an alternative to antibiotics to reduce the incidence and severity of necrotic enteritis in broiler chickens: a short review. *Food Science and Nutrition*, 14: 233-257. DOI: 10.4236/fns.2023.143016
- Diarra SS & Anand S. 2020. Impact of commercial feed dilution with copra meal or cassava leaf meal and enzyme supplementation on broiler performance. *Poultry Science*, 99(11): 5867-5873. DOI: 10.1016/j.psj.2020.08.028
- Diarra SS, Anand S, Lemuel T, Areta F, Mathew E, Lehi E, Hoponoa S, Kava F, Mafi J & Taliauli F. 2023. Commercial feed diluted with different fiber sources and enzyme product for broilers: growth performance, carcass and gut health. *Brazilian Journal of Poultry Science*, 25(1). DOI: 10.1590/1806-9061-2021-1546.
- Dibner JJ & Richards JD. 2005. Antibiotic growth promoters in agriculture: history and mode of action. *Poultry Science*, 84(4): 634-643. DOI: 10.1093/ps/84.4.634
- Erwan E, Irawati E, Nazir A, Fitra D, Rodiallah M & Chowdhury VS. 2021a. The Effect of Yellow and Red-fleshed Watermelon Rind Powder Dietary Supplementation on Plasma Metabolites in Indigenous Chickens. *Journal of World's Poultry Research*. 11(4): 475-480. DOI: 10.36380/jwpr.2021.56.
- Erwan E, Irawati E, Nazir A, Fitra D, Rodiallah M. 2021b. Body temperature, feed intake and plasma metabolites of indigenous chicks by oral administration of watermelon rind extract with two different colors of flesh: yellow and red. *International Journal of Zoology and Animal Biology*, 4(3):1-6. DOI: 10.23880/izab-16000300
- EP (European Parliament and Council), 2003. Regulation (EC) No 1831/2003 of the European Parliament and of the Council of 22 September 2003 on additives for use in animal nutrition. *Official Journal of the European Union*, 268: 29-43.
- FAO (World Health Organization). 2004. Second Joint FAO/OIE/WHO Expert Workshop on Non-Human Antimicrobial Usage and Antimicrobial Resistance: Management options: 15-18 March 2004, Oslo, Norway (No. WHO/CDS/CPE/ZFK/2004.8). World Health Organization.
- Fathima S, Shanmugasundaram R, Sifri M & Selvaraj R. 2023. Yeasts and yeast-based products in poultry nutrition. *Journal of Applied Poultry Research*, 32(2): 100345. DOI: 10.1016/j.japr.2023.100345
- Ferdous MF, Arefin MS, Rahman MM, Ripon MMR, Rashid MH, Sultana MR, Hossain MT, Ahammad MU & Rafiq K. 2019. Beneficial effects of probiotic and phytobiotic as growth promoter alternative to antibiotic for safe broiler production. *Journal of Advance Veterinary and Animal Research*, 6(3): 409–415. DOI: 10.5455/javar.2019.f361
- Ferlito C. 2020. The poultry industry and its supply chain in Malaysia: challenges from the Covid19 emergency. 1-37. DOI: 10.13140/RG.2.2.23221.91367
- Fujioka T, Kondou R & Fukuhara A. 2003. Efficacy of glycyrrhizin suppository for treatment of chronic hepatitis C: a pilot study. *Hepatology Research*, 26(1): 103-117. DOI: 10.1016/S1386-6346(02)00332-7
- Galli GM, Gerbet RR, Griss LG, Fortuoso BF, Petrolli TG, Boiago MM, Souza CF, Baldissera MD, Mesadri J, Wagner R, da Rosa G, Mendes RE, Gris A & Da Silva AS. 2020. Combination of herbal components (curcumin, carvacrol, thymol, cinnamaldehyde) in broiler chicken feed: impacts on response parameters, performance, fatty acid profiles, meat quality and control of coccidia and bacteria. *Microbial Pathogenesis*, 139: 1–11. DOI: 10.1016/J.Micpath.2019.103916
- Ganter MT, Ware LB, Howard M, Roux J, Gartland B, Matthay MA, Fleshner M & Pittet JF. 2006. Extracellular heat shock protein 72 is a marker of the stress protein response in acute lung injury. *American Journal of Physiology-Lung Cellular and Molecular Physiology*, 291(3): L354-L361. DOI: 10.1152/ajplung.00405.2005
- Gholamrezaie Sani L, Mohammadi M, Jalali SJ, Abolghasemi SA & Roostaei AM. 2013. Extract and leaf powder effect of *Artemisia annua* on performance, cellular and humoral immunity in broilers. *Iranian Journal of Veterinary Research*, 14(1): 15 -20.
- Giannini EG, Testa R & Savarino V. 2005. Liver enzyme alteration: a guide for clinicians, *Canadian Medical Association Journal*, 172(3): 367–379.
- Gilani SMH, Rashid Z, Galani S, Ilyas S, Sahar S, Al-Ghanim K, Zehra S, Azhar A, Al-Misned F, Ahmed Z, Al-Mulham N & Mahboob S. 2021. Growth performance, intestinal histomorphology, gut microflora and ghrelin gene expression analysis of broiler by supplementing natural growth promoters: a nutrigenomics approach. *Saudi Journal of Biological Sciences*, 28(6): 3438-3447. DOI: 10.1016/j.sjbs.2021.03.008
- Gilani SMH, Zehra S, Galani S & Ashraf A. 2018. Effect of natural growth promoters on immunity, and biochemical and haematological parameters

- of broiler chickens. *Tropical Journal of Pharmaceutical Research*, 17(4): 627-633. DOI: 10.4314/tjpr.v17i4.9
- Gruys E, Toussaint MJ, Upragarin N, Van Ederen AM, Adewuyi AA, Candiani D, Nguyen TKA & Sabeckiene J. 2005. Acute phase reactants, challenge in the near future of animal production and veterinary medicine. *Journal of Zhejiang University-SCIENCE B*, 6(10): 941-947. DOI: 10.1007/BF02888485
- Hansen PJ. 2009. Effects of heat stress on mammalian reproduction. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364(1534): 3341-3350. DOI: 10.1098/rstb.2009.0131
- Hao H, Cheng G, Iqbal Z, Ai X, Hussain HI, Huang L, Dai M, Wang Y, Liu Z & Yuan Z. 2014. Benefits and risks of antimicrobial use in food-producing animals. *Frontiers in Microbiology*, 5: 288. DOI: 10.3389/fmicb.2014.00288
- Hassan HMA, El-Moniary MM, Hamouda Y, El-Daly EF, Youssef AW & El-Azeem NAA. 2016. Effect of different levels of *Moringa oleifera* leaves meal on productive performance, carcass characteristics and some blood parameters of broiler chicks reared under heat stress conditions. *Asian Journal of Animal and Veterinary Advances*, 11: 60–66.
- Hidayat C, Irawan A, Jayanegara A, Sholikin MM, Prihambodo TR, Yanza YR, Wina E, Sadarman S, Krisnan R & Isbandi I. 2021. Effect of dietary tannins on the performance, lymphoid organ weight, and amino acid ileal digestibility of broiler chickens: a meta-analysis. *Veterinary World*, 14(6): 1405. DOI: 10.14202/vetworld.2021.1405-1411
- Hidayati NA, Indarto RE, Suryanto E & Dono ND. 2022. Intestinal health in broiler chickens treated with nanoencapsulation of *Terminalia catappa* leaf extract as an antibacterial agent. *Tropical Animal Science Journal*, 45(4): 443-450. DOI: 10.5398/tasj.2022.45.4.443
- Huang Q, Liu X, Zhao G, Hu T & Wang Y. 2018. Potential and challenges of tannins as an alternative to in-feed antibiotics for farm animal production. *Animal Nutrition*, 4: 137-150. DOI: 10.1016/j.aninu.2017.09.004
- Hughes P & Heritage J. 2004. Antibiotic growth-promoters in food animals. In F. A. Nations, Food and Agriculture Organization of the United Nations 2004 No. 160 (P. 152). Rome: Food and Agriculture Organization of the United Nations. DOI: 10.1002/jsf2.37
- Jamroz D, Wartecki T, Houszka M & Kamel C. 2006. Influence of diet type on the inclusion of plant origin active substances on morphological and histochemical characteristics of the stomach and jejunum walls in chicken. *Journal of Animal Physiology and Animal Nutrition*, 90(5-6): 255-268. DOI: 10.1111/j.1439-0396.2005.00603.x
- Janeway CA, Travers P, Walport M & Shlomchik MJ. 2001. *Immunobiology*, 5th edition, p732. London (UK): Taylor and Francis.
- Jha R & Mukku KK. 2019. Uremic toxins, oxidative stress, and inflammation in chronic kidney disease. *Journal of Renal Nutrition and Metabolism*, 5: 48-50. DOI: 10.4103/jrn.jrn_57_19
- Jimenez-Garcia SN, Vazquez-Cruz MA, Garcia-Mier L, Contreras-Medina LM, Guevara-González RG, Garcia-Trejo JF & Feregrino-Perez AA. 2018. Phytochemical and pharmacological properties of secondary metabolites in berries. *Therapeutic Foods*: 397–427. DOI: 10.1016/b978-0-12-811517-6.00013-1
- Khan I, Zaneb H, Masood S, Yousaf MS, Rehman HF & Rehman H. 2017. Effect of *Moringa oleifera* leaf powder supplementation on growth performance and intestinal morphology in broiler chickens. *Journal of Animal Physiology and Animal Nutrition*, 101: 114-121. DOI: 10.1111/jpn.12634
- Khan SH & Iqbal J. 2016. Recent advances in the role of organic acids in poultry nutrition. *Journal of Applied Animal Research*, 44(1): 359-369. DOI: 10.1080/09712119.2015.1079527
- Kuralkar P & Kuralkar SV. 2021. Role of herbal products in animal production—an updated review. *Journal of Ethnopharmacology*, 278: 114246. DOI: 10.1016/j.jep.2021.114246
- Lala AO, Ajayi OL, Okwelum N, Oso AO, Fakorede TV, Adebayo TA & Jagbojo JE. 2018. Haematological, biochemical and organ changes in broiler chickens fed varying levels of *Morinda lucida* (brimstone) leaf meal supplementation in the diets. *Tropical Animal and Health Production*, 50: 1005-1010. DOI: 10.1007/s11250-018-1524-8.
- Li JY, Lu Y, Hu S, Sun D & Yao YM. 2002. Preventive effect of glutamine on intestinal barrier dysfunction induced by severe trauma. *World Journal of Gastroenterology*. 8: 168–171. DOI: 10.3748/wjg.v8.i1.168
- Li Y, Zhang H, Chen YP, Yang MX, Zhang LL, Lu ZX, Zhou YM & Wang T. 2015. *Bacillus amyloliquefaciens* supplementation alleviates immunological stress and intestinal damage in lipopolysaccharide-challenged broilers. *Animal Feed Science and Technology*, 208: 119-131. DOI: 10.1016/j.anifeedsci.2015.07.001
- Liaqat S, Mahmood S, Ahmad S, Kamran Z & Koutoulis KC. 2016. Replacement of canola meal with *Moringa oleifera* leaf powder affects performance and immune response in broilers. *Journal of Applied Poultry Research*, 25: 352–358. DOI: 10.3382/japr/pfw018

- Low S. 2015. Signal grass (*Brachiaria decumbens*) toxicity in grazing ruminants. *Agriculture*, 5(4): 971-990. DOI: 10.3390/agriculture5040971
- Mandal AB, Biswas A, Yadav AS & Biswas AK. 2014. Effect of dietary *Moringa oleifera* leaves powder on growth performance, blood chemistry, meat quality and gut microflora of broiler chicks. *Animal Nutrition and Feed Technology*, 14(2): 349-357. DOI: 10.5958/0974-181X.2014.01324.9
- Ming J, Xie J, Xu P, Liu W, Ge X, Liu B, He Y, Cheng Y, Zhou Q & Pan L. 2010. Molecular cloning and expression of two HSP70 genes in the Wuchang bream (*Megalobrama amblycephala* Yih). *Fish & Shellfish Immunology*, 28(3): 407-418. DOI: 10.1016/j.fsi.2009.11.018
- Mohammadi Gheisar M & Kim IH. 2018. Phytobiotics in poultry and swine nutrition—a review. *Italian Journal of Animal Science*, 17(1): 92-99. DOI: 10.1080/1828051X.2017.1350120
- Muazu U & Aliyu-Paiko M. 2020. Evaluating the potentials of *Carica* papaya seed as phytobiotic to improve feed efficiency, growth performance and serum biochemical parameters in broiler chickens. *Journal of Biochemistry and Biotechnology*, 6(1): 8-18. DOI: 10.9790/264X-0601010818
- Mustafa MA. 2019. Effect of Eucalyptus leaves and its supplementation with diet on broiler performance, microbial and physiological statuses to alleviate cold stress. *Iraqi Journal of Agricultural Sciences*, 50(1). DOI: 10.36103/ijas.v50i1.302
- Nazifi S, Dadras H, Hoseinian SA, Ansari-Lari M & Masoudian M. 2010. Measuring acute phase proteins (haptoglobin, ceruloplasmin, serum amyloid A, and fibrinogen) in healthy and infectious bursal disease virus-infected chicks. *Comparative Clinical Pathology*, 19: 283-286. DOI: 10.1007/s00580-009-0858-z
- Ncube S, Saidi PT, Tivapasi MT, Imbayarwo-Chikosi VE & Halimani TE. 2018. Serum lipid, enzyme, and hematological responses of broilers fed *Acacia angustissima* leaf meal-based diets. *Tropical Animal and Health Production*, 50: 665-670. DOI: 10.1007/s11250-017-1484-4
- Ngamukote S, Makynen K, Thilawech T & Adisakwattana S. 2011. Cholesterol-lowering activity of the major polyphenols in grape seed. *Molecules*, 16: 5054-5061. DOI: 10.3390/molecules16065054
- Ogwuegbu MC, Ani AO, Oyeagu CE, Osita CO, Oyeagu U, Ugwuoke WI & Lewu FB. 2021. Sodium butyrate and rosemary leaf meal inclusion in broiler diet: effects on gut micro-floral, growth performance, ileum, jejunum and duodenal histological traits. *Advances in Animal and Veterinary Sciences*, 9(7): 1095-1112. DOI: 10.17582/journal.aavs/2021/9.7.1095.1112
- Ojo O, Ojo A, Barnabas M, Iyobhebhe M, Elebiyo T, Evbuomwan IO, Michael T, Ajiboye BO, Oyinloye BE & Oloyede O. 2022. Phytochemical properties and pharmacological activities of the genus *Pennisetum*: a review. *Scientific African*, 16. DOI: 10.1016/J.Sciaf.2022.E01132
- Ojo RJ, Oguche PI, Kube GD & Udzer TE. 2013. Effect of *Jatropha curcas* supplemented diet on broilers. *Scholars Academic Journal of Biosciences*, 1(6): 329–336.
- Oloruntola OD, Ayodele SO, Omoniyi IS, Adeyeye SA & Adegbeye MJ. 2022. The effect of dietary supplementation of mucuna leaf meal on the growth performance, blood parameters, and carcass quality of broiler. *Acta Scientiarum: Animal Sciences*, 44. DOI: 10.4025/actascianimsci.v44i1.55362
- Olukosi OA & Dono ND. 2014. Modification of digesta pH and intestinal morphology with the use of benzoic acid or phytobiotics and the effects on broiler chicken growth performance and energy and nutrient utilization. *Journal of Animal Science*, 92(9): 3945-3953. DOI: 10.2527/jas.2013-6368
- Pohl SA, Caldwell DJ, Lee JT, Coppedge JR, Dunn-Horrocks SL, Stringfellow KD, Jessen K & Farnell MB. 2012. Effects of dietary calcium formate inclusion on broiler performance, skeletal development, and gut maturation. *Journal of Applied Poultry Research*, 21(2): 311-317. DOI: 10.3382/japr.2011-00400
- Rahman MM & Yang DK. 2018. Effects of *Ananas comosus* leaf powder on broiler performance, haematology, biochemistry, and gut microbial population. *Revista Brasileira de Zootecnia*, 47: 1–6. DOI: 10.1590/rbz4720170064
- Reda FM, El-Saadony MT, Elnesr SS, Alagawany M & Tufarelli V. 2020. Effect of dietary supplementation of biological curcumin nanoparticles on growth and carcass traits, antioxidant status, immunity and caecal microbiota of Japanese quails. *Animals*, 10(5): 754. DOI: 10.3390/ani10050754
- Rehman MS, Mahmud A, Mehmood S, Pasha TN, Hussain J & Khan MT. 2017. Blood biochemistry and immune response in Aseel chicken under free range, semi-intensive, and confinement rearing systems. *Poultry Science*, 96(1): 226-233. DOI: 10.3382/ps/pew278
- Reis JH, Gebert RR, Barreta M, Baldissera MD, Dos Santos ID, Wagner R, Campigotto G, Jaguezeski AM, Gris A, De Lima JLF, Mendes RE, Fracasso M, Boiogo MM, Stefani LM, dos Santos DS, Robazza WS & da Silva AS. 2018. Effects of phytochemical feed additive based on thymol, carvacrol and cinnamic aldehyde on body weight, blood parameters and environmental bacteria in

- broilers chickens. *Microbial Pathogenesis*, 125: 168-176. DOI: 10.1016/j.micpath.2018.09.015
- Reis JH, Gebert RR, Barreta M, Boiogo MM, Souza CF, Baldissera MD, Santos ID, Wagner R, Laporta LV, Stefani LM & da Silva AS. 2019. Addition of grape pomace flour in the diet on laying hens in heat stress: impacts on health and performance as well as the fatty acid profile and total antioxidant capacity in the egg. *Journal of Thermal Biology*, 80: 141-149. DOI: 10.1016/j.jtherbio.2019.01.003
- Rizwanuddin S, Kumar V, Naik B, Singh P, Mishra S, Rustagi S & Kumar V. 2023. Microbial phytase: their sources, production, and role in the enhancement of nutritional aspects of food and feed additives. *Journal of Agriculture and Food Research*, 100559. DOI: 10.1016/j.jafr.2023.100559
- Saadatmand N, Toghyani M & Gheisari A. 2019. Effects of dietary fiber and threonine on performance, intestinal morphology and immune responses in broiler chickens. *Animal Nutrition*, 5(3): 248-255. DOI: 10.1016/J.Aninu.2019.06.001
- Saharan V, Pathak AK, Sharma RK, Sharma N, Sarma K & Kumar H. 2022. Influence of *Psidium guajava* leaf meal supplementation on growth performance, nutrient utilization, intestinal micrometry, caecal fermentative metabolites and microbiota of broiler chickens. *Animal Nutrition and Feed Technology*, 22(2): 367-382. DOI: 10.51227/ojafr.2023.12
- Saeed F, Afzaal M, Tufail T & Ahmad A. 2019. Use of natural antimicrobial agents: a safe preservation approach. *Active Antimicrobial Food Packaging*, 18(0).
- Sahin N, Tuzcu M, Orhan C, Onderci M, Eroksuz Y & Sahin K. 2009. The effects of vitamin C and E supplementation on heat shock protein 70 response of ovary and brain in heat-stressed quail. *British Poultry Science*, 50(2): 259- 265. DOI: 10.1080/00071660902758981
- Sebola NA & Mokoboki HK. 2019. Influence of dietary *Moringa oleifera* leaf meal on haematological parameters, serum biochemical indices and weight of internal organs of chickens. *Advances in Animal and Veterinary Sciences*, 7(12): 1042-1048. DOI: 10.17582/journal.aavs/2019/7.12.1042.1048
- Song Z, Cheng K, Zhang L & Wang T. 2017. Dietary supplementation of enzymatically treated *Artemisia annua* could alleviate the intestinal inflammatory response in heat-stressed broilers. *Journal of Thermal Biology*, 69: 184-190. DOI: 10.1016/j.jtherbio.2017.07.015
- Staib JL, Quindry JC, French JP, Criswell DS & Powers SK. 2007. Increased temperature, not cardiac load, activates heat shock transcription factor 1 and heat shock protein 72 expression in the heart. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 292(1): R432-R439. DOI: 10.1152/ajpregu.00895.2005
- Sugiharto S, Widiastuti E, Isroli I, Yudiarti T, Sartono TA & Wahyuni HI. 2020. Effect of feeding fermented mixture of cassava pulp and *Moringa oleifera* leaf meal on immune responses, antioxidative status, biochemistry indices, and intestinal ecology of broilers. *Veterinary World*, 13(2): 392. DOI: 10.14202/vetworld.2020.392-399
- Surai PF. 2014. Polyphenol compounds in the chicken/animal diet: from the past to the future. *Journal of Animal Physiology and Animal Nutrition*, 98(1): 19-31. DOI: 10.1111/jpn.12070.
- Suresh G, Das RK, Kaur Brar S, Rouissi T, Avalos Ramirez A, Chorfi Y & Godbout S. 2017. Alternatives to antibiotics in poultry feed: molecular perspectives. *Critical Reviews in Microbiology*, 44(3): 318-335. DOI: 10.1080/1040841x.2017.137306
- Tamzil MH, Noor RR, Hardjosworo PS, Manalu W & Sumantri C. 2013. Acute heat stress responses of three lines of chickens with different heat shock protein (HSP)-70 genotypes. *International Journal of Poultry Science*, 12(5): 264-272.
- Tang S, Yin B, Xu J & Bao E. 2018. Rosemary reduces heat stress by inducing CRYAB and HSP70 expression in broiler chickens. *Oxidative Medicine and Cellular Longevity*. DOI: 10.1155/2018/7014126
- Tavangar P, Gharahveysi S, Rezaeipour V & Irani M. 2021. Efficacy of phytobiotic and toxin binder feed additives individually or in combination on the growth performance, blood biochemical parameters, intestinal morphology, and microbial population in broiler chickens exposed to aflatoxin B1. *Tropical Animal and Health Production*, 53(3): 335. DOI: 10.1007/s11250-021-02778-0
- Tokofai MB, Idoh K & Agbonon A. 2020. Growth performance, haematological and biochemical parameters in broilers fed diets with varying levels of *Vernonia amygdalina* leaf meal. *European Poultry Science*, 84. DOI: 10.1399/eps.2020.321
- Valenzuela-Grijalva NV, Pinelli-Saavedra A, Muhlia-Almazan A, Domínguez-Díaz D & González-Ríos H. 2017. Dietary inclusion effects of phytochemicals as growth promoters in animal production. *Journal of Animal Science and Technology*, 59(1): 1-17. DOI: 10.1186/s40781-017-0133-9
- Viveros A, Chamorro S, Pizarro M, Arijia I, Centeno C & Brenes A. 2011. Effects of dietary polyphenol-rich grape products on intestinal microflora and gut morphology in broiler chicks.

- Poultry Science, 90(3): 566-578. DOI: 10.3382/ps.2010-00889
- Widiastuti E, Sartono TA, Wahyuni HI, Murwani R, Yudiarti T, Pratama AR & Sugiharto S. 2021. Acidification of papaya leaf and seed meal using *Averrhoa bilimbi* L. fruit filtrate and their effect on growth and carcass traits of broiler chickens. *Advances in Animal and Veterinary Sciences*, 9(9): 1368-1375. DOI: 10.17582/journal.aavs/2021/9.9.1368.1375
- Yilmaz B & Li H. 2018. Gut microbiota and iron: the crucial actors in health and disease. *Pharmaceuticals*, 11: 98. DOI: 10.3390/ph11040098
- Zdunczyk Z, Gruzauskas R, Juskievicz J, Semaskaite A, Jankowski J, Godycka-Klos I, Jarule V, Miezeliene A & Alencikiene G. 2010. Growth performance, gastrointestinal tract responses, and meat characteristics of broiler chickens fed a diet containing the natural alkaloid sanguinarine from *Macleaya cordata*. *Journal of Applied Poultry Research*, 19(4): 393-400. DOI: 10.3382/japr.2009-00114
- Zhang Y, Yang L, Zu Y, Chen X, Wang F & Liu F. 2010. Oxidative stability of sunflower oil supplemented with carnosic acid compared with synthetic antioxidants during accelerated storage. *Food Chemistry*, 118(3): 656-662. DOI: 10.1016/j.foodchem.2009.05.038