



## ORIGINAL ARTICLE

## Impact of Calcium Supplementation on Photosynthetic Pigments, Compatible Osmolytes Contents and Membrane Stability Index in Triticale (x *Triticosecale* Wittmack) Exposed to Salinity Stress

Malihe Jahani<sup>\*1,2</sup>, Mohammad Reza Hadi<sup>2</sup>, Mojtaba Jafarinia<sup>2</sup>, Sedighe Jahani<sup>1</sup>

<sup>1</sup>Department of Biology, Science and Research Branch, Islamic Azad University, Tehran, Iran

<sup>2</sup>Department of Biology, Marvdasht Branch, Islamic Azad University, Marvdasht, Iran

(Received: 31 May 2020

Accepted: 18 September 2021)

## KEYWORDS

Ameliorative impact;  
Carbohydrate;  
Sodium-calcium  
interactions;  
Triticale

**ABSTRACT:** In many areas, salinization is considered as one of the most serious dangers to environmental resources and human health. Calcium has a crucial role in plant resistance to salinity stress. In order to investigate the impact of calcium supplementation on photosynthetic pigments, compatible osmolytes contents and membrane stability index (MSI) in triticale (x *Triticosecale* Wittmack) exposed to salinity stress, an experiment as a completely randomized design with 3 replications in greenhouse condition ( $25 \pm 2^\circ\text{C}$ , 35% relative humidity, 16-hour photoperiod) was conducted. The seeds were germinated in soil. One week old triticale seedlings (with two leaves) were imposed by 0, 50, 100 and 150  $\text{mmol L}^{-1}$  NaCl and 0, 6 and 10  $\text{mmol L}^{-1}$   $\text{CaCl}_2$  for 5 weeks and assayed for some morpho-physiological parameters including fresh weight (FW) and dry weight (DW) of shoot, photosynthetic pigments (chlorophyll (Chl) a and Chl b, total Chl and carotenoids (Car)) contents, proline and glycine betaine (GB) contents, soluble sugars and starch contents and MSI in leaves. Results showed that with incrementing salinity meaningfully decremented FW and DW of shoot, photosynthetic pigments, starch content and MSI while proline, GB and soluble sugars contents incremented in leaves. Calcium treatment meaningfully incremented FW and DW of shoot, photosynthetic pigments, starch content and MSI but caused a meaningful decline in proline, GB and soluble sugars contents in leaves. It can be concluded that calcium had exerted an ameliorative impact on triticale under salinity stress. Maximum ameliorative impact of calcium was observed in plants exposed to 6  $\text{mmol L}^{-1}$   $\text{CaCl}_2$ .

## INTRODUCTION

At the current, most agricultural areas are seriously affected by salinity, which has emerged as a serious global issue owing to the potential detrimental impacts on human and animal health [1]. Calcium is considered as an essential element in most plants processes [2]. Moreover, calcium is as a crucial mineral for human health and it aids in the formation of teeth and bones. Also, calcium assists muscle movement and nerve communication [3]. Calcium ( $\text{Ca}^{2+}$ )

has been demonstrated to be a major indicator of plant salt resistance, conferring preserving benefits on plants growing in sodic soils [4].  $\text{Ca}^{2+}$  is required for the functional and structural integrity of cell membranes of plant, as well as the stabilization of cell wall structures, the regulation of ion transport and selectivity, and the control of ion-exchange behavior and cell wall enzyme activities [5]. Depending on the genotype of the plant, the nature of these reactions will

\*Corresponding author: malihe.jahani2009@gmail.com (M. Jahani)  
DOI: 10.22034/jchr.2021.1901194.1141

differ [4, 5].  $\text{Ca}^{2+}$  regulates the function, structure, and signaling of membranes by forming bonds with the phospholipid bilayer, therefore maintaining and enhancing the structural integrity of membrane organelles in plants under stress [6].  $\text{Ca}^{2+}$  serves as a second messenger in signaling of stress, which is one of its most important roles [4,7].

Plant growth is slowed by salinity stress, which is one of the most major osmotic stresses [8]. The adverse impact of salinity stress in plants is its osmotic impacts, ion toxicity impact and interference in nutrient absorption [9, 1]. In a study on rice, it was reported that salt stress diminished the fresh weight (FW) and dry weight (DW) of shoot, while the addition of calcium treatment diminished the adverse impacts of salinity stress and incremented FW and DW of shoot [10].

Salinity stress promotes the buildup of reactive oxygen species (ROS), which is a critical factor in plant growth retarding [8, 11]. Plants create a multitude of compatible solutes and antioxidant components that are triggered to give secondary protection against oxidative stress in order to minimize cellular damage caused by ROS buildup [12].

The buildup of adaptive solutions is one of the strategies against salinity stress in plants [11]. These solutions include essential metal ions such as  $\text{K}^+$  and they are basically organic solutions. The most major organic osmotic solutions which accumulate in plants due to salinity include soluble sugars, proline, glycine betaine (GB), and etc. [13, 14].

Salinity stress raises proline synthesis in the plant [15]. Proline is an essential amino acid and important osmolyte that is soluble in water and it is a response to the initial defense to balance osmotic pressure in cells under salinity stress [9]. It was reported that salinity stress raised proline levels in cotton, while addition of calcium treatment to the saline environment diminished proline levels [16].

Photosynthesis is one of the important physiological processes in plants, which is affected by genetic and environmental factors [11]. It was reported that the addition of calcium treatment to sweet sorghum exposed to salt stress diminished the adverse impacts of stress and raised the chlorophyll (Chl) level [17].

Triticale ( $\times$ *Triticosecale* Wittmack) is a human-made cereal that combines the functionality and high production of wheat (*Triticum* spp.) with the durability of rye (*Secale cereale* L.) [18]. Triticale has been planted all over the world, mostly for grain and fodder production, but also for bioenergy generation in recent years [19]. Furthermore, in the human food market, it likewise has a developing potential [20]. Triticale contains a lot of non-starch polysaccharides (such as arabinoxylans), phenolic acid, and anthocyanin, which can cause to prevent heart disease, cancer, diabetes, and neurodegenerative disorders [21]. Also, triticale has antioxidant and antihypertensive properties [22]. It will become a key cereal crop for world populations in the near future [19]. Due to the value of triticale as a cereal crop plant with great developmental potential in the food market and the expansion of saline land areas, identifying a component for salinity stress tolerance in triticale is critical.

Since salinity is one of the most major problems in different regions of the world and calcium plays a crucial role in resistance of plants to salinity, it is necessary to study the sodium-calcium interaction in plants. To the best of the authors' knowledge, there is no study on the interaction of sodium-calcium with triticale. This study provides baseline information on the influence of calcium chloride on triticale exposed to salinity stress. The present research was carried out to investigate the impact of calcium supplementation on photosynthetic pigments, compatible osmolytes contents and membrane stability index (MSI) in triticale ( $\times$  *Triticosecale* Wittmack) exposed to salinity stress. This will provide a simple, cheap and economic solution to deal with salinity and further enhance sustainable agriculture.

## MATERIALS AND METHODS

### *Growth condition and treatments*


To investigate the interactive impact of sodium chloride and calcium chloride on physio-biochemical parameters in triticale leaves, a research as a completely randomized design with three replications was conducted in greenhouse condition (temperature  $25\pm 2^\circ\text{C}$ , relative humidity 35%, with a 16-hour photoperiod). Seeds of triticale cultivar Moreno

were planted in soil. Physico-chemical properties of soil are shown in Table 1. One week after seed cultivation in the soil, seedlings were subjected to various doses of sodium chloride and calcium chloride. Treatments included 0, 50, 100 and 150 mmol L<sup>-1</sup> NaCl levels and 0, 6 and 10mmol L<sup>-1</sup> CaCl<sub>2</sub> levels. Applying the treatments (as addition to the


soil) was done for 5 weeks. After the treatment period, 42-day-old plants (Figure 1-4) harvested and then evaluated some morpho-physiological parameters including FW and DW of shoot, photosynthetic pigments, proline and GB, soluble sugars and starch contents and MSI in the leaves.

**Table 1.** Physico-chemical properties of soil.

EC (dS m <sup>-1</sup> )	pH	Organic matter (%)	Saturation percentage (%)	Sand (%)	Silt (%)	Clay (%)	Texture
0.838	7.31	1.231	54.713	63.5	26.5	10	Sandy Loam
Soluble Cations (mEq 100 g <sup>-1</sup> soil)				Soluble Anions (mEq 100 g <sup>-1</sup> soil)			
Na <sup>+</sup>	K <sup>+</sup>	Ca <sup>+2</sup>	Mg <sup>+2</sup>	SO <sub>4</sub> <sup>-2</sup>	Cl <sup>-</sup>	HCO <sub>3</sub> <sup>-</sup>	
0.059	0.037	0.353	0.1105	0.074	0.0191	0.402	




**Figure 1.** The impact of various doses of sodium chloride on vegetative growth of triticale plants (photograph of 42-day-old plants).



**Figure 2.** The impact of various doses of calcium chloride on vegetative growth of triticale plants exposed 50 mmol L<sup>-1</sup> sodium chloride treatment (photograph of 42-day-old plants).



**Figure 3.** The impact of various doses of calcium chloride on vegetative growth of triticale plants exposed 100 mmol L<sup>-1</sup> sodium chloride treatment (photograph of 42-day-old plants).



**Figure 4.** The impact of various doses of calcium chloride on vegetative growth of triticale plants exposed 150 mmol L<sup>-1</sup> sodium chloride treatment (photograph of 42-day-old plants).

### Growth analysis

To determine shoot FW, FW of each sample was measured with a Sartorius digital scale model TE214S with an accuracy level of 0.0001 g. Then for measuring shoot DW, each sample was separately inserted into an aluminum foil and dried in an oven at 70 °C for 48 h and then weighed with the same digital scale.

### Physio-biochemical analysis

The contents of Chls (Chl-a, Chl-b, and total Chl) and Car were assessed by a spectrophotometric method at wavelengths of 646.8, 663.2, and 470 nm [23].

The amount of proline was assayed by a spectrophotometric method [24]. Utilized technique focuses on the creation of a color combination under acidic circumstances at 100 °C via a reaction between the proline and ninhydrin reagent. At 520 nm, the proline content was measured and finally computed utilizing a standard curve. Moreover, amount of GB was analyzed at 365 nm utilizing a spectrophotometer [25].

The phenol sulfuric acid technique [26] was utilized to assess the amount of soluble sugars and insoluble sugars (starch). A spectrophotometer was utilized to detect the absorbance at 485 nm, and the soluble sugars and starch content were computed utilizing a standard glucose curve.

For MSI estimation, leaf discs were washed in diH<sub>2</sub>O and put in two groups of vials containing 15 mL of ddH<sub>2</sub>O. At

25 °C, one group was incubated for 2 hours. After that, the solution's electrical conductivity (EC<sub>1</sub>) was estimated. The conductivity (EC<sub>2</sub>) of the second group was estimated after it was heated in a water bath for 20 minutes at 95 °C. Afterwards, MSI was defined utilizing the following formula [27]:


$$MSI = [1 - (EC_1 / EC_2)] \times 100$$

### Statistical analysis

The experiment was conducted in a completely randomised design with three independent repetitions. Statistical analyses were performed utilizing analysis of variance (ANOVA) by SPSS v.22 software and were expressed as the mean values  $\pm$  SD. The significance of differences between treatments was evaluated utilizing Tukey's test at 5% probability level.

## RESULTS

The results showed that, as salinity incremented, shoot FW and DW diminished meaningfully, but addition of calcium treatment to the salinity environment incremented FW and DW of shoot meaningfully. The highest FW and DW was related to the control plant, and the lowest FW and DW was related to the plant imposed by 150 mmol L<sup>-1</sup> sodium chloride, which meaningfully diminished 67% and 68.67% respectively compared to the control plant (Figure 5).




**Figure 5.** The interaction impacts of NaCl and CaCl<sub>2</sub> on (A) fresh weight and (B) dry weight of triticale shoot. Values are mean of three replicates ± SD. Different letters represent a meaningful difference between treatments (P<0.05).

The results showed that, as salinity incremented, Chl-a, Chl-b, total Chl and Car content in leaf diminished meaningfully, while addition of calcium treatment to the saline environment incremented Chl-a, Chl-b and total Chl content in leaf meaningfully (Figure 6).

The highest Chl-a, Chl-b and total Chl content was observed in the control and the lowest Chl-a content was observed in the plant imposed by 150 mmol L<sup>-1</sup> sodium chloride, had decremented 53.33% respectively compared with the control plant (Figure 6). The lowest Chl-b content was observed in the plant imposed by 150 mmol L<sup>-1</sup> sodium chloride which decremented 79.22% compared to the control plant (Figure 6).


The lowest total Chl content was observed in the plant imposed by 150 mmol L<sup>-1</sup> sodium chloride which decremented 59.60% compared to the control plant (Figure 6). The highest carotenoid content was observed in the control and the plant imposed by 50 mmol L<sup>-1</sup> sodium chloride and 6 mmol L<sup>-1</sup> calcium chloride. The lowest carotenoid content was observed in the plant imposed by 150 mmol L<sup>-1</sup> sodium chloride and the plant imposed by 150 mmol L<sup>-1</sup> sodium chloride and 10 mmol L<sup>-1</sup> calcium chloride, had decremented 68.44% and 63.53% respectively compared with the control plant (Figure 6).



**Figure 6.** Interaction impacts of NaCl and CaCl<sub>2</sub> on (A) chlorophyll (Chl) a, (B) Chl-b, (C) total Chl and (D) carotenoids (Car) contents in triticale leaf. Values are mean of three replicates ± SD. Different letters represent a meaningful difference between treatments (P≤0.05).

Results showed that the highest proline and GB contents were observed in the plant exposed to 150 mmol L<sup>-1</sup> sodium chloride with an increment of 2.28 and 2.08 times

respectively compared with the control plant and the lowest proline content was related to the control plant (Figure 7).



**Figure 7.** Interaction impacts of NaCl and CaCl<sub>2</sub> on (A) the proline and (B) glycine betaine contents in triticale leaf. Values are mean of three replicates ± SD. Different letters represent a meaningful difference between treatments (P≤0.05).

The results showed that, as salinity incremented, soluble sugars contents incremented meaningfully, but addition of calcium treatment to the salinity environment decremented soluble sugars content meaningfully. The highest soluble


sugars content was related to the plant exposed to 150 mmol L<sup>-1</sup> sodium chloride, which meaningfully incremented 1.38 times compared to the control plant, and the lowest



soluble sugars content was related to the control plant (Figure 8).

Also, the results showed that starch content decremented under salinity stress but incremented slightly with calcium supplementation. The highest starch content was observed


in the control and the lowest starch content was observed in the plant imposed by 150 mmol L<sup>-1</sup> sodium chloride with a diminish of 39.26% compared with the control plant (Figure 8).



**Figure 8.** Interaction impacts of NaCl and CaCl<sub>2</sub> on (A) the soluble sugars and (B) starch contents in triticale leaf. Values are mean of three replicates ± SD. Different letters represent a meaningful difference between treatments (P≤0.05).

The results showed that leaf MSI decremented under salinity stress but raised with calcium addition. The highest MSI was related to the control plant and the lowest MSI

was related to the plant exposed to 150 mmol L<sup>-1</sup> sodium chloride, which meaningfully decremented 32.24% compared to the control plant (Figure 9).



**Figure 9.** Interaction impacts of NaCl and CaCl<sub>2</sub> on the membrane stability index (MSI) in triticale leaf. Values are mean of three replicates ± SD. Different letters represent a meaningful difference between treatments (P≤0.05).

## DISCUSSION

In the present study, salinity resulted in a meaningful decline in the shoot biomass while the addition of calcium incremented the shoot biomass meaningfully (Figure 5). The reason for the weight loss caused by salinity could probably be due to the presence of many harmful ions such as Na<sup>+</sup> and Cl<sup>-</sup> in the saline environment where these ions are harmful themselves or cause interference in other nutrients absorption, for example the competition of K<sup>+</sup> and

Na<sup>+</sup> or Cl<sup>-</sup> and NO<sup>-3</sup> causes interference in nutrients absorption in the plant [28].

In studies conducted on tomato [29, 30, 31], barley [32], sorghum [33], rice [34], cotton [16] and soybean [35], it was reported that salinity stress decremented shoot FW and DW, while calcium addition to the saline environment incremented shoot FW and DW, which is in agreement with the results of the present study. Also, salinity stress may

cause a secondary osmotic stress or a physiological drought stress in the plant which prevents water lost by the closure of stomata in the leaves. Stress also diminishes leaf area and ultimately diminishes photosynthesis and dry matter [28].

It has been reported that the addition of calcium to saline environments decrements the adverse impacts of salinity on plant growth, by maintaining the structure and integrity of cell membrane and incrementing cell division, diminishing sodium absorption and transport to the upper organs, incrementing potassium absorption and therefore incrementing the potassium to sodium ratio in the plant, enhancing nitrogen metabolism and photosynthetic activity [36, 5].

In the current study, parallel to the inhibitory impacts of salinity on growth, there was a meaningful decrement in the photosynthetic pigments content under salt stress but addition of calcium treatment incremented meaningfully photosynthetic pigments content in treated seedlings (Figure 6). In studies conducted on soybean [35], wheat [37] indian mustard [38], chili pepper [39], tomato [40] and rice [34, 41], it was reported that salinity stress diminished the Chl content while calcium addition to the saline environments incremented the Chl content, which is consistent with the results of the current research.

In salinity condition, the decline in photosynthesis and biomass has been related to stomatal closure [42], and the incremented generation of ROS in chloroplasts [43]. Calcium application under salt stress could cause to simultaneous increment in stomatal conductance and intercellular CO<sub>2</sub> dose and finally cause to increment of photosynthesis and plant biomass [44].

Data presented here showed that salinity treatments led to rising proline content in the treated triticale seedlings while addition of calcium treatment meaningfully diminished the proline content in seedlings (Figure 7a). Proline is an osmotic regulator that responds to salinity stress by incrementing [1, 15]. The incremented proline during stress is probably due to the increment of the proline biosynthesis enzymes activities such as P5CS (proline-5-carboxylate synthase) and P5CR (proline-5-carboxylate reductase) and decrement of the proline degradation enzymes activities

such as PROX (proline oxidase) and PDH (proline dehydrogenase) [45, 46].

In study on *Catharanthus roseus* seedlings, salt stress caused to increment in P5CS activity and proline content and decrement in PROX activity while addition of calcium to salinity-exposed plants caused to diminish in proline content by incrementing in PROX activity and decrement in P5CS activity [47]. Calcium application under salt stress could cause to increment in PROX activity and decrement in P5CS activity and finally cause to diminish in proline content [47]. In studies carried out on rice [41, 48], mung bean [49], peanut [50] and broad bean [51], it was reported that salinity stress incremented proline content, while addition of calcium to saline environments diminished the proline content, which is consistent with the results of the current research.

Furthermore, in the present study, salinity resulted in a meaningful increment in GB while the addition of calcium decremented GB meaningfully (Figure 7b). As compatible osmolytes, GB and proline preserve plants against environmental stressors [52, 53]. According to this study, the incremented amounts of proline and GB in response to salt stress might represent a metabolic adaptation to scavenge ROS. Similarly, a rise in GB levels owing to salt stress has been documented in tomato [54].

In the present study, salinity resulted in a meaningful increment in soluble sugars and a meaningful decrement in starch while the addition of calcium supplementation meaningfully decremented soluble sugars and slightly incremented starch (Figure 8). Probably the reason for the accumulation of soluble sugars during salinity stress is the insoluble sugars (starch) decompose and form soluble sugars to maintain osmotic potential and reduce the risk of dehydration [55]. In addition, stopping the growth and synthesis of sugars by non-photosynthetic pathway can be another factor in incrementing the concentration of soluble sugars in during of stress [55]. Calcium may decrement salinity stress in the plant due to reduced sodium accumulation, thereby reducing the need for the plant to accumulate soluble sugars as osmotic protectors [36]. Similarly, it was reported that salinity stress raised the amount of soluble sugars in rice [56]. Also, in cowpea, it



was reported that salinity stress incremented the soluble sugars content while adding calcium supplementation diminished the soluble sugars content [57].

Data presented here showed that MSI decremented under salinity stress but incremented with calcium addition (Figure 9). Here, the heightened  $\text{Na}^+$  levels owing to salinity stress may compromise cell membrane integrity, which results in ion leakage and disruption of cellular metabolism. Incremented permeability for ions, which may be easily detected by electrolyte leakage, is a great sign of cellular membrane disorder caused by salt stress [50]. Calcium ions may compete for membrane-binding sites with sodium ions. As a result, it's been proposed that high calcium levels can preserve cell membranes against the negative impacts of salt [5, 58]. The plasma membrane permeability to  $\text{Na}^+$  can be diminished by high  $\text{Ca}^{2+}$  levels. The passive influx of  $\text{Na}^+$  is decreased when membrane permeability to  $\text{Na}^+$  is decreased by  $\text{Ca}^{2+}$  [58]. Similarly, it was reported a diminish in MSI in tomato subjected to salinity stress and an improvement in MSI by calcium supplementation [40]. Also, in olive plant subjected to salinity stress was documented a decrement in electrolyte leakage by supplementary calcium [59].

### CONCLUSIONS

Calcium is a crucial mineral nutrient, and it is effective in diminishing the adverse impacts of salinity. In addition, because of the importance of triticale as a cultivating plant and the increment of extents of salty lands, determination of the optimal concentration of calcium is important for plants resistance in stress conditions. Results of this research revealed that the addition of calcium to saline environments decrements the adverse impacts of salinity stress and the maximum improving impacts of calcium was observed at a concentration of  $6 \text{ mmol L}^{-1}$  calcium chloride which can be a simple, cheap and economical strategy to deal with salinity stress and incrementing soil productivity and it may be recommended for utilize by farmers as a suitable option against salinity stress.

### ACKNOWLEDGMENTS

This research is as a part of M.Sc. thesis of Malihe Jahani and it received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

### Conflict of interests

The authors declare that there are no conflicts of interest related to this article.

### REFERENCES

1. Polash M.A.S., Sakil M.A., Hossain M.A., 2019. Plants responses and their physiological and biochemical defense mechanisms against salinity: a review. The Journal of the Society for Tropical Plant Research. 6(2), 250-274. doi: 10.22271/tpr.2019.v6.i2.035
2. Kader M.A., Lindberg S., 2010. Cytosolic calcium and pH signaling in plants under salinity stress. Plant Signaling and Behavior. 5(3), 233-238. doi: 10.4161/psb.5.3.10740
3. Pravina P., Sayaji D., Avinash M., 2013. Calcium and its role in human body. International Journal of Research in Pharmaceutical and Biomedical Sciences. 4(2), 659-668.
4. White P.J., Broadley M.R., 2003. Calcium in plants. Annals of Botany. 92(4), 487-511. <https://doi.org/10.1093/aob/mcg164>
5. Hadi M.R., Karimi N., 2012. The role of calcium in plants salt tolerance. Journal of Plants' Nutrition. 35(13), 2037-2054. <https://doi.org/10.1080/01904167.2012.717158>
6. Ahmad P., Abd\_Allah E.F., Alyemeni M.N., Wijaya L., Alam P., Bhardwaj R., Siddique K.H.M., 2018. Exogenous application of calcium to 24-epibrassinosteroid pre-treated tomato seedlings mitigates NaCl toxicity by modifying ascorbate–glutathione cycle and secondary metabolites. Scientific Reports. 8, 13515. <https://doi.org/10.1038/s41598-018-31917-1>
7. Manishankar P., Wang N., Köster P., Alatar A.A., Kudla J., 2018. Calcium signaling during salt stress and in the regulation of ion homeostasis, Journal of Experimental Botany. 69(17), 4215-4226.
8. Parihar P., Singh S., Singh R., Singh V.P., Prasad S.M., 2015. Effect of salinity stress on plants and its tolerance strategies: a review. Environmental Science and Pollution

- Research. 22(6), 4056-4075. doi: 10.1007/s11356-014-3739-1
9. Gupta B., Huang B., 2014. Mechanism of salinity tolerance in plants: physiological, biochemical, and molecular characterization. *International Journal of Genomics*. 2014(1), 1-18.
10. Cha-um S., Singh H.P., Samphumphuang T., Kirdmanee C., 2012. Calcium-alleviated salt tolerance in indica rice (*Oryza sativa* L. spp. indica): physiological and morphological changes. *Australian Journal of Crop Science*. 6(1), 176-182. [https:// search.informit.com. au/ documentSummary; dn=054025225882532;res=IELHSS](https://search.informit.com.au/documentSummary;dn=054025225882532;res=IELHSS)
11. Acosta-Motos J.R., Ortuño M.F., Bernal-Vicente A., Diaz-Vivancos P., Sanchez-Blanco M.J., Hernandez J.A., 2017. Plant responses to salt stress: adaptive mechanisms. *Agronomy*. 7(1), 1-38. [https://doi .org/ 10.3390/ agronomy7010018](https://doi.org/10.3390/agronomy7010018)
12. Nxele X., Klein A., Ndimba B.K., 2017. Drought and salinity stress alters ROS accumulation, water retention, and osmolyte content in sorghum plants. *South African Journal of Botany*. 108, 261-266. [https:// doi.org/10.1016 /j.sajb.2016.11.003](https://doi.org/10.1016/j.sajb.2016.11.003)
13. Parvaiz A., Satyawati S., 2008. Salt stress and phyto-biochemical responses of plants – a review. *Plant Soil and Environment*. 54(3), 89-99.
14. Suprasanna P., Nikalje G.C., Rai A.N., 2016. Osmolyte Accumulation and Implications in Plant Abiotic Stress Tolerance. In: Iqbal N., Nazar R., A. Khan N. (eds) *Osmolytes and Plants Acclimation to Changing Environment: Emerging Omics Technologies*. Springer, New Delhi. [https://doi.org/10.1007/978-81-322-2616-1\\_1](https://doi.org/10.1007/978-81-322-2616-1_1)
15. Alhasnawi A.N., 2019. Role of proline in plant stress tolerance: a mini review. *Research on Crops*. 20(1), 223-229. doi: 10.31830/2348-7542.2019.032
16. Amuthavalli P., Anbu D., Sivasankaramoorthy S., 2012. Effect of calcium chloride on growth and biochemical constituents of cotton (*Gossypium hirsutum* L.) under salt stress. *International Journal of Research in Botany*. 2(3), 9-12.
17. Zhou G., Ma B.L., 2012. Calcium addition affects germination and early seedling growth of sweet sorghum under saline conditions. *Agricultural Science and Technology*. 13(12), 2538-2543. [http:// eprints.icrisat. ac.in/id/eprint/9831](http://eprints.icrisat.ac.in/id/eprint/9831)
18. McGoverin C.M., Snyders F., Muller N., Botes W., Fox G., Manley M., 2011. A review of triticale uses and the effect of growth environment on grain quality. *Journal of the Science of Food and Agriculture*. 91(7), 1155-1165. doi: 10.1002/jsfa.4338
19. Mergoum M., Sapkota S., ElDoliefy A.E.A., Naraghi S.M., Pirseyedi S., Alamri M.S., AbuHammad W., 2019. Triticale (x *Triticosecale* Wittmack) Breeding. In: Al-Khayri J, Jain S, Johnson DV (Eds). *Advances in Plant Breeding Strategies: Cereals*. Springer, Cham. doi: 10.1007/978-3-030-23108-8
20. Langó B., Bóna L., Ács E., Tömösközi S., 2017. Nutritional features of triticale as affected by genotype, crop year, and location. *Acta Alimentaria*. 46(2), 238-245. <https://doi.org/10.1556/066.2017.46.2.14>
21. Chen H., Chen Z., Fu Y., Liu J., Lin S., Zhang Q., Liu Y., Wu D., Lin D., Han G., Wang L., Qin W., 2019. Structure, antioxidant, and hypoglycemic activities of arabinoxylans extracted by multiple methods from triticale. *Antioxidants*. 8(12), 584. doi: 10.3390/antiox8120584
22. Deng C., Zhang Z., Yan G., Wang F., Zhao L., Liu N., Abudurezike A., Li Y., Wang W., Shi S., 2020. Salt-responsive transcriptome analysis of triticale reveals candidate genes involved in the key metabolic pathway in response to salt stress. *Scientific Reports*. 10, 1-9. <https://doi.org/10.1038/s41598-020-77686-8>
23. Lichtenthaler H.K., 1987. Chlorophylls and carotenoids; pigments of photosynthetic biomembranes. In: Colowick SP, Kaplan NO (eds). *Methods in Enzymology*, Academic Press, San Diego, New York. 148, 350-382.
24. Bates L.S., Waldren R.P., Teare I.D., 1973. Rapid determination of free proline for water-stress studies. *Plant and Soil*. 39(1), 205-207. [https://doi.org/10. 1007/BF00018060](https://doi.org/10.1007/BF00018060)
25. Grieve C.M., Grattan, S.R., 1983. Rapid assay for determination of water soluble quaternary ammonium compounds. *Plant and Soil*. 70, 303-307. [https://doi. org/10.1007/BF02374789](https://doi.org/10.1007/BF02374789)
26. Kochert G., 1978. Carbohydrate determination by the phenol-sulfuric acid method. In: Hellebust JA, Craigie JS

[eds.], Handbook of Phycological Methods: Physiological and Biochemical Methods. 95-97. Cambridge University Press, Cambridge.

27. Premachandra G.S., Saneoka H., Ogata S., 1990. Cell membrane stability, an indicator of drought tolerance as affected by applied nitrogen in soybean. *Journal of agricultural science*. (Cambridge, Eng.). 115, 63-66.
28. Parida A.K., Das A.B., 2005. Salt tolerance and salinity effects on plants. a review. *Ecotoxicology and Environmental Safety*. 60(3), 324-349. doi: 10.1016/j.ecoenv.2004.06.010
29. Parvin K., Ahamed K.U., Islam M.M., Haque M.N., 2015. Response of tomato plant under salt stress: role of exogenous calcium. *Journal of Plant Sciences*. 10(6), 222-233. doi: 10.3923/jps.2015.222.233
30. Ahmad P., Abd Allah E.F., Alyemeni M.N., Wijaya L., Alam P., Bhardwaj R., Siddique K.H.M., 2018. Exogenous application of calcium to 24-epibrassinosteroid pre-treated tomato seedlings mitigates NaCl toxicity by modifying ascorbate-glutathione cycle and secondary metabolites. *Sci Rep*. 8(1), 13515. doi: 10.1038/s41598-018-31917-1
31. Tanveer K., Gilani S., Hussain Z., Ishaq R., Adeel M., Ilyas N., 2020. Effect of salt stress on tomato plant and the role of calcium. *Journal of Plant Nutrition*. 43(1), 28-35. <https://doi.org/10.1080/01904167.2019.1659324>
32. Jahani S., Lahouti M., Jahani M., 2014. Investigation Na<sup>+</sup>-Ca<sup>2+</sup> interaction on biomass and enzymes activity of peroxidase and polyphenol oxidase in leaf of barley (*Hordeum vulgare* L.). *Crop Physiology Journal*. 5(20), 15-24. <http://cpj.iauhvaz.ac.ir/article-1-186-en.html>
33. Shariat Jafari M.H., Kafi M., Astarai A.R., 2009. Interactive effects of NaCl induced salinity, calcium and potassium on physiomorphological traits of sorghum (*Sorghum bicolor* L.). *Pakistan Journal of Botany*. 41(6), 3053-3063.
34. Sivasankaramoorthy S., 2013. Effect of supplementary calcium enhances plant growth, photosynthetic pigments and uptake of nutrient in *Oryza sativa* L. under NaCl stress. *International Journal of Chemical and Life Sciences*. 2(7), 1189-1192.
35. Elkesh A.A., Alnusaire T.S., Soliman M.H., Gawayed S., Senousy H.H., Fahad S., 2019. Calcium availability regulates antioxidant system, physio-biochemical activities and alleviates salinity stress mediated oxidative damage in soybean seedlings. *Journal of Applied Botany and Food Quality*. 92, 258-266. doi: 10.5073/JABFQ.2019.092.036
36. Cramer G.R., 2002. Sodium-calcium interactions under salinity stress. *Salinity: Environment - Plants - Molecules*. Chapter 10, 205-227. doi: 10.1007/0-306-48155-3\_10
37. Yücel C.N., Heybet E.H., 2016. Salicylic acid and calcium treatments improves wheat vigor, lipids and phenolics under high salinity. *Acta Chimica Slovenica*. 63(4), 738-746. doi: 10.17344/acsi.2016.2449
38. Yousuf P.Y., Ahmad A., Hemant Ganie A.H., Aref I.M., Iqbal M., 2015. Potassium and calcium application ameliorates growth and oxidative homeostasis in salt-stressed Indian mustard (*Brassica juncea*) plants. *Pakistan Journal of Botany*. 47(5), 1629-1639.
39. Mashela P.W., Aluvilu A., Pofu K.M., 2012. Influence of sodium chloride with and without calcium chloride on growth and productivity of chilli pepper. *International Journal of AgriScience*. 2(11), 1001-1008.
40. Tabatabaieian J., 2014. Effect of calcium nutrition on reducing the effects of salinity on tomato plant. *American Journal of Plant Nutrition and Fertilization Technology*. 4(1), 11-17. doi: 10.3923/ajpnft.2014.11.17
41. Tahjib-Ul-Arif M., Roy P.R., Al Mamun Sohag A., Afrin S., Rady M.M., Afzal Hossain M., 2018. Exogenous calcium supplementation improves salinity tolerance in *BRR1 Dhan28*; a salt-susceptible high-yielding *Oryza sativa* cultivar. *Journal of Crop Science and Biotechnology*. 21, 383-394. <https://doi.org/10.1007/s12892-018-0098-0>
42. Steduto P., Albrizio R., Giorio P., Sorrentino G., 2000. Gas-exchange response and stomatal and non-stomatal limitations to carbon assimilation of sunflower under salinity. *Environmental and Experimental Botany*. 44(3), 243-255. doi: 10.1016/s0098-8472(00)00071-x
43. Gunes A., Inal A., Alpaslan M., Eraslan F., Bagci E.G., Cicek N., 2007. Salicylic acid induced changes on some physiological parameters symptomatic for oxidative stress and mineral nutrition in maize (*Zea mays* L.) grown under salinity. *Journal of Plant Physiology*. 164(6), 728-736. doi: 10.1016/j.jplph.2005.12.009.

44. Xu D., Wang W., Gao T., Fang X., Gao X., Li J., Bu H., Mu J., 2017. Calcium alleviates decreases in photosynthesis under salt stress by enhancing antioxidant metabolism and adjusting solute accumulation in *Calligonum mongolicum*. Conservation Physiology, 5(1), cox060, <https://doi.org/10.1093/conphys/cox060>
45. Ashraf M., Foolad M., 2007. Roles of glycine betaine and proline in improving plant abiotic stress resistance. Environmental and Experimental Botany. 59(2), 206-216. doi: 10.1016/j.envexpbot.2005.12.006
46. Hayat S., Hayat Q., Alyemini M.N., Wani A.S., Pichtel J., Ahmad A., 2012. Role of proline under changing environments: a review. Plant signaling & behavior. 7(11), 1456-1466. doi: 10.4161/psb.21949
47. Jaleel C.A., Manivannan P., Sankar B., Kishorekumar A., Panneerselvam R., 2007. Calcium chloride effects on salinity-induced oxidative stress, proline metabolism and indole alkaloid accumulation in *Catharanthus roseus*. Comptes Rendus Biologies. 330(9), 674-683. doi: 10.1016/j.crv.2007.07.002
48. Rahman A., Nahar K., Hasanuzzaman M., Fujita M., 2016. Calcium supplementation improves Na<sup>+</sup>/K<sup>+</sup> ratio, antioxidant defense and glyoxalase systems in salt-stressed rice seedlings. Frontiers in Plant Science. 7, 609. <https://doi.org/10.3389/fpls.2016.00609>
49. Manivannan P., Jaleel C.A., Sankar B., Somasundaram R., Murali P.V., Sridharan R., Panneerselvam R., 2007. Salt stress mitigation by calcium chloride in *Vigna radiata* L. Wilczek. Acta Biologica Cracoviensia Series Botanica. 49(2), 105-109.
50. Girija C., Smith B.N., Swamy P.M., 2002. Interactive effects of sodium chloride and calcium chloride on the accumulation of proline and glycine betaine in peanut (*Arachis hypogaea* L.). Environmental and Experimental Botany. 47(1), 1-10. [https://doi.org/10.1016/S0098-8472\(01\)00096-X](https://doi.org/10.1016/S0098-8472(01)00096-X)
51. El-Samad H.M.A., Barakat N.A.M., 2013. The physiological mechanisms of calcium chloride application on broad bean plants grown under salinity stress. Journal of Ecology and the Natural Environment. 5(12), 371-377. doi: 10.5897/JENE10.089
52. Giri J., 2011. Glycine betaine and abiotic stress tolerance in plants. Plant Signaling & Behavior. 6(11), 1746-1751. doi: 10.4161/psb.6.11.17801
53. Singh M., Kumar J., Singh S., Singh V.P., Prasad S.M., 2016. Roles of osmoprotectants in improving salinity and drought tolerance in plants: a review. Reviews in Environmental Science and Bio/Technology. 14(3), 407-426.
54. Umar J., Aliyu A., Shehu K., Abubakar L., 2018. Influence of salt stress on proline and glycine betaine accumulation in tomato (*Solanum lycopersicum* L.). Journal of Horticulture and Plant Research. 1, 19-25. doi: 10.18052/www.scipress.com/JHPR.1.19
55. Hisao T., 1973. Plant responses to water stress. Annual Review of Plant Physiology. 24, 519-570.
56. Amirjani M.R., 2011. Effect of salinity stress on growth, sugar content, pigments and enzyme activity of rice. International Journal of Botany. 7(1), 73-81. doi: 10.3923/ijb.2011.73.81
57. Silva J.V., Lacerda C.F., Costa P.H.A., Filho J.E., Enéas-Filho G., Enéas G.F., Prisco J.T., 2003. Physiological responses of NaCl stressed cowpea plants grown in nutrient solution supplemented with CaCl<sub>2</sub>. Brazilian Journal of Plant Physiology. 15(2), 99-105. doi: 10.1590/S1677-04202003000200005
58. Tuna A.L., Kaya C., Ashraf M., Altunlu H., Yokas I., Yagmur B., 2007. The Effects of calcium sulphate on growth, membrane stability and nutrient uptake of tomato plants grown under salt stress. Environmental and Experimental Botany. 59(2), 173-178. doi: 10.1016/j.envexpbot.2005.12.007
59. Larbi A., Kchaou H., Gaaliche B., Gargouri K., Boulal H., Morales F., 2020. Supplementary potassium and calcium improves salt tolerance in olive plants. Scientia Horticulturae. 260, 1-10. doi: 10.1016/j.scienta.2019.108912