



Managing drought on rangelands: adaptive strategies as perceived by pastoralists in Jiroft county

Azam Khosravi Mashizi^{1*} , Mohsen Sharafatmandrad² 

¹ Associate Professor, Department of Ecological Engineering, Faculty of Natural Resources, University of Jiroft, Jiroft, Iran

² Associate Professor, Department of Ecological Engineering, Faculty of Natural Resources, University of Jiroft, Jiroft, Iran

Article Info	Abstract
<p>Article type: Research Article</p> <p>Article history: Received: November 2022 Accepted: June 2023</p> <p>Corresponding author: Azam.khosravi@ujiroft.ac.ir</p> <p>Keywords: Arid land Ecosystem Drought Shrubland</p>	<p>Drought poses a significant challenge to pastoralists in arid and semi-arid regions, necessitating a reassessment of drought management strategies on rangelands. This study aimed to identify optimal strategies for drought management on rangelands based on the perceptions of nomadic and rural pastoralists in Jiroft County, Kerman Province. Data were collected through the administration of questionnaires. The findings, determined through multiple regression analysis, underscore the utmost significance of specific management strategies for pastoralists during drought conditions. Notably, water management ($P < 0.0001$), forage management ($P < 0.001$), and income diversification ($P < 0.05$) emerged as crucial elements in effective drought management. Principal Component Analysis (PCA) further revealed that employing suitable irrigation techniques, engaging in recreational activities, purchasing forage, and efficiently storing water are critical sub-criteria for enhancing drought management strategies. Nomadic pastoralists identified moving livestock to rangelands with better conditions and changing livestock types as important drought management strategies. Given the reluctance of rural and nomadic pastoralists to reduce their livestock numbers during drought periods, the implementation of such strategies requires governmental arrangements and financial support. The findings emphasize pastoralists' preference for reactive drought management methods over preventive strategies, underscoring the need to raise awareness among pastoral communities about the risks associated with drought. Overall, the indigenous knowledge held by pastoralists has the potential to enhance existing management plans aimed at mitigating the consequences of drought.</p>

Cite this article: Khosravi Mashizi, Azam; Sharafatmandrad, Mohsen. 2023. Managing drought on rangelands: Adaptive strategies as perceived by pastoralists in Jiroft County. *Environmental Resources Research*, 11(2), 167-178.



© The Author(s).

DOI: 10.22069/ijerr.2024.20994.1396

Publisher: Gorgan University of Agricultural Sciences and Natural Resources

Introduction

Drought is a prevalent and recurring natural phenomenon, manifesting in various climates. Regions characterized by arid and semi-arid conditions, which exhibit significant fluctuations in rainfall, are particularly susceptible to drought (Karrous & Elmouidm, 2008). The repercussions of drought span across multiple domains, including social, economic, and environmental aspects, and these consequences can differ from one location to another (Nafarzadegan et al., 2012), contingent on the specific region (Brown et al., 2017).

Drought can have detrimental effects on natural resources, particularly rangelands. Macon et al. (2016) showed that drought reduced rangeland forage production by 80% in California. Rangelands cover a significant part of Iran. These ecosystems are generally fragile lands prone to degradation and drought. Hence, recognizing the consequences of drought is the first step to cope with it in rangelands. Padiab et al. (2009) investigated the impacts of drought on rangelands around Hamoon wetland in Sistan region and reported that 200,000 hectares of rangelands around the wetland and 15,000 hectares of rehabilitated reeds in the wetland bed have experienced adverse effects due to decreased rainfall and reduced water inflow to the lake in 2000-2001.

As drought may cause severe damage to the human communities, rangeland managers seek solutions to help mitigate the impacts of drought based on the ecological, social and economic impacts of the past droughts (Kelley et al., 2016). Drought management programs should aim to mitigate drought impacts, adapt to drought variability, and recover from drought (Moghaddas Farimani et al. 2017). Drought management is a crucial tool for minimizing drought risk and expediting drought recovery. Many drought management programs have failed due to their neglect of the interactions between humans and the environment (Urquijo-Reguera, 2022). Local stakeholders as the main exploiters of ecosystems can play important role to help mitigate the impacts of drought on ecosystems (Singh and Chudasama, 2017). The pastoralists all

around the world have endured recurrent drought which negatively affect their economy from rangelands (Smart et al., 2021). Accumulated financial losses and psychological stress caused by drought affect the emotional and social dynamism of pastoralists' families (Wilmer and Fernández-Giménez, 2016).

As frequency and severity of future droughts are largely unpredictable, awareness of pastoralists' flexibility is one of the key features of drought management (Fazey et al., 2010). Pastoralism is highly dynamic and complex (Rhoades et al., 2014) and their perceptions of drought adaptation strategies are different. They choose different strategies based on their needs and goals (Wilmer and Fernández-Giménez, 2015). According to Wilmer et al. (2016), effective drought management should take into account the perceptions of those directly affected, and successful drought management necessitates learning from individuals on the 'front line' of rangeland drought who have endured recurrent drought. Wilmer and Fernández-Giménez (2015) also examined the US rangeland stakeholders' attitude on drought management and concluded that future plans are likely to be more successful if they are designed based on the complex experiences of pastoralists. Moghaddas Farimani et al. (2017) introduced the most effective drought coping strategies based on pastoralists' experiences in Fars province, Iran, as reducing stocking rates and gradually phasing out inefficient, old, and sick livestock. As drought stands as one of the most existential natural disasters threatening pastoralists in arid and semi-arid regions, it becomes imperative to reevaluate managerial strategies for drought on rangelands. While drought is a complex phenomenon, this study aims to explore the most effective drought management strategies by tapping into the experiences and perceptions of rural and nomadic pastoralists who have endured prolonged drought in their rangelands.

Materials and methods

Study Area

This study was conducted in Jiroft County, Kerman Province, south east of Iran (57°01' to 57°35' E and 28°40' to 29° 21' N, Figure 1). The average annual rainfall varies between 136 to 466 mm. The region has

experienced frequent droughts. In recent years, wet spells have decreased and dry

spells have increased (Bigne and Ekhtesasi, 2013).




Figure 1. Location of the study area

Methods

This study is a descriptive-analytical research. Questionnaires were employed to collect pastoralists' opinions in two parts. The first part gathered personal information, including gender, age, income, level of education, and employment status. The second part assessed the importance of drought management strategies using a five-point Likert scale ranging from 1=very low to 5 = very high. Criteria and sub-criteria for drought management were determined based on previous studies (Table 1). The questionnaire's validity was confirmed by university professors and experts, and necessary corrections were made in multiple stages. Cronbach's alpha was used to observe the principles and techniques of work and measure the degree of questionnaire reliability.

The required sample size was determined based on the following equation:

$$n = \frac{Z^2 p(1 - p)}{d^2}$$

where n is the sample size, Z is the statistic corresponding to level of confidence, P is the

expected prevalence that can be obtained from same studies or a pilot study conducted by the researchers, and d is precision (corresponding to effect size). Thus, 100 local adults with sufficient experience in pastoralism were randomly selected for interview.

Data Analysis

Multiple regression tests and Principal Component Analysis (PCA) were used to analyze the data and determine the most important criteria and sub-criteria of drought management (Curz-Cardenas et al., 2014). The multiple regression tests showed the contribution of drought management criteria (forage management, grazing management, flock management, water management and income diversification during drought) (Kalantari, 2010). Before running the multiple regression tests, Kolmogorov–Smirnov normality test was used to check data for normal distribution. IBM SPSS Statistics V22.0 was used for both descriptive and analytical data analysis.

Table 1. Criteria and sub-criteria for drought management

Criteria	Sub-criteria
Forage management (Treydte et al., 2017)	Sowing of forage plants (Guo et al., 2022)
	Planting of drought tolerant species for rangeland rehabilitation (Abu-Zanat et al., 2020)
	Planting of fast growing species for rangeland rehabilitation (Abu-Zanat et al., 2020)
	Forage purchase (Vrieling et al., 2016)
	Fodder storage in years with normal rainfall (Vrieling et al., 2016)
Grazing management (Liebig et al., 2014)	Conservation of species diversity (Wen et al., 2021)
	Reducing grazing time (Liebig et al., 2014)
	Implementing rotational grazing (Hughes, 2017)
	Implementing rangeland rest (Moghaddas Farimani et al., 2017)
Flock management (Mengistu, 2016)	Moving livestock to rangelands with better condition
	Reducing livestock numbers (Moghaddas Farimani et al., 2017)
	Changing livestock type (Mengistu, 2016)
Income diversification (Aliyar et al., 2022)	Breeding yearlings (Haigh et al., 2019)
	Keeping livestock of various breeds (Mengistu, 2016)
	Developing recreation (Bakhtiari et al., 2018)
Water management (Rahaman et al., 2019)	Developing hunting (Kachergis et al., 2014)
	Developing handicrafts (Aliyar et al., 2022)
	Controlling water loss (Singh et al., 2021)
	Harvesting rainwater and snowmelt (Rahaman et al., 2019)
	Promoting and teaching the correct techniques of water distribution and consumption (Lankford et al., 2020)
	Identifying and making proper use of other available sources for water required by livestock (Moghaddas Farimani et al., 2017)

Results

The personal information of respondents is summarized in Table 2. Approximately 71% of respondents were male, while 29% were female. The majority of respondents (94%) had a high school education or less, with six percent holding a 4-year college degree. In terms of age distribution, 19% of respondents were aged 30-40, 33% were aged 40-50, 35% were aged 50-60, and 13%

were aged over 60 years. Among the respondents, 56% identified as rural pastoralists, while the remaining 44% identified as nomadic pastoralists. The income of 59% of responders was less than 5,000,000 Rials per month. Additionally, 36% reported an income ranging from 5,000,000 to 10,000,000 Rials per month, and 11% reported an income exceeding 10,000,000 Rials per month.

Table 2. Personal information of pastoralists participated in the study.

Variable	Value	Frequency	Frequency (%)
Age	30-40	25	19
	40-50	44	33
	50-60	46	35
	60<	18	13
Income (Rials/month)	<5000000	44	33
	5000000-10000000	74	56
	10000000<	11	11
Education	Less than high school	79	59
	High school	46	35
	4year college degree	8	6
Gender	Female	38	29
	Male	95	71
Pastoralism type	Rural	75	56
	Nomadic	58	44

Table 3. Mean importance of each of the sub-criteria for drought management

Sub-criteria	Rural		Nomadic	
	Mean	SD	Mean	SD
Planting of forage plants	3.97	1.2	3.04	0.98
Planting of drought tolerant species for rangeland rehabilitation	2.46	0.78	2.37	0.75
Planting of fast growing species for rangeland rehabilitation	3.89	0.78	3.24	0.67
Forage purchase	4.12	0.46	4.65	0.89
Fodder storage in years with normal rainfall	2.23	0.32	1.38	0.38
Conservation of species diversity	2.38	0.98	2.86	0.67
Reducing grazing time	3.12	0.52	3.05	1.24
Implementing rotational grazing	3.89	0.67	3.24	1.23
Implementing rangeland resting	2.35	0.34	2.53	0.49
Moving livestock to rangelands with better condition	2.38	0.68	4.06	0.35
Reducing livestock numbers	1.61	1.20	1.59	0.36
Changing livestock type	2.24	0.57	2.35	0.67
Breeding yearlings	3.25	0.56	2.34	0.31
Keeping livestock of various breeds	3.37	1.20	3.56	0.97
Developing recreation	4.38	0.37	3.68	0.65
Developing hunting	2.38	1.23	2.98	0.66
Developing handicrafts	3.58	0.38	3.84	0.90
Controlling water loss	3.98	0.32	4.12	0.12
Harvesting rainwater and snowmelt	4.86	0.03	4.03	0.58
Promoting and teaching the correct techniques of water distribution and consumption	4.21	0.17	3.15	1.30
Identifying and making proper use of other available sources for livestock water	3.24	1.23	4.65	0.17

Forage purchase was the most important sub-criteria of forage management for rural and nomadic pastoralists with mean values of 0.46 ± 0.46 and 4.65 ± 0.89 respectively. Implementing rotational grazing (3.89 ± 0.67) and moving livestock to rangelands with better condition (4.12 ± 0.46) were the most important sub-criterion for grazing management based on rural and nomadic pastoralists respectively. Among the flock management sub-criteria, keeping livestock of various breeds held the highest importance for both rural and nomadic pastoralists, with mean values of 3.37 ± 1.20 and 3.56 ± 0.97 , respectively. The most important sub-criteria of income diversification were developing recreation for rural pastoralists with mean value of 3.38 ± 0.37 and developing handicrafts for nomadic pastoralists with mean value of 3.84 ± 0.90 . The most important sub-criteria

of water management were harvesting rainwater and snowmelt for rural pastoralists with mean value of 4.86 ± 0.03 and identification and proper use of other available sources for livestock water for nomadic pastoralists with mean value of 4.65 ± 0.17 (Table 3).

The PCA results revealed that, for rural pastoralists, promoting and teaching the correct techniques of water distribution and consumption, developing recreation, forage purchase, and harvesting rainwater and snowmelt were the most important sub-criteria of drought management. In contrast, for nomadic pastoralists, the key sub-criteria included identifying and making proper use of other available sources for livestock water, developing recreation, forage purchase, moving livestock to rangelands with better conditions, and changing livestock type (Table 4).

Table 4. The most important sub-criteria for drought management based on PCA

Sub-criteria	Rural		Nomadic	
	PCA1	PCA2	PCA1	PCA2
Planting of forage plants	0.202	0.038	0.213	0.029
Planting of drought tolerant species for rangeland rehabilitation	0.113	0.096	0.130	0.077
Planting of fast growing species for rangeland rehabilitation	0.134	0.137	0.239	0.067
Forage purchase	0.389	0.195	0.459	0.485
Fodder storage in years with normal rainfall	0.032	0.021	0.035	0.013
Conservation of species diversity	0.123	0.312	0.132	0.049
Reducing grazing time	0.032	0.037	0.035	0.064
Implementing rotational grazing	0.074	0.018	0.038	0.087
Implementing rangeland resting	0.031	0.237	0.031	0.019
Moving livestock to rangelands with better condition	0.037	0.234	0.358	0.438
Reducing livestock numbers	0.098	0.138	0.068	0.087
Changing livestock type	0.123	0.023	0.238	0.758
Breeding yearlings	0.137	0.038	0.237	0.035
Keeping livestock of various breeds	0.139	0.223	0.133	0.012
Developing recreation	0.454	0.213	0.432	0.234
Developing hunting	0.221	0.132	0.234	0.123
Developing handicrafts	0.237	0.231	0.239	0.137
Controlling water loss	0.298	0.138	0.492	0.123
Harvesting rainwater and snowmelt	0.192	0.438	0.253	0.035
Promoting and teaching the correct techniques of water distribution and consumption	0.447	0.135	0.231	0.137
Identifying and making proper use of other available sources for livestock water	0.237	0.190	0.482	0.121
Eigenvalue	8.96	1.27	8.93	1.46
Variance	64.06	9.13	63.08	10.48

According to the results, all five management criteria were found to be effective in drought management, with a coefficient of determination (R^2) of 84% at 99% confidence level. Additionally, the multiple correlation coefficient was 92% based on rural pastoralists (see Table 5). According to Table 6, all five independent variables (forage management, flock management, grazing management, water management and income diversification) were effective in predicting the dependent

variable (drought management) at 99% confidence level. Standardized (regression) coefficient (Beta coefficient) was used to assess the importance and role of independent variables in predicting the dependent variable. Based on rural pastoralists, water management had a greater contribution than other variables in predicting drought management. Hence, drought management can increase by 0.38 units with 1-unit increase in water management.

Table 5. The relationship between drought management criteria based on multiple regression for rural pastoralists.

	df	Sum of squares	F	sig	R^2	R
Regression	4	2.490	3.952	0.001	0.84	0.92
Residual	129	0.63				
Total	133					

Table 6. The relative importance of variables and their impact on drought management for rural pastoralists.

Drought management criteria	Standard B	t	p-value
Forage management	0.365	4.35	0.001
Grazing management	0.265	3.02	0.03
Flock management	0.213	2.86	0.04
Income diversification	0.331	3.98	0.001
Water management	0.386	5.32	0.000

According to the results, all five management criteria were effective on drought management, so that the coefficient of determination (R^2) was 84% at 99% confidence level. Also in this analysis, the rate of multiple correlation coefficient was 92% based on nomadic pastoralists (Table 7). According to Table 8, all five independent variables (forage management, flock management, grazing management,

water management, and income diversification) were statistically significant predictors of the dependent variable (drought management) at 95% confidence level. Among rural pastoralists, the water management variable had the most substantial contribution to predicting drought management. Hence, drought management can increase by 0.36 units with 1-unit increase in water management.

Table 7. The relationship between criteria based on multiple regression for nomadic pastoralists

	df	Sum of squares	F	sig	R2	R
Regression	4	3.856	6.76	0.000	0.90	0.95
Residual	129	0.57				
Total	133					

Table 8. The relative importance of variables and their impact on drought management for nomadic pastoralists

Drought management criteria	Standard B	t	p-value
Forage management	0.350	5.62	0.001
Grazing management	0.219	3.62	0.04
Flock management	0.313	4.86	0.01
Income diversification	0.281	3.58	0.04
Water management	0.360	6.32	0.000

Discussion

Our results showed that water management, forage management and income diversification were the most important strategies for drought management. Safaei et al. (2020) emphasized on the importance of water management during drought in arid and semi-arid ecosystems. Moghaddas Farimani (2017) also showed that fodder scarcity during drought is one of the most important challenges for pastoralists in rangeland. Flock management was also the most important management strategies during drought for nomadic pastoralists. Since goats tolerate low forage rangelands more than other livestock types, pastoralists can increase the proportion of goats in the

herd in drought years (Zhang et al., 2013). The reduction or absence of rainfall, diminished hydrological flows, and declining soil moisture levels are the most significant indicators of drought, with far-reaching economic and social impacts on communities (Nafarzadegan et al., 2012). Since, the highest water consumption is in agriculture section in Iran, water management was more important for the rural pastoralists than nomadic ones. Drought can reduce the sustainable production of natural ecosystems such as rangelands. Precipitation is the most important climatic factor for plant growth and soil moisture (Engda et al., 2016). Therefore, precipitation is the main limiting

factor of forage production. The findings from Noori et al. (2010) revealed a significant relationship between rangeland production and precipitation, with the lowest rangeland production occurring in years with low precipitation. Rangelands are composed of diverse plant species with different water requirements (Karimi et al. 2015). Therefore, plants' dependence on rainfall is different based on their growth form, root system as well as the temporal and spatial distribution of precipitation. Determining grazing capacity based on forage supply in arid rangelands is hard because of fluctuations in annual forage production (Moghaddam, 1998). In general, pastoralists encounter three crises related to forage shortages during drought: (1) a shortage and decline in rangeland forage or cultivated forage, (2) a reduction in forage storage for autumn and winter, and (3) the impacts of drought on crop storage and diminished plant growth in the subsequent growing season (Kachergis et al., 2014). Pastoralists also consider income diversification and reducing dependence on rangelands as important management strategies. Derner et al. (2014) and Kachergis et al. (2014) concluded that income diversification is a climate-independent strategy that can be employed to address drought. For nomadic pastoralists, the decrease in forage production and financial pressure are the most significant consequences of drought (Bagheri et al., 2012). Therefore, purchasing forage may prevent the reduction of livestock production and income.

Promoting and teaching correct techniques for water distribution and consumption, developing recreation, purchasing forage, and harvesting rainwater and snowmelt are the most crucial sub-criteria for drought management among rural pastoralists. Tingsanchali and Piriyaawong (2018) assessed drought risk based on surface water irrigation systems and demonstrated that 83 billion cubic meters of the total renewable water resources are annually consumed in the agriculture sector, with 65% lost due to the use of traditional and incorrect irrigation methods (Mamoudian, 2008; Aryanfar et al., 2020). Adopting modern irrigation systems

not only alleviates plant stress but also provides the required amount of water with minimal loss, thereby improving agricultural growth and development (Ghasemizadeh and Salemi, 2009). Thomas et al. (2013) emphasized that social and organizational networks, along with improved communication mechanisms with the media and the public, can significantly enhance adaptive capacity and result in fewer economic losses during drought. Recreation development serves as a viable source of income for villagers (Zhang et al., 2020) and can enhance the livelihoods of local people while reducing income dependence on rangelands. However, previous studies indicate a decrease in tourist numbers during drought (Bakhtiari et al., 2018), as drought can diminish the aesthetic value of natural landscapes by depleting water resources and damaging vegetation. Furthermore, the sale of handicrafts may decline due to reductions in agricultural and livestock production.

Nomadic pastoralists regard moving livestock to rangelands with better conditions and changing livestock types as crucial strategies for drought management. These findings align with the results of Mazaheri and Safari (2009) and Derner et al. (2014). Altering livestock composition, preventing livestock grazing in degraded rangelands with lower production potential, and relocating livestock to high-potential rangelands are essential aspects of drought management (McAllister et al., 2009, Coppock, 2011). Rangeland size can influence flock management strategies, with larger rangelands offering ecological and economic benefits such as forage heterogeneity and increased capital and forage storage (McAllister et al., 2009). However, previous studies have indicated that adaptive grazing management is one of the best strategies for drought management due to the temporal and spatial heterogeneity of forage (Derner et al., 2016).

Conclusion

In this study, it was observed that rural and nomadic pastoralists showed limited interest in altering the number of their livestock. Implementing such a strategy necessitates financial assistance from the government,

possibly through insurance or incentive programs. Additionally, both rural and nomadic pastoralists placed the least emphasis on preventive practices for drought, such as fodder storage during years with normal rainfall. The results of Kachergis et al. (2014) showed that most pastoralists use reactive drought management techniques such as forage purchase and fewer pastoralists use prevention strategies defined in their drought management plans such as forage stock. Given the destructive impacts of drought on ecosystems, pastoralists need to be prepared to cope with drought. Therefore, it is essential to educate young pastoralists about

drought risks, with a focus on preventive practices. In general, this study suggests that leveraging the indigenous knowledge of pastoralists could enhance existing management plans for the development of drought coping strategies. Various strategies must be implemented for drought management because what may be the 'best' management strategy during one drought may not be the most effective strategy in the next drought (McAllister et al., 2009). Therefore, operations with a broader range of management options during a drought may possess greater resilience to withstand drought conditions.

References

- Abu-Zanat, M.M.W., Al-Ghathithi, A.K., and Akash, M.W. 2020. Effect of Planting Atriplex seedlings in micro-catchments on attributes of natural vegetation in arid rangelands. *Journal of Arid Environments*. 180, 104199.
- Aliyar, Q., Zulfiqar, F., Datta, A., Kuwornu, J.K.M., and Shrestha, S. 2022. Drought perception and field-level adaptation strategies of farming households in drought-prone areas of Afghanistan. *International Journal of Disaster Risk Reduction*. 72, 102862.
- Aryanfar, Y. 2020. A review on the water sector in Iran: Current forecasts, scenario and sustainability issue. *International Journal of Progressive Sciences and Technologies*. 22, 13-18.
- Bagheri, M., Zibaei, M., Esmaili, A.K., 2012. Long Term Evaluation of Livestock Management Strategies in Drought Conditions: Case Study of Nomads in Fars Province. *Agriculture Economic Research*. 15, 113-142.
- Bigne, S., and Ekhtesasi, M.R. 2013. Investigation of the effects of drought in Jiroft plain using SPI index, the first national conference on environment, energy and bio-defense, Tehran.
- Brown, J., Alvarez, P., Byrd, K., Deswood, H., Elias, E., and Spiegel, S. 2017. Coping with historic drought in California rangelands: Developing a more effective institutional response. *Rangelands*, 39:73-78.
- Cochran, W. G. 1977. *Sampling techniques* (3rd ed.). New York: John Wiley & Sons.
- Coppock, D.L., 2011. Ranching and multiyear droughts in Utah: production impacts, risk perceptions, and changes in preparedness. *Rangeland Ecology and Management*. 64, 607-618.
- Curz-Cardenas, G., Lopez-Mata, L., Villasenor, J.L., and Ortiz, E. 2014. Potential species distribution modeling and the use of principal component analysis as predictor variables. *Rev. Mex. Biodivers.* 85, 189-199.
- Derner, J.D., and Augustine, D.J. 2016. Adaptive management for drought on rangelands. *Rangelands*. 38, 211-215.
- Engda, T.A., Kelleners, T.J., Paige, G.B., and Hild, A.L. 2016. Rainfall, evapotranspiration, and soil moisture as herbage production predictors for Wyoming rangelands. *Arid Land Research and Management*. 30, 445-459.
- Ghasemizadeh, A., and Salemi, H. 1388. Changing the method of surface irrigation to drip of old trees of model apple fruit Suitable for dealing with drought and water shortage crisis. *The Second National Conference on Drought Impacts and Solutions Manage it*. Ahwaz. Chamran University. 5 p.
- Guo, H., Feng, L., Wu, Y., Wang, J., and Liang, Q. 2022. Assessment of smallholders' vulnerability to drought based on household-scale planting strategies and adaptability: A survey study of Xinghe County. *International Journal of Disaster Risk Reduction*. 72, 102820.
- Haigh, T.R., Schacht, W., Knutson, C.L., Smart, A.J., Volesky, J., Allen, C., Hayes, M., and Burbach, M. 2019. Socioecological determinants of drought impacts and coping strategies for ranching operations in the Great Plains. *Rangeland Ecology and Management*. 72, 561-571.
- Havstad, K.M., Peters, D.P., Skaggs, R., Brown, J., Bestelmeyer, B., Fredrickson, E., Herrick, J., and Wright, J. 2007. Ecological services to and from rangelands of the United States. *Ecological Economics*. 64(2), 261-268.

- Hughes, L.E. 2017. Plant species diversity, drought, and a grazing system on the Arizona strip. *Rangelands*. 39, 20-27.
- Kachergis, E., Derner, J.D., Cutts, B.B., Roche, L.M., Eviner, V.T., Lubell, M.N., and Tate, K.W. 2014. Increasing flexibility in rangeland management during drought. *Ecosphere*. 5, 1-14.
- Kachergis, E., Derner, J.D., Cutts, B.B., Roche, L.M., Eviner, V.T., Lubell, M.N., and Tate, K.W. 2014. Increasing flexibility in rangeland management during drought. *Ecosphere*. 5, 1-14.
- Karimi, Gh., Yeganeh, H., Abbasi Khaleki, M., Moammeri, M., and Afra, H. 2015. Changes in *Bromus tomentellus* Boiss production and consumption in Kurdan rangelands, Alborz. *Iranian Journal of Natural Resources (Journal of Range & Watershed Management)*. 68 (2), 359-370.
- Karrou, M., and Elmouidm, M. 2008. Drought management and planning strategies in semi-arid and arid agro-pastoralist systems of West Asia and North Africa: A review. In: & Lopez- Francos A, editor. *Drought management: scientific and technological innovations*. Zaragoza: CIHEAM. p. 179-184
- Kelley, W.K., DerekScasta, J., and Derner, J.D. 2016. Advancing knowledge for proactive drought planning and enhancing adaptive management for drought on rangelands: Introduction to a Special Issue. *Rangelands*. 38(4), 159-161.
- Lankford, B., Closas, A., Dalton, J., Gunn, E.L., Hess, T., Knox, J.W., Der Kooij, S., and Lautze, J. 2020. A scale-based framework to understand the promises, pitfalls and paradoxes of irrigation efficiency to meet major water challenges. *Global Environmental Change*. 65, 102182.
- Liebig, M.A., Kronberg, S.L., Hendrickson, J.R., and Gross, J.R. 2014. Grazing management, season, and drought contributions to near-surface soil property dynamics in semiarid rangeland. *Rangeland Ecology and Management*, 67(3), 266-274.
- Macon, D.K., Barry, S., Beccetti, T., Davy, J.S., Doran, M.P., Finzel, J.A., George, H., Harper, J.M., Hunt-Singer, L., Ingram, R.S., and Lancaster, D.E. 2016. Coping with drought on California rangelands. *Rangelands*. 38, 222-228.
- Mamoudian, S.A., 2008. Secretary of the IWA National Committee for Iran: "Iran. Water and wastewater management across the country", IWA Yearbook, p. 28
- Mazaheri, Z., and Safari, N. 2009. A study of the effects of drought on the life of a nomadic community: A case study of Chaharmahal and Bakhtiari province, the second national conference on the effects of drought and its management strategies, Isfahan.
- McAllister, R. R. J., Stafford Smith, D. M., Stokes, C. J., and Walsh, F. J. 2009. Patterns of accessing variable resources across time and space: Desert plants, animals and people. *Journal of Arid Environment*. 73, 338-346.
- Mengistu, D. 2016. Impacts of drought and conventional coping strategies of Borana Community, Southern Ethiopia. *Research Journal on Humanities Social Sciences*. 6(23), 37.
- Moghaddam, M., 1998. Range and range management. Tehran University Press, 470 pp.
- MoghaddasFarimani, Sh., Raufirad, V., Hunter, R., and Lebailly, Ph. 2017. Coping strategies during drought: The case of rangeland users in Southwest Iran. *Rangelands*. 39, 133-142.
- Nafarzadegan, A.R., Rezaeian Zadeh, M., Kherad, M., Ahani, H., Gharekhani, A., Karampoor, M.A., Kousari, M.R. 2012. Drought area monitoring during the past three decades in Fars province, Iran. *Quat International*. 250, 27-36.
- Noori, Gh., Khosravi, M., Javdani, R. and Karimi, S. 2010. Determining the relationship between drought and variability in pasture production in Sistan and Baluchestan province during the statistical period of 1370-1386 (Case study: Iranshahr city). *The 4th International Congress of the Islamic World Geographers (ICIWG 2010)*, pp. 1-11
- Padiab, M., Ghasemi Arian, Y., and Razavi Moghadam, M. 2009. Investigating the impact of drought on Sistan rangeland Appearance, National Conference on Water Crisis in Agriculture and Natural Resources, Tehran.
- Ppplley, H.W., Briske, D.D., Morgan, J.A., Wolter, K., Bailey, D.W., and Brown, J.R. 2013. Climate change and North American rangelands: trends, projections, and implications. *Rangeland Ecology and Management*. 66,493-511
- Rahaman, Md. F., Jahan, Ch.S., and Mazumder, Q.H. 2019. Rainwater harvesting: Practiced potential for Integrated Water Resource Management in drought-prone Barind Tract, Bangladesh. *Groundwater for Sustainable Development*. 9, 100267.
- Safaei, A., Choramin, M., and Korkani, K. 2020. Using several drought indices in Water resources management in drought condition (Case Study of Shahid Abbaspour Dam Station). *Water Engineering*. 8, 1-10

- Singh, Ch., Jain, G., Sukhwani, V., and Shaw, R. 2021. Losses and damages associated with slow-onset events: urban drought and water insecurity in Asia. *Current Opinion in Environmental Sustainability*. 50, 72-86.
- Singh, P.K., and Chudasama, H. 2017. Pathways for drought resilient livelihoods based on people's perception. *Climatic Change*. 140, 179–193.
- Smart, A.J., Harmoney, K., Scasta, D., Stephenson, M.B., Volesky, J.D., Vermeire, L.T., Mosley, J.C., Sedivec, K., Meehan, M., Haigh, T., Derner, J.D., and McClaran, M.P. 2021. Forum: Critical Decision Dates for Drought Management in Central and Northern Great Plains Rangelands. *Rangeland Ecology and Management*. 78, 191-200.
- Smith, A., and Katz, R. 2013. U.S. Billion-dollar weather and climate disasters: data sources, trends, accuracy and biases. *Natural Hazards*. 6(2), 387-410.
- Tingsanchali, T., and Piriya Wong, Th. 2018. Drought risk assessment of irrigation project areas in a river basin. *Engineering Journal*. 22, 279-287.
- Treydte, A.C., Schmiedgen, A., Berhane, G., and Tarekegn, K.D. 2017. Rangeland forage availability and management in times of drought – A case study of pastoralists in Afar, Ethiopia. *Journal of Arid Environments*. 139, 67-75.
- Urquijo-Reguera, J., Gómez-Villarino, M.T., Pereira, D., and De Stefano, L. 2022. An assessment framework to analyze drought management plans: The case of Spain. *Agronomy*. 12(4), 970. <https://doi.org/10.3390/agronomy12040970>
- Vrieling, A., Meroni, M., Mude, A.G., Chantarat, S., Ummenhofer, C.C., and de Bie, K. 2016. Early assessment of seasonal forage availability for mitigating the impact of drought on East African pastoralists. *Remote Sensing of Environment*. 174, 44-55.
- Wen, Zh., Zheng, H., Smith, J.R., and Ouyang, Zh. 2021. Plant functional diversity mediates indirect effects of land-use intensity on soil water conservation in the dry season of tropical areas. *Forest Ecology and Management*. 480, 118646.
- Wilmer, H., York, E., Kelley, W.K., and Brunson, M.W. 2016. "In Every Ranger's Mind": Effects of Drought on Ranch Planning and Practice. *Rangelands*. 38, 216-221.
- Wilmer, H., and Fernández-Giménez, M.E. 2015. Rethinking ranger decision-making: a grounded theory of ranching approaches to drought and succession management. *The Rangeland Journal*. 37, 517-528.
- Zhang, C., Wenjun, L., and Mingming, F. 2013. Adaptation of herders to droughts and privatization of rangeland-use rights in the arid Alxa Left Banner of Inner Mongolia. *Journal Environment Management*. 126, 182-190.
- Zhang, K., Sun, X., Jin, Y., Liu, J., Wang, R., and Zhang, Sh. 2020. Development models matter to the mutual growth of ecosystem services and household incomes in developing rural neighborhoods. *Ecological Indicators*. 115, 106363.

