

JENSEN'S INEQUALITY AND tgs-CONVEX FUNCTIONS WITH APPLICATIONS

H. Barsam [™], Y. Sayyari[™], and S. Mirzadeh[™]

Article type: Research Article

(Received: 25 July 2022, Received in revised form 16 January 2023) (Accepted: 31 January 2023, Published Online: 01 February 2023)

ABSTRACT. In recent years, many researches have been done on the tgsconvex functions and their applications. In this article, we present some properties of the tgs-convex functions by interesting examples. Then we investigate the non-positive property of the tgs-convex functions. Also, we derive types of the Jensen's inequality for the tgs-convex functions and obtain several inequalities with respect to the Jensen's inequality. Finally, we give some applications of these inequalities.

Keywords: Jensen's inequality, tqs-convex function, Global bounds. 2020 MSC: Primary 26A51, 26B25, 26D20. 26D15, 26D10.

1. Introduction and Preliminaries

Recently, many researchs and surveis have been published on mathematical inequalities and their applications in ergodic theory (see [11, 12, 14, 20–22]) and convex analysis (see [1, 3-8, 13, 15-19, 23, 24]). In the last few decades, mathematical inequalities and their generalization for convex functions have attracted wide attention. Convex analysis has an important role in the development of inequalities theory. Applying the convexity property of functions, researchers have extracted many inequality theories. In fact, the convexity property of functions is base of some inequalities such as the arithmetic mean, harmonic mean inequality also in inequality with respect to entropies including Shannon's inequality, Ky Fan's inequality and etc. In applied literature of mathematical inequalities, the Jensen inequality is a well-known, paramount, and extensively used inequality. This inequality is as follows: Let $f: I \longrightarrow \mathbb{R}$ be a convex function. Then the inequality

$$f(\sum_{i=1}^{n} p_i x_i) \le \sum_{i=1}^{n} p_i f(x_i)$$

holds for every convex combination $\sum_{i=1}^{n} p_i x_i$ of points $x_i \in I$. Recently, generalizations and improvements of Jensen's inequality have been considered by many researchers. It has been generalized to some functions including, sconvex, m-convex, etc. The concept of the tgs-convex functions was introduced

⊠ hasanbarsam@ujiroft.ac.ir, ORCID: 0000-0003-4487-5434 DOI: 10.22103/jmmr.2023.19962.1307

Publisher: Shahid Bahonar University of Kerman

459



How to cite: H. Barsam, Y. Sayyari, S. Mirzadeh, Jensen's inequality and tgs-convex functions with applications, J. Mahani Math. Res. 2023; 12(2): 459-469.

H. Barsam, Y. Sayyari, S. Mirzadeh

by the authors in [25]. They have obtained some results on these functions. Here we examine this definition and conclude the only non-negative tgs-convex function is zero. In fact, we prove that any tgs-convex function is non-positive. Hence, we assume that f is an arbitrary function that means that f is not necessarily non-negative. In this paper, we try to find Jensen's inequality for tgs-convex functions. Also, we obtain some inequalities with respect to Jensen's inequality with some applications.

Definition 1.1. Let $I \subseteq \mathbb{R}$ be an interval. A function $f: I \longrightarrow \mathbb{R}$ is said to be convex if the inequality

$$f(\lambda x + (1 - \lambda)y) \le \lambda f(x) + (1 - \lambda)f(y)$$

holds for every $x, y \in I$ and every $\lambda \in [0, 1]$.

The following definition for non-negative functions can be found in ([9], [16], [23], [24] [25]).

Definition 1.2. Let $f: I \subseteq \mathbb{R} \to \mathbb{R}$ be a function. The function f is called a *tgs*-convex function on I if the inequality

$$f(tx + (1 - t)y) \le t(1 - t)(f(x) + f(y))$$

holds for all $x, y \in I$ and $t \in (0, 1)$.

Notice that in this definition, the non-negativity costraint on f is removed.

Definition 1.3. Let $x_1, ..., x_n \in I$ be *n* points, and let $p_1, ..., p_n \in [0, 1]$ be *n* coefficients such that $\sum_{i=1}^n p_i = 1$. The summation $\sum_{i=1}^n p_i x_i$ is called the convex combination of points x_i (with coefficients p_i).

2. Results and proofs

First, we prove that tgs-convex functions cannot be positive.

Proposition 2.1. Let $f: I \subseteq \mathbb{R} \to \mathbb{R}$ be a tgs-convex function. Then $f \leq 0$.

Proof. Suppose that $x = y \in I$ and $t = \frac{1}{2}$. Since f is a tgs-convex, we have

$$f(\frac{1}{2}x + \frac{1}{2}x) \le \frac{1}{4}(2f(x))$$

Archive of SID.ir

that is, $f(x) \leq 0$ for every x.

Remark 2.2. Let $f: I \subseteq \mathbb{R} \to \mathbb{R}$ be a non-negative tgs-convex function. Then, f = 0.

Lemma 2.1 introduces an example of tgs-convex functions.

Lemma 2.3. For $a \ge 1$, let $f : [a,b] \to \mathbb{R}$ be a function defined by $f(x) = -\log x$. Then f is a tgs-convex function.

460

Jensen's inequality and tgs-convex functions with applications – JMMR Vol. 12, No. 2 (2023) 461

Proof. Let $x, y \in [a, b]$ and 0 < t < 1 be arbitrary. Since $a \ge 1$, we have $x, y \ge 1$. Hence, $x^{t(1-t)} \le x^t$ and $y^{t(1-t)} \le y^{1-t}$. Thus

(1)
$$(xy)^{t(1-t)} = x^{t(1-t)}y^{t(1-t)} \le x^t y^{t(1-t)} \le x^t y^{1-t}$$

On the other hand, by Young's inequality, we have

 $x^{t}y^{1-t} \le tx + (1-t)y.$

Then by Equation (2.1), we obtain

$$(xy)^{t(1-t)} \le tx + (1-t)y.$$

Hence,

$$\log((xy)^{t(1-t)}) \le \log(tx + (1-t)y),$$

we get

$$t(1-t)(\log x + \log y) \le \log(tx + (1-t)y),$$

and also,

$$-\log(tx + (1-t)y) \le t(1-t)(-\log x - \log y).$$

Archive of SID.ir

In the following, we consider the relationship between convex functions and tgs-convex functions in view of Definition 1.2.

Theorem 2.4. Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a negative convex function. Then f is a tgs-convex function.

Proof. Let $t \in [0,1]$ and $x, y \in I$. Then $t(1-t) \le t$, $t(1-t) \le 1-t$. Since $f \le 0$, we get $tf(x) \le t(1-t)f(x)$ and $(1-t)f(y) \le t(1-t)f(y)$. Hence

$$f(tx + (1 - t)y) \le tf(x) + (1 - t)f(y)$$

$$\le t(1 - t)f(x) + t(1 - t)f(y)$$

$$= t(1 - t)(f(x) + f(y)).$$

Example 2.5. The function $f(x) = -\sqrt{x}$ on $[0, +\infty)$ is tgs-convex.

Example 2.6. The function $f(x) = -\ln x$ on $[1, +\infty)$ is tys-convex.

In the following, we present a function f such that f is a tgs-convex which it is not convex.

Example 2.7. Let $f : [0, 4] \longrightarrow \mathbb{R}$ be a function defined by

$$f(x) = \begin{cases} -1 & \text{if } 0 \le x < 2\\ -2 & \text{if } 2 \le x \le 4 \end{cases}$$

Then f is tgs-convex. In fact, let $x, y \in [0, 4]$ and $t \in [0, 1]$. We have $f(tx + (1 - t)y) \le -1 \le -4t(1 - t) \le t(1 - t)(f(x) + f(y)).$

H. Barsam, Y. Sayyari, S. Mirzadeh

Note that $t(1-t) \leq \frac{1}{4}$ for each $t \in [0,1]$. The function f is not convex because for $x = 0, y = 2, t = \frac{1}{2}$ we have

$$f(1) \not\leq \frac{f(0) + f(2)}{2}.$$

Theorem 2.8. Let $f : I \to \mathbb{R}$ be a tgs-convex function and r be a root of f such that $r \in int(I)$. Then f is identically 0.

Proof. Let $x, y \in I$ be arbitrary such that x < r < y. There exists $t \in (0, 1)$ such that r = tx + (1 - t)y. Hence

$$0 = f(r) = f(tx + (1 - t)y) \le t(1 - t)(f(x) + f(y)) \le 0.$$

Thus, f(x) = f(y) = 0 (since $f \le 0$) which implies that f is zero on I.

Remark 2.9. The condition $r \in int(I)$ in Theorem 2.8 is essential, see the following example,

Example 2.10. Let $f : [0,1] \longrightarrow \mathbb{R}$ be a function defined by

$$f(x) = \begin{cases} 0 & if \ x = 0 \\ -1 & if \ 0 < x \le 1 \end{cases}$$

Then f is tgs-convex which it is not identically 0. In fact, let $x, y \in [0, 1]$ and $t \in [0, 1]$. We have

$$f(tx + (1-t)y) = -1 \le -\frac{1}{2} \le t(1-t)(f(x) + f(y)).$$

Note that $t(1-t) \leq \frac{1}{4}$ for each $t \in [0,1]$.

Remark 2.11. Let $f: I \to \mathbb{R}$ be a function, b > 0 and $-2b \le f(x) \le -b$. Then f is a tgs-convex function.

Proof. Let $x, y \in I$ and $t \in [0, 1]$. Then we have

$$f(tx + (1-t)y) \le -b \le -4bt(1-t) \le t(1-t)(f(x) + f(y)).$$

In the following, we present some results on Jensen's inequality.

Theorem 2.12. Let $f : [a,b] \to \mathbb{R}$ be a tgs-convex function and $t_i \in [0,1]$ such that $\sum_{i=1}^{n} t_i = 1$. Then

$$f(\sum_{i=1}^{n} t_i x_i) \le \frac{1}{n} \sum_{i=1}^{n} t_i (1-t_i) f(x_i).$$

Proof. Let $t_1, t_2 \in [0, 1]$ such that $t_1 + t_2 = 1$. We have

$$\begin{aligned} f(t_1x_1 + t_2x_2) &\leq t_1t_2(f(x_1) + f(x_2)) \\ &= t_1t_2f(x_1) + t_1t_2f(x_2) \\ &\leq t_1t_2f(x_1) \; (\text{and} \; \leq t_1t_2f(x_2)) \; \text{since} \; f(x) \leq 0 \\ &= t_1(1 - t_1)f(x_1) \; (\text{and} \; \leq t_2(1 - t_2)f(x_2)). \end{aligned}$$

Archive of SID.ir

Jensen's inequality and tgs-convex functions with applications – JMMR Vol. 12, No. 2 (2023) 463

So $2f(t_1x_1+t_2x_2) \leq t_1t_2(f(x_1)+f(x_2))$. Hence, $f(t_1x_1+t_2x_2) \leq \frac{1}{2}t_1t_2(f(x_1)+f(x_2))$. Now, we prove that the result holds for n. Let $\alpha_j = \sum_{k=1, k\neq j}^n t_k$. We have,

$$f(\sum_{k=1}^{n} t_k x_k) = f(\alpha_j(\sum_{k=1, k \neq j}^{n} \frac{t_k}{\alpha_j} x_k) + t_j x_j) \le t_j(1-t_j)f(x_j), \ j = 1, 2, \dots, n.$$

In fact, we prove that $f(\sum_{i=1}^{n} t_i x_i)$ is less than $t_i(1-t_i)f(x_i)$ for each $1 \le i \le n$. By summing the above inequalities, we have

$$nf(\sum_{i=1}^{n} t_i x_i) \le \sum_{i=1}^{n} t_i (1-t_i) f(x_i).$$

Since $f(\sum_{i=1}^{n} t_i x_i)$ is less than $t_i(1-t_i)f(x_i)$ for each $1 \le i \le n$, we can improve the result of Theorem 2.12 as follows:

Example 2.13. Assume that $x_1, \ldots, x_n \in \mathbb{R}$. Then we have

(1) if $x_i \ge 1$, then we

$$\sum_{i=1}^n t_i x_i \ge \prod_{i=1}^n x_i^{\frac{t_i(1-t_i)}{n}}$$

(2) if $x_i \ge 1$, then we have

$$\frac{\sum_{i=1}^{n} x_i}{n} \ge \left(\prod_{i=1}^{n} x_i\right)^{\frac{n-1}{n^3}}$$

(3) if $n \in \mathbb{N}$, then we have

$$n+1 \ge 2[n!]^{\frac{n-1}{n^3}}.$$

Proof. (1): In Theorem 2.12, put $f(x) = -\log x$. (2): In (1), put $t_i = \frac{1}{n}$. (3): In (2), put $x_i = i$.

Remark 2.14. Let $f: I \to \mathbb{R}$ be a tgs-convex function $n \ge 2$ and $t_i \in [0, 1]$ such that $\sum_{i=1}^n t_i = 1$. Then we have

$$f(\sum_{i=1}^{n} t_i x_i) \le \min_{1 \le j \le n} \{ t_j (1 - t_j) f(x_j) \}.$$

We give another relation with respect to Jensen's inequality.

Theorem 2.15. Let $f: I \to \mathbb{R}$ be a tgs-convex function $n \ge 2$ and $t_i \in [0,1]$ such that $\sum_{i=1}^{n} t_i = 1$. Then we have

Archive of SID.ir

H. Barsam, Y. Sayyari, S. Mirzadeh

Proof.

$$f(\sum_{i=1}^{n} t_i x_i) \le \prod_{i=1}^{n} t_i f(x_1) + t_1 \prod_{i=2}^{n} t_i f(x_2) + (t_1 + t_2) \prod_{i=3}^{n} t_i f(x_4) + (t_1 + t_2 + t_3) \prod_{i=4}^{n} t_i f(x_i) + \dots + (\sum_{i=1}^{n-2} t_i) t_{n-1} t_n f(x_{n-1}) + (\sum_{i=1}^{n-1} t_i) t_n f(x_n).$$

Proof by induction. For n = 2, the result is trivially held. Assume that for n - 1, the result holds. Specifically, we have

$$f(\sum_{j=1}^{n-1} t_i x_j) \le \prod_{j=1}^{n-1} t_j f(x_1) + t_1 \prod_{j=2}^{n-1} t_j f(x_2) + (t_1 + t_2) \prod_{j=3}^{n-1} t_j f(x_4) + (t_1 + t_2 + t_3) \prod_{j=4}^{n-1} t_j f(x_j) + \dots + (\sum_{j=1}^{n-2} t_j) t_{n-1} f(x_{n-1}),$$

which $\sum_{j=1}^{n-1} t_j = 1$. Now, we prove that the result holds for n. For this purpose, assume that $\sum_{i=1}^{n} t_i = 1$, $t_i \in [0,1]$ and $\alpha = \sum_{i=1}^{n-1} t_i$. We

$$f(\sum_{i=1}^{n} t_i x_i) = f(\sum_{i=1}^{n-1} t_i x_i + t_n x_n) = f(\alpha \sum_{i=1}^{n-1} \frac{t_i}{\alpha} x_i + t_n x_n)$$

$$\leq \alpha t_n (f(\sum_{i=1}^{n-1} \frac{t_i}{\alpha} x_i) + f(x_n))$$

$$= \alpha t_n f(\sum_{i=1}^{n-1} \frac{t_i}{\alpha} x_i) + \alpha t_n f(x_n)$$

$$\leq \alpha t_n [\prod_{i=1}^{n-1} \frac{t_i}{\alpha} f(x_1) + \frac{t_1}{\alpha} \prod_{i=2}^{n-1} \frac{t_i}{\alpha} f(x_2) + (\frac{t_1}{\alpha} + \frac{t_2}{\alpha}) \prod_{i=3}^{n-1} \frac{t_i}{\alpha} f(x_4)$$

Archive of SID.ir

Jensen's inequality and tgs-convex functions with applications – JMMR Vol. 12, No. 2 (2023) 465

$$+ \left(\frac{t_1}{\alpha} + \frac{t_2}{\alpha} + \frac{t_3}{\alpha}\right) \prod_{i=4}^{n-1} \frac{t_i}{\alpha} f(x_i) + \dots + \left(\sum_{i=1}^{n-2} \frac{t_i}{\alpha}\right) \frac{t_{n-1}}{\alpha} f(x_{n-1}) \right] + \alpha t_n f(x_n)$$

$$\le \frac{t_n}{\alpha^{n-2}} \prod_{i=1}^{n-1} t_i f(x_1) + \frac{t_n}{\alpha^{n-2}} t_1 \prod_{i=2}^{n-1} t_i f(x_2) + \frac{t_n}{\alpha^{n-3}} (t_1 + t_2) \prod_{i=3}^{n-1} t_i f(x_4)$$

$$+ \frac{t_n}{\alpha^{n-4}} (t_1 + t_2 + t_3) \prod_{i=4}^{n-1} t_i f(x_i) + \dots + \frac{t_n}{\alpha} (\sum_{i=1}^{n-2} t_i) t_{n-1} f(x_{n-1})$$

$$+ \left(\sum_{i=1}^{n-1} t_i\right) t_n f(x_n)$$

$$\le \prod_{i=1}^n t_i f(x_1) + t_1 \prod_{i=2}^n t_i f(x_2) + (t_1 + t_2) \prod_{i=3}^n t_i f(x_4)$$

$$+ (t_1 + t_2 + t_3) \prod_{i=4}^n t_i f(x_i) + \dots + \left(\sum_{i=1}^{n-2} t_i\right) t_{n-1} t_n f(x_{n-1})$$

$$+ \left(\sum_{i=1}^{n-1} t_i\right) t_n f(x_n).$$

Note that since $0 < \alpha \leq 1$ and $1 \leq j \leq n$, we have $\frac{1}{\alpha} \geq 1$ and $f(x) \leq 0$. We get

$$\frac{1}{\alpha^{j}} (\sum_{k=1}^{m} t_{k}) \prod_{s=m+1}^{n} t_{s} f(x_{s}) \leq (\sum_{k=1}^{m} t_{k}) \prod_{s=m+1}^{n} t_{s} f(x_{s}).$$

Example 2.16. Let $f : I \to \mathbb{R}$ be a tys-convex function and $x_1, \ldots, x_n \in \mathbb{R}$. Then

(1) we have

$$f(\frac{\sum_{i=1}^{n} x_i}{n}) \le \frac{f(x_1) + \sum_{i=2}^{n} (i-1)n^{i-2}f(x_i)}{n^n};$$

(2) if $x_i \ge 1$, then we

$$\left(\frac{\sum_{i=1}^{n} x_i}{n}\right)^{n^n} \ge x_1 \prod_{i=2}^{n} x_i^{(i-1)n^{i-2}}.$$

Proof. (1): In Theorem 2.15, put $t_i = \frac{1}{n}$. (2): In (1), put $f(x) = -\log x$.

Theorem 2.17. Let $f : [a, b] \to \mathbb{R}$ be a tgs-convex function and $t_i \in [0, 1]$ such that $\sum_{i=1}^{n} t_i = 1$. Then

$$f(\sum_{i=1}^{n} t_i x_i) \le \prod_{i=1}^{n} t_i(\sum_{i=1}^{n} f(x_i)).$$

Archive of SID.ir

H. Barsam, Y. Sayyari, S. Mirzadeh

Proof. Proof by induction. For m = 2, the result is trivially held. Assume that for m, the result holds Specifically, we have

$$f(\sum_{i=1}^{m} t_i x_i) \le \prod_{i=1}^{m} t_i(\sum_{i=1}^{m} f(x_i)),$$

which $\sum_{i=1}^{m} t_i = 1$. Now, we prove that the result holds for m + 1. For this purpose, assume that $\sum_{k=1}^{m+1} t_k = 1$, we have

$$\begin{split} f(\sum_{k=1}^{m+1} t_k x_k) &= f(\sum_{k=1}^{m-1} t_k x_k + (t_m + t_{m+1}) \frac{t_m x_m + t_{m+1} x_{m+1}}{t_m + t_{m+1}}) \\ &\leq \prod_{k=1}^{m-1} t_k (t_m + t_{m+1}) (\sum_{k=1}^{m-1} f(x_k) + f(\frac{t_m x_m + t_{m+1} x_{m+1}}{t_m + t_{m+1}})) \\ &\leq \prod_{k=1}^{m-1} t_k (t_m + t_{m+1}) (\sum_{k=1}^{m-1} f(x_k) + \frac{t_m t_{m+1}}{(t_m + t_{m+1})^2} [f(x_m) + f(x_{m+1})] \\ &= \prod_{k=1}^{m-1} t_k (t_m + t_{m+1}) (\sum_{k=1}^{m-1} f(x_k)) + \frac{1}{t_m + t_{m+1}} \prod_{k=1}^{m+1} t_k [f(x_m) + f(x_{m+1})] \end{split}$$

Note that since $f(x) \leq 0$, we have

$$(t_m + t_{m+1}) \prod_{k=1}^{m-1} t_k \ge \prod_{k=1}^{m+1} t_k,$$
$$\frac{1}{t_m + t_{m+1}} \prod_{k=1}^{m+1} t_k \ge \prod_{k=1}^{m+1} t_k.$$

3. Applications

In this section, we present an application of our results.

Proposition 3.1. Let $a_0, a_1, \ldots, a_n \in \mathbb{R}$ where $a_0 \neq 0$, and suppose that for $i = 1, 2, \ldots, n, x_i \in \mathbb{R}$ and $t_i \in [0, 1]$ such that $\sum_{i=1}^n t_i = 1$. Then we have (1) if $x_i \geq 1$, then we

$$\left(\prod_{i=1}^{n} x_i\right)^{\prod_{i=1}^{n} t_i} \le \sum_{i=1}^{n} t_i x_i;$$

(2) if $x_i \ge 1$, then we

$$\sqrt[n^n]{\prod_{i=1}^n x_i} \le \frac{\sum_{i=1}^n x_i}{n};$$

Archive of SID.ir

Jensen's inequality and tgs-convex functions with applications – JMMR Vol. 12, No. 2 (2023) 467

(3) if
$$0 < a_0 \le a_1 \le \dots \le a_n$$
 then
 $\sqrt[n^n]{\frac{a_n}{a_0}} \le \frac{\sum_{i=1}^n \frac{a_i}{a_{i-1}}}{n}$

Proof. (1): Using Lemma 2.3, it is proved that the function $f(x) = -\log x$ is tgs-convex on $[1, +\infty]$. Now, appling Theorem 2.17 for $-\log x$, we have

$$-\log(\sum_{i=1}^{n} t_i x_i) \le (\prod_{i=1}^{n} t_i)(-\sum_{i=1}^{n} \log x_i),$$

hence

$$\log(\sum_{i=1}^{n} t_i x_i) \ge (\prod_{i=1}^{n} t_i)(\sum_{i=1}^{n} \log x_i) = \log\left((\prod_{i=1}^{n} x_i)^{\prod_{i=1}^{n} t_i}\right).$$

(2): In (1), put $t_i = \frac{1}{n}$. (3): In (2), put $x_i = \frac{a_i}{a_{i-1}}$.

Archive of SID.ir

4. Conclusion

This paper investigated the tgs-convex functions. It was proven that if we consider the tgs-convex function as a non-negative function, it must be the zero function. So we conclude that any tgs-convex function is non-positive. Also, we presented three versions of Jensen inequality for tgs-convex functions.

5. Aknowledgement

We would like to thank the reviewers for their thoughtful comments and efforts towards improving our paper

References

- I. Abbas Baloch, A. A. Mughal, Y. M. Chu, et al, A variant of Jensen-type inequality and related results for harmonic convex functions, AIMS Math., 5 (2020), 6404–6418.
- [2] M. Adil Khan, M. Hanif, Z. A. Khan, et al, Association of Jensen's inequality for sconvex function with Csiszar divergence, J. Inequal. Appl., 2019 (2019), 1–14.
- [3] M. Adil Khan, S. Khan, Y.-M. Chu, A new bound for the Jensen gap with applications in information theory, IEEE Access20(2020), 98001-98008.
- [4] M. Adil Khan, J. Pecaric, Y. M. Chu, Refinements of Jensen's and McShane's inequalities with applications, AIMS Math., 5 (2020), 4931–4945.
- [5] H. Barsam, Some New Hermite-Hadamard Type Inequalities for Convex Functions. Wavelets linear algebr., 7(2)(2020), 11-22.
- [6] H. Barsam, A. R. Sattarzadeh, Hermite-Hadamard inequalities for uniformly convex functions and Its Applications in Means, Miskolc Math. Notes, 21(2) (2020), 621-630.
- [7] H. Barsam, A. R. Sattarzadeh, Some results on Hermite-Hadamard inequalities, Mahani Math. Res. Cent. 9(2) (2020), 79-86.
- [8] H. Barsam, Y. Sayyari, Jensen's inequality and m-convex functions, Wavelets linear algebr., 8(2) (2020), 43-51.
- S. S. Dragomir, A converse result for Jensen's discrete inequality via Grüss inequality and applications in information theory, An. Univ. Oradea. Fasc. Mat.7 (1999-2000), 178-189.

H. Barsam, Y. Sayyari, S. Mirzadeh

- [10] S. S. Dragomir, Bounds for the normalized Jenson functional, Bull. Austral. Math. Soc. 74 (2006), 471-478.
- [11] M. Ebrahimi, A. Mehrpooya, An application of geometry in algebra: uncertainty of hyper MV -algebras, in Proceedings of the 7th seminar on geometry, topology, Tehran (2014), 529–534
- [12] A. Mehrpooya, M. Ebrahimi, B. Davvaz, Two dissimilar approaches to dynamical systems on hyper MV-algebras and their information entropy, Eur. Phys. J. Plus, 132(2017), 379–405.
- [13] B. Feng, M.Ghafoor, Y. M. Chu, M. I. Qureshi, X. Feng, C. Yao, X. Qiao, Hermite-Hadamard and Jensen's type inequalities for modified (p, h)-convex functions, AIMS Math., 5(6) (2020), 6959–6971.
- [14] S.S. Dragomir, C.J. Goh, Some bounds on entropy measures in Information Theory, Appl. Math. Lett., 10 (3) (1997) 23-28.
- [15] S. Khan, M.A. Khan, S.I. Butt, Y.M. Chu, A new bound for the Jensen gap pertaining twice differentiable functions with applications Adv. Differ. Equ., 2020 (1) (2020), 1-11.
- [16] N. Mehreen, M. Anwar, Inequalities for tgs-convex functions via some conformable fractional integrals, Int. J. Nonlinear Anal. Appl., 12(2) (2021), 425-436.
- [17] Y.Sayyari, H. Barsam, Hermite-Hadamard type inequality for m-convex functions by using a new inequality for differentiable functions, J. Mahani Math. Res. Cent. 9(2) (2020), 55-67.
- [18] Y.Sayyari, H. Barsam, On some inequalities of differentable uniformly convex mapping with applications, Numer. Funct. Anal. Optim. (2023), (accepted).
- [19] Y. Sayyari, H. Barsam, A. R. Sattarzadeh, On New Refinement of the Jensen inequality Using Uniformly convex functions with Applications. Appl Anal.(2023). (accepted). doi.org/10.1080/00036811.2023.2171873
- [20] Y. Sayyari, New bounds for entropy of information sources, Wavelet and Linear Algebra, 7 (2) (2020), 1-9.
- [21] Y. Sayyari, A refinement of the Jensen-Simic-Mercer inequality with applications to entropy, j. korean soc. math. edu. ser. b-pure and appl. math., 29 (1) (2022), 51–57.
- [22] S. Simic, Jensen's inequality and new entropy bounds, Appl. Math. Lett. 22 (8) (2009), 1262-1265.
- [23] S. Simic, On an upper bound for Jensen's inequality, Journal of Inequalities in Pure and Applied Mathematics, 10(2), article 60, 5 pages, 2009.
- [24] S. Simic, On a global upper bound for Jensen's inequality, J. Math. Anal. Appl., 343 (1) (2008), 414-419.
- [25] M. Tunc, E. Gov, U. Sanal, On tgs-convex function and their inequalities, Facta Univ. Ser. Math. Inf. 30(5) (2015), 679-691.

Archive of SID.ir

468

Jensen's inequality and $\it tgs\mbox{-}convex$ functions with applications – JMMR Vol. 12, No. 2 (2023) 469

HASAN BARSAM ORCID NUMBER: 0000-0003-4487-5434 DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE UNIVERSITY OF JIROFT, P.O. BOX 78671-61167 JIROFT, IRAN Email address: hasanbarsam@ujiroft.ac.ir

Yamin Sayyari Orcid number: 0000-0001-8019-3655 Department of Mathematics Sirjan University of Technology Sirjan, Iran *Email address*: y.sayyari@gmail.com

Somayeh Mirzadeh Orcid number: 0000-0002-2804-7068 Department of Mathematics University of Hormozgan, P.O. Box 3995 Bandar Abbas, Iran *Email address*: mirzadeh@hormozgan.ac.ir