

## REVIEW ARTICLE

# Diagnostic Accuracy of Ultrasonography and Radiography in Detection of Pulmonary Contusion; a Systematic Review and Meta-Analysis

Mostafa Hosseini<sup>1,2</sup>, Parisa Ghelichkhani<sup>3</sup>, Masoud Baikpour<sup>4</sup>, Abbas Tafakhori<sup>5,6</sup>, Hadi Asady<sup>7</sup>,  
 Mohammad Javad Haji Ghanbari<sup>8</sup>, Mahmoud Yousefifard<sup>9\*</sup>, Saeed Safari<sup>10</sup>

1. Sina Trauma and Surgery Research Center, Tehran University Medical Sciences, Tehran, Iran.

2. Department of Epidemiology and Biostatistics, school of Public Health, Tehran University of Medical Sciences, Tehran, Iran.

3. Department of Intensive Care Nursing, School of Nursing and Midwifery, Tehran University of Medical Sciences, Tehran, Iran.

4. Department of Medicine, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran.

5. Department of Neurology, School of Medicine, Imam Khomeini Hospital, Tehran University of Medical Sciences, Tehran, Iran.

6. Iranian Center of Neurological Research, Tehran University of Medical Sciences, Tehran, Iran.

7. Department of Occupational Health Engineering, Faculty of Public Health, Tehran University of Medical Sciences, Tehran, Iran.

8. Emergency Medicine Department, Rasoul Akram Hospital, Iran University of Medical Sciences, Tehran, Iran.

9. Department of Physiology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran.

10. Department of Emergency Medicine, Shohadaye Tajrish Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

\*Corresponding Author: Mahmoud Yousefifard, Department of Physiology, School of Medicine, Tehran University of Medical Sciences, Poursina Ave, Tehran, Iran. Email: usefifard@razi.tums.ac.ir; Phone/Fax: +982122721155.

Received: June 2015; Accepted: August 2015

## Abstract

**Introduction:** Ultrasonography is currently being used as one of the diagnostic modalities in various medical emergencies for screening of trauma patients. The diagnostic value of this modality in detection of traumatic chest injuries has been evaluated by several studies but its diagnostic accuracy in diagnosis of pulmonary contusion is a matter of discussion. Therefore, the present study aimed to determine the diagnostic accuracy of ultrasonography and radiography in detection of pulmonary contusion through a systematic review and meta-analysis. **Methods:** An extended systematic search was performed by two reviewers in databases of Medline, EMBASE, ISI Web of Knowledge, Scopus, Cochrane Library, and ProQuest. They extracted the data and assessed the quality of the studies. After summarization of data into true positive, false positive, true negative, and false negative meta-analysis was carried out via a mixed-effects binary regression model. Further subgroup analysis was performed due to a significant heterogeneity between the studies. **Results:** 12 studies were included in this meta-analysis (1681 chest trauma patients, 76% male). Pooled sensitivity of ultrasonography in detection of pulmonary contusion was 0.92 (95% CI: 0.81-0.96; I<sup>2</sup>= 95.81, p<0.001) and its pooled specificity was calculated to be 0.89 (95% CI: 0.85-0.93; I<sup>2</sup>= 67.29, p<0.001) while these figures for chest radiography were 0.44 (95% CI: 0.32-0.58; I<sup>2</sup>= 87.52, p<0.001) and 0.98 (95% CI: 0.88-1.0; I<sup>2</sup>= 95.22, p<0.001), respectively. Subgroup analysis showed that the sources of heterogeneity between the studies were sampling method, operator, frequency of the transducer, and sample size. **Conclusion:** Ultrasonography was found to be a better screening tool in detection of pulmonary contusion. Moreover, an ultrasonography performed by a radiologist / intensivist with 1-5MHz probe has a higher diagnostic value in identifying pulmonary contusions.

**Key words:** Pulmonary contusion; ultrasonography; radiography; diagnostic tests, routine

**Cite this article as:** Hosseini M, Ghelichkhani P, Baikpour M, et al. Diagnostic accuracy of ultrasonography and radiography in detection of pulmonary contusion; a systematic review and meta-analysis. *Emergency*. 2015;3(4):127-34.

## Introduction:

Pulmonary contusion is a common complication of traumatic thoracic injuries. Reports indicate that 25 to 80% of thoracic traumas are associated with pulmonary contusion (1, 2). Various techniques have been proposed for detection of this lesion including clinical assessment, chest radiography (CXR), arterial blood gas, and computed tomography (CT) scan (3, 4). CXR is

the most common diagnostic tool in detection of pulmonary contusion but presence of hemothorax or pneumothorax might complicate the diagnosis (5-7). Moreover, identification of this lesion in CXR is not possible in the first 6 hours after injury (8, 9). CT scan is the most accurate diagnostic tool for pulmonary contusion and can detect the lesion right after the injury (10, 11).

Ultrasonography reported to have acceptable sensitivity



and specificity in detection of pulmonary contusion (11-13). In the last 10 years many studies have evaluated the diagnostic values of ultrasonography and radiography in detection of traumatic thoracic injuries including pulmonary contusion (14-16), but reaching a consensus has been hindered by the vast disagreements on this subject. One of the ways to overcome this problem is conducting a systematic review and meta-analysis (17, 18). In this regard, we aimed to compare the diagnostic values of these two modalities in detection of pulmonary contusion through a meta-analysis of the available literature.


## Methods:

### Search strategy and selection criteria

Search strategy was based on the keywords related to ultrasonography and chest radiography including "Ultrasonography" OR "Sonography" OR "Ultrasound" OR "Chest Film" OR "Chest Radiograph" combined with pulmonary contusion-related terms including "Contusions" OR "Pulmonary Contusion" OR "Lung Contusion". The

systematic search was carried out in databases of Medline (via PubMed), EMBASE, ISI Web of Knowledge, Scopus, Cochrane Library, and ProQuest directed at finding retrospective and prospective original articles. We run a hand search using Google Scholar for extracting further studies. Bibliographies of the related and review articles were scanned in order to find relevant undiscovered studies in our systematic search. The search keywords were extracted from Medical Subject Heading (MeSH) terms and Emtree.

Review and editorial articles, case reports, letters to editors, poster presentations, and meeting abstracts were excluded from this survey. Application of a reference test other than CT scan and conducting the study on animal samples were also considered as exclusion criteria. Two reviewers (M.Y, P.G) extracted data in true positive (TP), true negative (TN), false positive (FP), and false negative (FN). In cases where these values could not be obtained neither from the article nor by contacting the authors, the survey were excluded from the study.



**Figure 1:** Flowchart of the study.



**Table 1:** Charecteristics of included studies

Study	No. of Patient (+ / -) <sup>1</sup>	Age <sup>2</sup> (years)	Male (%)	Reference / Index	Transducer / Operator	Sampling	Weaknesses
Lichtenstein 2004 (19)	184 / 200	58±15	NR	CT/ US, CXR	5 MHz / Intensivist	Consecutive	----
Soldati 2006 (11)	37 / 51	35 (18-89)	72.4	CT/ US, CXR	3.5- to 5-MHz / EP	Consecutive	The most patients were as- sessed retrospectively
Elmali 2007 (20)	39 / 21	43 (16-85)	80	CT / CXR	NA / Radiologist	Consecutive	Low sample size
Traub 2007 (21)	44 / 97	47 (18-89)	75	CT / CXR	NA / Radiologist	Convenience	Retrospective design
Rocco 2008 (22)	63 / 117	42±14	66.7	CT/ US, CXR	3.5 MHz / Intensivist	Consecutive	Possibility of selection bias Low sample size
Xirouchaki 2011 (13)	54 / 30	57±21.5	81	CT/ US, CXR	5-to 9-MHz / Intensivist	Convenience	Low sample size
Hyacinthe 2012 (23)	147 / 90	39 (22-51)	82	CT / US, CXR	5-to 2-MHz / EP	Consecutive	Possibility of selection bias
Blasińska 2013 (24)	11 / 49	NR	NR	CT / CXR	NA / Radiologist	Consecutive	Low sample size
Chardoli 2013 (25)	11 / 189	38 (16-90)	84	CT / CXR	NA / EP	Convenience	Lack of Blinding
Leblanc 2014 (26)	38 / 7	36 (15-56)	71	CT / US, CXR	5-to 1-MHz / Intensivist	Convenience	Possible selection bias Low sample size
Helmy 2015 (27)	40 / 10	39 (18-67)	70	CT / US, CXR	5 MHz / Radiologist	Convenience	Possibility of selection bias Low sample size
Vafaei 2015 (12)	48 / 104	31 (4-67)	77.6	CT/ US, CXR	3.5-to 7-MHz / EP	Convenience	Possibility of selection bias Possibility of selection bias

1, (+ / -): (number of patient with contusion / number of patient without contusion); 2, Number are presented as mean ± standard deviation or (range). CT: Computed tomography; CXR: Chest radiography; EP: Emergency physician; NA: Not applicable; NR: Not Reported; US: Ultrasonography.



### Data extraction

Two reviewers (M.Y, P.G) independently assessed the titles and abstracts of the articles found in the systematic search. Then the full texts of the potentially relevant articles were evaluated and the data from the studies that met the inclusion criteria were precisely summarized in details. No time or language limitations were established. Quality assessment of the articles was performed according to the guidelines suggested by 14-Item Quality Assessment of Diagnostic Accuracy Studies (QUADAS-2) tool (28). Based on this criteria, all included studies were screened for presence of selection, performance, recording, and reporting biases.

Demographic characteristics of the patients including age, gender, the number of patients with/without pulmonary contusion according to the results of CT scans, characteristics of ultrasound device (transducer, frequency) and its operator, blinding status, and sampling method (consecutive, convenience). Finally, the number of TP, TN, FN, and FP cases were recorded. Disagreements were solved by the third author (M.H). The method proposed by Siström and Mergo (29) was used to extract the data presented as charts. Web-based programs were utilized to calculate the number of TP, TN, FN, and FP cases from the articles in which only the sensitivity and specificity were presented.

### Statistical analysis

Analysis was done by STATA 11.0 statistical software via "MIDAS" module. Summary receiver operative curves (SROC), sensitivity, specificity, positive likelihood ratio and negative likelihood ratio of ultrasonography and radiography in detection of pulmonary contusion with 95% confidence interval (95% CI) were estimated. In cases where data were presented separately for each hemi-thorax the information were included separately as presented in the original article. Due to the significant heterogeneity between the included studies, mixed effects binary regression model was applied. Heterogeneity was evaluated through calculation of  $I^2$  and  $\chi^2$  tests and a p value of less than 0.1 along with an  $I^2$  greater than 50% were considered as presence of considerable heterogeneity (30).

In order to recognize the sources of heterogeneity, subgroup analysis was performed considering the sampling method (consecutive/ convenience), operator (emergency physician/ other specialists) or the interpreting physician, the ultrasound device's frequency of the transducer (1-5 MHz/ 5-10 MHz), and sample size (less than 100 patients/ more than 100 patients). In all the analyses, p value of less than 0.05 was considered as statistically significant.

## Results:

### Study characteristics

Search in the mentioned databases yielded 15 studies

that met the inclusion criteria. Further manual search resulted in finding 3 more related surveys. After summarization and quality assessment, 12 studies were included (11-13, 19-27) (Figure 1). A total of 716 patients with pulmonary contusion and 965 subjects without were evaluated. Their age ranged from 4 to 90 years old and male patients comprised 76% of the study population. The summary of included surveys is presented in Table 1. Diagnostic accuracy of ultrasonography and radiography in detection of pulmonary contusion were assessed simultaneously in eight studies (11-13, 19, 22, 23, 26, 27) and the accuracy of radiography was evaluated individually in four surveys (20, 21, 24, 25). Considerable heterogeneity was observed between the studies ( $P < 0.001$ ). No publication bias was observed in evaluation of the diagnostic accuracy of ultrasonography ( $p = 0.97$ ) and chest radiography ( $p = 0.15$ ) (Figure 2).

### Meta-analysis

#### - Ultrasonography

Area under the curve of SROC for ultrasonography in pulmonary contusion diagnosis was found to be 0.93 (95% CI: 0.91 - 0.95) (Figure 3-A). Pooled sensitivity of ultrasonography in this regard was 0.92 (95% CI: 0.81 - 0.96;  $I^2 = 95.81$ ,  $p < 0.001$ ) and its pooled specificity was estimated to be 0.89 (95% CI: 0.85 - 0.93;  $I^2 = 67.29$ ,  $p < 0.001$ ). Ultrasonography had pooled positive and negative likelihood ratios of 8.94 (95% CI: 5.95 - 93.36;  $I^2 = 67.92$ ,  $p < 0.001$ ) and 0.09 (95% CI: 0.04 - 0.22;  $I^2 = 06.36$ ,  $p < 0.001$ ), respectively (Figure 4).

Table 2 demonstrates the results of subgroup analysis. The sensitivity of this modality was lower when consecutive sampling method was used (0.87 vs. 0.97), procedure was performed via an emergency specialist (0.77 vs. 0.95), sample sizes of higher than 100 patients, the sensitivity (0.86 vs. 0.96), and frequencies of ultrasonography probe was higher than 5 MHz (0.86 vs. 0.93).

#### - Chest Radiography

Data from 12 surveys were included in this part of meta-analysis (11-13, 19-27). Area under the SROC for radiography in detection of pulmonary contusion was 0.72 (95% CI: 0.67 - 0.75) (Figure 3-B). Pooled sensitivity and specificity of this diagnostic tool were 0.44 (95% CI: 0.32 - 0.58;  $I^2 = 87.52$ ,  $p < 0.001$ ) and 0.98 (95% CI: 0.88 - 1.0;  $I^2 = 95.22$ ,  $p < 0.001$ ), respectively. Pooled positive and negative likelihood ratios were also calculated to be 19.69 (95% CI: 3.59 - 108.07;  $I^2 = 88.75$ ,  $p < 0.001$ ) and 0.57 (95% CI: 0.45 - 0.72;  $I^2 = 93.13$ ,  $p < 0.001$ ), respectively (Figure 5).

Subgroup analysis showed that the sensitivity of radiography is affected by the interpreting physician of the plain film (emergency physician/ other specialists) and sample size (Table 2). According to the results of this analysis, the sensitivity of this imaging modality is higher when the radiographs were interpreted by a radiologist or intensivist (0.49; 95% CI: 0.30-0.68) compared to an



**Table 2:** Subgroup analysis of diagnostic accuracy for chest radiography and ultrasonography in detection of pulmonary contusion

Covariate	No. of studies	Bivariate random-effect model					
		Sensitivity (95% CI)	P	Specificity (95% CI)	p	heterogeneity, I <sup>2</sup>	P*
Ultrasonography							
Patient enrollment							
Consecutive	5	0.87 (0.78-0.96)	0.16	0.90 (0.86-0.95)	<0.001	49.0 %	0.14
Convenience	3	0.97 (0.94-1.00)		0.88 (0.82-0.95)			
Operator							
Emergency physician	3	0.77 (0.62-0.93)	<0.001	0.90 (0.84-0.95)	<0.001	68.0 %	0.04
Other physician	5	0.95 (0.92-0.99)		0.89 (0.84-0.94)			
Sample size							
< 100	4	0.96 (0.90-1.00)	0.38	0.94 (0.89-0.99)	<0.001	55.0 %	0.11
≥ 100	4	0.86 (0.73-0.99)		0.88 (0.84-0.95)			
Frequency							
1-5 MHz	5	0.93 (0.87-1.0)	0.41	0.88 (0.84-0.91)	<0.001	58.0 %	0.09
5-10 MHz	3	0.86 (0.65-1.0)		0.93 (0.89-0.98)			
Radiography							
Patient enrollment							
Consecutive	6	0.45 (0.26-0.63)	0.90	0.99 (0.96-1.00)	0.72	0.0 %	0.61
Convenience	6	0.44 (0.24-0.63)		0.95 (0.85-1.00)			
Operator							
Emergency physician	6	0.40 (0.21-0.58)	0.74	0.98 (0.94-1.00)	0.03	0.0 %	0.79
Other physician	6	0.49 (0.30-0.68)		0.97 (0.90-1.00)			
Sample size							
< 100	6	0.55 (0.38-0.72)	0.15	0.94 (0.82-1.00)	0.99	36.0 %	0.21
≥ 100	6	0.35 (0.19-0.51)		0.99 (0.97-1.00)			

\*, P value &lt; 0.1 was considered as significant for heterogeneity; CI: Confidence interval.





emergency specialist (0.40; 95% CI: 0.21 - 0.58).

### Discussion:

The present meta-analysis is the first to assess the diagnostic accuracy of ultrasonography and radiography in detection of pulmonary contusion. The results illustrate a higher sensitivity of ultrasonography compared to radiography (0.92 vs. 0.44) in this regard, whereas the specificity of radiography was slightly higher (0.97 vs. 0.89). Since these two imaging modalities are the first diagnostic tools for assessment of traumatic thoracic injuries, their screening accuracy is of utmost importance. Accordingly, ultrasonography has better screening performance characteristics in detection of pulmonary contusion compared to radiography.

Various studies have pointed out the fact that diagnostic accuracy of ultrasonography is directly dependent on the skills of the operator (14, 15, 31, 32). Findings of the present survey were also congruent with this statement to some extent. The results demonstrated a higher sensitivity of ultrasonography in detection of pulmonary contusion when performed by a radiologist or an intensivist compared to emergency specialists. This might be due to the nature of pulmonary contusion whose diagnostic signs are very challenging to detect. The most important signs of pulmonary contusion identified by ultrasonography include multiple B-lines and an irregularly delineated tissue image which might be a moderately hypo-echoic blurred lesion (16). Furthermore, after observation of these signs, the operator should rule out pneumothorax as well. Therefore, experience plays an important role in pulmonary contusion diagnosis.

Frequency of transducer was another factor affecting the diagnostic accuracy of ultrasonography. Application of transducers with frequencies lower than 5MHz yield greater diagnostic values compared to higher. This finding is also related to the nature of the lesion. Contusion is characterized by parenchymal injuries and accumulation of fluid and blood in the lung tissues (16). These tissues lie in the deepest layers of chest cavity and so the penetrating power of ultrasound wave is more important than the image resolution (which is directly related to the wave's frequency). Since ultrasound waves with lower frequencies have greater penetrating powers, application of these probes increases the chances of pulmonary contusion diagnosis. Sample size was also found to have an effect on diagnostic values of ultrasonography and radiography in detection of pulmonary contusion. The sensitivity of both these modalities was found to be higher in the studies with sample sizes of less than 100 patients. This might be due to possible selection bias in these studies (33). Selection of patients with severe traumas and so the higher chances of injury identification via imaging would be prominent in these studies. Moreover in some of these surveys, pneumothorax

patients had been excluded which might have made the diagnosis easier (27).

Utilization of three strategies has improved the quality of the present meta-analysis. Firstly, comprehensive search in databases to include the maximum number of related surveys and secondly, elimination of publication bias. Thirdly, the effects of heterogeneity between the studies were controlled by subgroup analysis.

On the other hand, simultaneous inclusion of retrospective and prospective studies might be considered as a limitation of this study. However, evaluation of outliers on the scatterplot based on standardized predictive random effects revealed that retrospective surveys are not the cause of diversity between the studies. Moreover, due to the observational nature of included studies, precise assessment of causal relationships was impossible.

### Conclusion:

The results of present meta-analysis revealed the better screening performance characteristics of chest ultrasonography compared to radiography in detection of pulmonary contusion. It should be mentioned that these characteristics were dependent on operator and characteristics of device.

### Acknowledgments:

None

### Conflict of interest:

None

### Funding support:

This research has been supported by Tehran University of Medical Sciences & health Services grant number: 93-02-38-25618.

### Authors' contributions:

All authors passed four criteria for authorship contribution based on recommendations of the International Committee of Medical Journal Editors.

### References:

1. Balci AE, Kazez A, Eren S, Ayan E, Ozalp K, Eren MN. Blunt thoracic trauma in children: review of 137 cases. *Eur J Cardiothorac Surg.* 2004;26(2):387-92.
2. Skinner DL, den Hollander D, Laing GL, Rodseth RN, Muckart DJ. Severe blunt thoracic trauma: differences between adults and children in a level I trauma centre. *S Afr Med J.* 2015;105(1):47-51.
3. Keough V, Pudelek B. Blunt chest trauma: review of selected pulmonary injuries focusing on pulmonary contusion. *AACN Clin Issues.* 2001;12(2):270-81.
4. Simon B, Ebert J, Bokhari F, et al. Management of pulmonary contusion and flail chest: an Eastern Association for the Surgery of Trauma practice management guideline. *J Trauma Acute Care Surg.* 2012;73(5):S351-S61.
5. Chapagain D, Reddy D, Shah S, Shrestha K. Diagnostic modalities x-ray and CT chest differ in the management of thoracic injury. *Journal of College of Medical Sciences-Nepal.* 2015;10(1):22-31.
6. Kea B, Gamarallage R, Vairamuthu H, et al. What is the clinical significance of chest CT when the chest x-ray result is normal



in patients with blunt trauma? *Am J Emerg Med.* 2013;31(8):1268-73.

7. Forouzanfar MM, Safari S, Niazazari M, et al. Clinical decision rule to prevent unnecessary chest X-ray in patients with blunt multiple traumas. *Emerg Med Australas.* 2014;26(6):561-6.

8. Allen GS, Coates NE. Pulmonary contusion: a collective review. *Am Surg.* 1996;62(11):895-900.

9. Wanek S, Mayberry JC. Blunt thoracic trauma: flail chest, pulmonary contusion, and blast injury. *Crit Care Clin.* 2004;20(1):71-81.

10. Cohn SM, DuBose JJ. Pulmonary contusion: an update on recent advances in clinical management. *World J Surg.* 2010;34(8):1959-70.

11. Soldati G, Testa A, Silva FR, Carbone L, Portale G, Silveri NG. Chest ultrasonography in lung contusion. *CHEST Journal.* 2006;130(2):533-8.

12. Vafaei A, Hatamabadi HR, Heidary K, Alimohammadi H, Tarbiat M. Diagnostic Accuracy of Ultrasonography and Radiography in Initial Evaluation of Chest Trauma Patients. *Emergency.* 2015;3:[In press].

13. Xirouchaki N, Magkanas E, Vaporidi K, et al. Lung ultrasound in critically ill patients: comparison with bedside chest radiography. *Intensive Care Med.* 2011;37(9):1488-93.

14. Bui-Mansfield LT, Chen DC, O'Brien SD. Accuracy of ultrasound of musculoskeletal soft-tissue tumors. *Am J Roentgenol.* 2015;204(2):W532-40.

15. Fine D, Perring S, Herbetko J, Hacking C, Fleming J, Dewbury K. Three-dimensional (3D) ultrasound imaging of the gallbladder and dilated biliary tree: reconstruction from real-time B-scans. *Br J Radiol.* 1991;64(767):1056-7.

16. Ganie FA, Lone H, Lone GN, et al. Lung Contusion: A Clinico-Pathological Entity with Unpredictable Clinical Course. *Bull Emerg Trauma.* 2013;1(1):7-16.

17. Hosseini M, Yousefifard M, Aziznejad H, Nasirinezhad F. The Effect of bone marrow derived mesenchymal stem cell transplantation on allodynia and hyperalgesia in neuropathic animals: A systematic review with meta-analysis. *Biol Blood Marrow Transplant.* 2015; 21(9): 1537-44.

18. Ebrahimi A, Yousefifard M, Kazemi HM, et al. Diagnostic Accuracy of Chest Ultrasonography versus Chest Radiography for Identification of Pneumothorax: A Systematic Review and Meta-Analysis. *Tanaffos.* 2014;13(4):29-40.

19. Lichtenstein D, Goldstein I, Mourgeon E, Cluzel P, Grenier P, Rouby JJ. Comparative diagnostic performances of auscultation, chest radiography, and lung ultrasonography in acute respiratory distress syndrome. *Anesthesiology.* 2004;100(1):9-15.

20. Elmali M, Baydin A, Nural MS, Arslan B, Ceyhan M, Gurmen N. Lung parenchymal injury and its frequency in blunt thoracic

trauma: the diagnostic value of chest radiography and thoracic CT. *Diagn Interv Radiol.* 2007;13(4):179-82.

21. Traub M, Stevenson M, McEvoy S, et al. The use of chest computed tomography versus chest X-ray in patients with major blunt trauma. *Injury.* 2007;38(1):43-7.

22. Rocco M, Carbone I, Morelli A, et al. Diagnostic accuracy of bedside ultrasonography in the ICU: feasibility of detecting pulmonary effusion and lung contusion in patients on respiratory support after severe blunt thoracic trauma. *Acta Anaesthesiol Scand.* 2008;52(6):776-84.

23. Hyacinthe AC, Broux C, Francony G, et al. Diagnostic accuracy of ultrasonography in the acute assessment of common thoracic lesions after trauma. *Chest.* 2012;141(5):1177-83.

24. Błasińska-Przerwa K, Pachó R, Bestry I. The application of MDCT in the diagnosis of chest trauma. *Pneumonol Alergol Pol.* 2013;81(6):518-26.

25. Chardoli M, Hasan-Ghaliaye T, Akbari H, Rahimi-Movaghar V. Accuracy of chest radiography versus chest computed tomography in hemodynamically stable patients with blunt chest trauma. *Chin J Traumatol.* 2013;16(6):351-4.

26. Leblanc D, Bouvet C, Degiovanni F, et al. Early lung ultrasonography predicts the occurrence of acute respiratory distress syndrome in blunt trauma patients. *Intensive Care Med.* 2014;40(10):1468-74.

27. Helmy S, Beshay B, Hady MA, Mansour A. Role of chest ultrasonography in the diagnosis of lung contusion. *Egypt J Chest Dis Tuberc.* 2015;64:469-75.

28. Whiting PF, Rutjes AW, Westwood ME, et al. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Ann Intern Med.* 2011;155(8):529-36.

29. Siström CL, Mergo PJ. A simple method for obtaining original data from published graphs and plots. *Am J Roentgenol.* 2000;174(5):1241-4.


30. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ.* 2003;327(7414):557.

31. Abbasi S, Farsi D, Hafezimoghadam P, Fathi M, Zare MA. Accuracy of emergency physician-performed ultrasound in detecting traumatic pneumothorax after a 2-h training course. *Eur J Emerg Med.* 2013;20(3):173-7.


32. Lee JH, Jeong YK, Park KB, Park JK, Jeong AK, Hwang JC. Operator-dependent techniques for graded compression sonography to detect the appendix and diagnose acute appendicitis. *Am J Roentgenol.* 2005;184(1):91-7.

33. Deeks JJ, Macaskill P, Irwig L. The performance of tests of publication bias and other sample size effects in systematic reviews of diagnostic test accuracy was assessed. *J Clin Epidemiol.* 2005;58(9):882-93.






**Figure 2:** Deeks' funnel plot asymmetry test for assessment of publication bias. P values < 0.05 were considered as significant. Ultrasonography (A); Radiography (B). ESS: Effective sample sizes.




**Figure 3:** Summary receiver operative curves (SROC) with prediction and confidence contours of ultrasonography (A) and chest radiography (B) in detection of pulmonary contusion. AUC: Area under the curve; SENS: Sensitivity; SPEC: Specificity.



A




B




**Figure 4:** Forest plot of screening performance characteristics of chest ultrasonography in detection of pulmonary contusion. Sensitivity and specificity (A); Diagnostic likelihood ratio (DLR) (B). CI: Confidence interval.

A



B



**Figure 5:** Forest plot of screening performance characteristics of chest radiography in detection of pulmonary contusion. Sensitivity and specificity (A); Diagnostic likelihood ratio (DLR) (B). CI: Confidence interval.

