

# **REVIEW ARTICLE**

# Application of Ultrasonography and Radiography in Detection of Hemothorax; a Systematic Review and Meta-Analysis

Vafa Rahimi-Movaghar<sup>1</sup>, Mahmoud Yousefifard<sup>2</sup>, Parisa Ghelichkhani<sup>3</sup>, Masoud Baikpour<sup>4</sup>, Abbas Tafakhori<sup>5,6</sup>, Hadi Asady<sup>7</sup>, Gholamreza Faridaalaee<sup>8</sup>, Mostafa Hosseini<sup>9\*</sup>, Saeed Safari<sup>10</sup>

Sina Trauma and Surgery Research Center, Tehran University Medical Sciences, Tehran, Iran.
Department of Physiology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran.
Department of Intensive Care Nursing, School of Nursing and Midwifery, Tehran University of Medical Sciences, Tehran, Iran.
Department of Medicine, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran.
Department of Neurology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran.
Department of Neurology, School of Medicine, Imam Khomeini Hospital, Tehran University of Medical Sciences, Tehran, Iran.
Iranian Center of Neurological Research, Tehran University of Medical Sciences, Tehran, Iran.
Department of Occupational Health Engineering, Faculty of Public Health, Tehran University of Medical Sciences, Tehran, Iran.
Department of Emergency Medicine, Maragheh University of Medical Sciences, Maragheh, Iran.
Department of Epidemiology and Biostatistics, school of Public Health, Tehran University of Medical Sciences, Tehran, Iran.
Department of Emergency Medicine, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran.

\*Corresponding Author: Mostafa Hosseini, Department of Epidemiology and Biostatistics School of Public Health, Tehran University of Medical Sciences, Poursina Ave, Tehran, Iran; Email: mhossein110@yahoo.com; Tel: +982188989125; Fax: +982188989127. Received: June 2015; Accepted: August 2015

#### Abstract

Introduction: Hemothorax is one of the most prevalent injuries caused by thoracic traumas. Early detection and treatment of this injury is of utmost importance in prognosis of the patient, but there are still controversial debates on the diagnostic value of imaging techniques in detection of hemothorax. Therefore, the present study aimed to evaluate the diagnostic value of chest ultrasonography and radiography in detection of hemothorax through a systematic review and meta-analysis. Methods: Two independent reviewers performed an extended systematic search in databases of Medline, EMBASE, ISI Web of Knowledge, Scopus, Cochrane Library, and ProQuest. Data were extract and quality of the relevant studies were assessed. The number of true positive, false positive, true negative and false negative cases were extracted and screening performance characteristics of two imaging techniques were calculated using a mixed-effects binary regression model. Results: Data from 12 studies were extracted and included in the meta-analysis (7361 patients, 77.1% male). Pooled sensitivity and specificity of ultrasonography in detection of hemothorax were 0.67 (95% CI: 0.41-0.86; I2= 68.38, p<0.001) and 0.99 (95% CI: 0.95-1.0; I2= 88.16, p<0.001), respectively. These measures for radiography were 0.54 (95% CI: 0.33-0.75; I2= 92.85, p<0.001) and 0.99 (95% CI: 0.94-1.0; I2= 99.22, p<0.001), respectively. Subgroup analysis found operator of the ultrasonography device, frequency of the transducer and sample size to be important sources of heterogeneity of included studies. **Conclusion:** The results of this study showed that although the sensitivity of ultrasonography in detection of hemothorax is relatively higher than radiography, but it is still at a moderate level (0.67%). The specificity of both imaging modalities were found to be at an excellent level in this regard. The screening characteristics of ultrasonography was found to be influenced of the operator and frequency of transducer.

Keywords: Hemothorax; ultrasonography; radiography; diagnostic tests, routine

**Cite this article as:** Rahimi-Movaghar V, Yousefifard M, Ghelichkhani P, et al. Application of ultrasonography and radiography in detection of hemothorax: a systematic review and meta-analysis. Emergency. 2016; 4(3):116-126.

#### Introduction:

hest traumas are one of the most important causes of mortality in the fourth decade of life (1, 2). 25% of trauma mortalities are due to these injuries (3). In this regard, imaging techniques play a vital role in management of these patients. Although some thoracic traumas are treated according to clinical findings of the patient before performing any imaging studies, but in many cases application of various imaging modalities such as computed tomography (CT) scan, plain chest Xray (CXR) and ultrasonography are necessary. Among these modalities, CT scan is the gold standard for identification of intra thoracic injuries following trauma with a significantly high diagnostic value for occult and soft tissue injuries (4-9). However, limited availability of CT scan in all medical centers, limitations in patient transfer



This open-access article distributed under the terms of the Creative Commons Attribution NonCommercial 3.0 License (CC BY-NC 3.0). Copyright © 2016 Shahid Beheshti University of Medical Sciences. All rights reserved. Downloaded from: www.jemerg.com///.SID.u/ to radiology department and radiation exposure led the researchers to look for other diagnostic tools (10).

CXR is the first diagnostic test for screening of thoracic traumas but the limitations of supine radiography in some traumatic injuries such as pneumothorax is confirmed in various studies (11, 12). Moreover, Low diagnostic yield of routine chest radiography in patients with thoracic injuries encouraged the researchers to search for alternative imaging techniques (11-14). Accordingly, in recent years scoring systems such as thoracic injury rule out criteria (TIRC) and national emergency X-Radiography utilization study (NEXUS) have been developed to lower the burden of unnecessary imaging studies (15, 16).

Major attention has recently been drawn to ultrasonography as a quick screening tool with minimum complications (17). It has shown to have superior diagnostic value in detection of thoracic traumatic injuries compared to chest radiography (18-22). However, diagnostic accuracy of ultrasonography is highly dependent on the skills of the operator and is usually not reliable in detection of injuries without bleeding or free fluid (23-25).

Hemothorax is one the traumatic thoracic injuries caused by accumulation of blood in pleural cavity. This lesion along with pneumothorax is present in 83% of thoracic traumas (26). However, detection of this complication via chest radiography is not possible unless the volume of hemothorax exceeds 175 milliliters (27). Moreover, the diagnostic value of ultrasonography for hemothorax is still a matter of debate as well (18, 19, 28, 29). Recently multiple systematic reviews have been published to evaluate the diagnostic value of ultrasonography and chest radiography in detection of thoracic traumas, but almost all of them have assessed pneumothorax. These reviews showed a higher sensitivity of ultrasonography in identification of pneumothorax compared to chest radiography (29, 30). None of these surveys has taken a meta-analytic approach towards assessing diagnostic value of these imaging modalities in detection of hemothorax. Therefore, the present study



Figure 1: Flowchart of the study.



This open-access article distributed under the terms of the Creative Commons Attribution NonCommercial 3.0 License (CC BY-NC 3.0). Copyright © 2016 Shahid Beheshti University of Medical Sciences. All rights reserved. Downloaded from: www.jemerg.com.w.j

Table 1: Charect	eristics of include	ed studies					
Study	No. of patient (+ / -) <sup>1</sup>	Age <sup>2</sup> (years)	Male (%)	Reference / Index	Transducer / Operator	Sampling	Weaknesses
Ma 1997 (27)	26/214	NR	NR	CT / US, CXR	3.5-to 2.5-MHz / EP	Consecutive	Retrospective design
Abboud 2003 (28)	14 / 126	38 (5–89)	NR	CT / US	3.75 MHz / EP	Convenience	Time interval between US and CT scan was varied; Blinding was not performed
Brooks 2004 (18)	12 / 49	NR	NR	CT / US	4- to 2-MHz / EP	Convenience	Small sample size; Possibility of selection bias
Traub 2007 (31)	16 / 125	47 (18-89)	75	CT / CXR	NA / Radiologist	Convenience	Retrospective design Possibility of selection bias
Hyacinthe 2012 (19)	35 / 202	39 (22-51)	82	CT / US	5- to 2-MHz / EP	Consecutive	Possibility of selection bias
Poveda 2012 (32)	47 / 21	39 (16-70)	89.7	Surgery / US	3.75 MHZ / Radiologist	Convenience	Possibility of selection bias
Błasińska 2013 (33)	24 / 36	NR	NR	CT / CXR	NA / Radiologist	Consecutive	Low sample size
Uz 2013 (34)	21 / 86	37 (19.8)	80.4	CT/ US	10-to 5-MHz / Radiologist	Consecutive	Low sample size
Chardoli 2013 (35)	14 / 186	38 (16-90)	84	CT / CXR	NA/ EP	Convenience	The interpretation of the CXR and CT were not in blind fashion Possible selection bias
Leblanc 2014 (36)	19 / 26	36 (15-56)	71	CT/ US, CXR	5-to 1-MHz / Intensivist	Convenience	Low sample size Possibility of selection bias
Langdorf 2015 (37)	230 / 5682	≥ 15	57.4	CT/ CXR	NA / Radiologist	Convenience	Possibility of selection bias Variation in timing of CT and CXR
Vafaei 2015 (38)	29 / 123	31 (4-67)	77.6	CT / US, CXR	3.5-to 7-MHz / EP	Convenience	Possibility of selection bias
1, (+ / -): Numbe (range). CT: Com phy.	r of patient with l puted tomograph	hemothorax / m ıy; CXR: Chest ra	umber of Idiograpł	patient without h 1y; EP: Emergency	lemothorax; 2, Nı y physician; NA: N	umber are pres Vot applicable;	ented as mean ± standard deviation or NR: Not Reported; US: Ultrasonogra-



This open-access article distributed under the terms of the Creative Commons Attribution NonCommercial 3.0 License (CC BY-NC 3.0). Copyright © 2016 Shahid Beheshti University of Medical Sciences. All rights reserved. Downloaded from: www.jemerg.com W. SID. U aimed to evaluate the diagnostic accuracy of chest ultrasonography and radiography in detection of hemothorax through a systematic review of available literature and meta-analysis.

## Methods:

## Search strategy and selection criteria

Two independent reviewer (M.Y, P.G) performed an extended systematic search in databases of Medline (via PubMed), EMBASE, ISI Web of Knowledge, Scopus, Cochrane Library, and ProQuest. We screened google scholar for further studies. The objective was to find surveys evaluating the diagnostic accuracy of ultrasonography or chest radiography in detection of hemothorax.

Keywords were chosen according to Medical Subject Heading (MeSH) terms and EMTREE. Similar keywords were used for search in other databases. These keywords were terms related to ultrasonography and radiography including "Ultrasonography" OR "Sonography" OR "Ultrasound" OR "Chest Film" OR "Chest Radiograph" combined with hemothorax related terms including "Hemothorax" OR "Haemothorax" OR "Haemorrhagic Pleural Effusion". In order to find further studies or unpublished surveys, hand-search was performed on the list of bibliography of relevant studies and the authors were contacted in cases where the data could not be extracted from the survey.

Only the original articles and the surveys conducted on human subjects were included. Review and editorial articles, case reports, letters to editors, poster presentations, and meeting abstracts were excluded. Studies were included only if they presented an evaluation of the diagnostic value of ultrasonography or chest radiography in hemothorax detection. Other inclusion criteria were as follows: confirmation of injury via CT scan or surgery, performance of radiography or ultrasonography for all patients, presentation of true positive, true negative, false positive, and false negative cases (in the article, through contacting the authors, or using webbased calculators). No time or language limitations were applied. Both retrospective and prospective studies were included.

### Data extraction

Two of the authors (M.Y, P.G) independently worked on summarizing the data including assessment of quality of studies, information related to the subjects (age, gender, the number of patients with/without hemothorax, the etiology of hemothorax), the characteristics of ultrasonography device (transducer, frequency), operators and the physicians in charge of interpreting the imaging, blinding status, sampling method (consecutive, convenience), study design (retrospective, prospective), reference test, and the number of true positive, false positive, true negative, and false negative cases. Disagreements were discussed with the third author (M.H) and a solution was proposed. In cases of data inaccessibility, the corresponding authors of the articles were contacted. Data presented as charts were extracted via the method proposed by Sistrom and Mergo (39). In cases where only the sensitivity and specificity were presented in the article, reliable web-based programs were used to calculate the number of true positive, false positive, true negative and false negative cases.

Quality of the studies were evaluated according to the Quality Assessment of Diagnostic Accuracy Studies (QUADAS2) guideline (40). Assessment was performed based on the criteria established for designing a diagnostic survey considering various biases including selection, performance, recording and reporting bias.

### Statistical analysis

Analysis was performed via STATA 11.0 statistical software through MIDAS module. The number of false positive, false negative, true positive and true negative cases were recorded. Then, pooled sensitivity, specificity, positive likelihood ratio, and negative likelihood ratio of chest ultrasonography and radiography in detection of hemothorax were calculated with 95% confidence interval (95% CI). In cases of data presented for each hemithorax separately, we also included the information, separately. In the present study, mixed-effects binary regression model, a type of random effect model, was used because of the presence of significant heterogeneity between the studies. Heterogeneity was assessed through application of I2 and  $\chi$ 2 tests. A p value of less than 0.1 along with an I2 greater than 50% were considered as positive heterogeneity (41).

To identify the source of heterogeneity, subgroup analysis was carried out using a bivariate mixed-effects binary regression model. Subgroup analyses were performed according to sampling method (consecutive / convenience), operator of the ultrasonography device (emergency physician/ other specialists) or the interpreting physician of CXR, the frequency of ultrasonography transducer (1-5 MHz/ 5-10 MHz) and sample size (less than 100 patients/ more than 100 patients).

# **Results:**

### Study characteristics

In literature review, 178 potentially relevant studies were identified, of which 37 met the inclusion criteria. Eventually 12 surveys were included in final meta-analysis (18, 19, 27, 28, 31-38) (Figure 1). Data on 7361 trauma patients including 487 with hemothorax and 6874 without were extracted (77.1% male). Table 1 summarizes the baseline characteristics of included studies. Diagnostic accuracy of ultrasonography and radiography were evaluated simultaneously in three studies (27, 36, 38), whereas the accuracy of ultrasonography and radiography were assessed individually in five



This open-access article distributed under the terms of the Creative Commons Attribution NonCommercial 3.0 License (CC BY-NC 3.0). Copyright © 2016 Shahid Beheshti University of Medical Sciences. All rights reserved. Downloaded from: www.jemerg.com *W*. SID. *i*  (18, 19, 28, 32, 34) and four (31, 33, 35, 37) surveys, respectively. Significant heterogeneity was observed between the studies (P<0.001). No publication bias was found (Figure 2).

## Meta-analysis

## - Ultrasonography

Area under the curve of summary Receiver Operative Curves (SROC) for ultrasonography in detection of hemothorax was 0.97 (95% CI, 0.95-0.98) (Figure 3-A). Its pooled sensitivity and specificity in detection of hemothorax were 0.67 (95% CI: 0.41-0.86; I2= 68.38, p<0.001) and 0.99 (95% CI: 0.95-1.0; I2= 88.16, p<0.001), respectively. In addition, positive and negative likelihood ratios were computed to be 52.88 (95% CI: 9.87-283.23; I2= 80.61, p<0.001) and 0.33 (95% CI: 0.16-0.68; I2= 95.66, p<0.001), respectively (Figure 4). Subgroup analysis found sampling method (consecutive/ convenience), operator (emergency physician/ other specialists), frequency of the transducer (1-5 MHz/ 5-10 MHz), and sample size (less than 100 patients/ more than 100 patients) to be important sources of heterogeneity among studies. Sensitivity of the surveys with consecutive sampling methods were significantly higher than the other studies (0.76 vs. 0.61), but their specificity did not differ considerably (1.0 vs. 0.97). Moreover the sensitivity of ultrasonography in detection of hemothorax was found to be significantly higher when the procedure was performed via an emergency physician (0.70 vs. 0.62) or using a 5-10 MHz transducer (0.75 vs. 0.64) (Table 2).

### - Radiography

As presented in Figure 3-B, area under the SROC curve for radiography was 0.92 (95% CI: 0.89-0.94). Pooled sensitivity and specificity of this modality in detection of hemothorax were 0.54 (95% CI: 0.33-0.75; I2= 92.85, p<0.001) and 0.99 (95% CI: 0.94-1.0; I2= 99.22, p<0.001), respectively. Its positive and negative likelihood ratios were also 46.01 (95% CI: 10.17-208.14; I2= 96.10, p<0.001) and 0.46 (95% CI: 0.29-0.75; I2= 95.66, p<0.001), respectively (Figure 5).

Subgroup analysis showed that sampling method (consecutive / convenience), the interpreting physician (emergency physician/ other specialists), and sample size (less than 100 patients/ more than 100 patients) were important sources of heterogeneity between the studies. Sensitivity of radiography was significantly higher in surveys with consecutive sampling methods (0.61 vs. 0.51) and with sample sizes of less than 100 patients (0.69 vs.0.46).

### **Discussion**:

Sonography, as one of the most available screening tools in emergency settings, is useful for various clinical applications but its diagnostic value in traumatic thoracic injuries is still a controversial subject. The present study is the first to conduct a systematic review with meta-analytic approach on one of the most important thoracic traumas. The results of this study illustrated the relatively higher sensitivity of ultrasonography in this regard, but it is still at a moderate level. The specificity and positive likelihood ratios calculated for both of these modalities were same and excellent.

The results of subgroup analysis showed that the sensitivity of ultrasonography was influenced by the operator of the ultrasound device and frequency of transducer but the specificity of this modality is not affected by them. Accordingly, as ultrasonography performed by an emergency physician has a higher diagnostic value compared to other physicians. This finding might be due to awareness of the emergency physician about the clinical condition of the patient. Although in 8 studies the operators were blinded and in the other 4 the setting was not mentioned, a complete unawareness of the emergency physician about the patients' clinical condition seems unlikely. Since these physicians are the first line of the medical team responsible for treatment of trauma patients, based on their experience they might suspect the presence of hemothorax according to the history and clinical findings of the subjects and consequently pay much more attention to find sonographic evidence of this complication. The diagnostic value of chest radiography in detection of hemothorax is neither affected by the operator nor by the interpreting physician, since the physician or the radiologist is not in direct contact with the patients when interpreting their radiographs.

Ebrahimi et al. (29) found no significant relation between frequency of transducer and detection of pneumothorax but the present survey yielded opposite results regarding detection of hemothorax. This might be due to the fact that the sound wave emitted from the transducer easily moves through fluids (high penetrating power in fluids), since the amount of energy absorbed by the fluids is very low. Therefore, ultrasonography with higher frequencies is able to produce clearer images with higher resolutions (42), an event that does not occur in pneumothorax because propagation of the sound wave through the air is associated with loss of energy and so pictures with higher resolutions are not necessarily yielded with higher frequencies.

The minimum amount of fluid that can be detected by each of these modalities is different, 175 milliliters for radiography and only 20 milliliters for ultrasonography (27).

According to subgroup analysis, diagnostic accuracy of radiography in detection of hemothorax is influenced by the sample size of the survey. The results showed that in the studies with sample sizes of less than 100 patients, the sensitivity of radiography was reported to be higher. This finding could be due to probable selection bias in studies with smaller sample sizes, which might have led



This open-access article distributed under the terms of the Creative Commons Attribution NonCommercial 3.0 License (CC BY-NC 3.0). Copyright © 2016 Shahid Beheshti University of Medical Sciences. All rights reserved. Downloaded from: www.jemerg.com.sciences.com.sci

Lable Z: Subgroup analysis o	of diagnostic accuracy	<i>i</i> for chest radiography ar	nd ultraso Biv	nography in detection of variate random-effect n	nemoth nodel	orax	
covariate	NO. OI SLUUIES	Sensitivity (95% CI)	ď	Specificity (95% CI)	d	Heterogeneity, I <sup>2</sup>	Å
Ultrasonography							
Patient enrollment							
Consecutive	m	0.76(0.45-1.00)	0.56	1.00(0.99-1.00)	0.86	10.00 %	0.33
Convenience	5	0.61 (0.31-0.92)		0.97~(0.93-1.00)			
Operator							
Emergency physician	ы	0.70 (0.42-0.99)	0.68	0.99~(0.98-1.00)	0.02	0.00 %	0.59
Other physician	3	0.62 (0.23-1.00)		$0.97\ (0.90-1.00)$			
Sample size							
< 100	IJ	0.70(0.41-0.98)	0.72	0.99~(0.97-1.00)	0.08	0.00 %	0.74
≥ 100	ŝ	0.63 (0.24-1.00)		$0.98\ (0.95-1.00)$			
Frequency of transducer							
1-5 MHz	Ŋ	0.64 (0.37-0.92)	0.55	0.99~(0.97-1.00)	0.12	0.00 %	0.94
5-10 MHz	33	0.75 (0.37-1.00)		0.99~(0.95-1.00)			
Radiography				C			
Patient enrollment							
Consecutive	ŝ	0.61 (0.24-0.98)	0.65	0.98 (0.94-1.00)	0.03	0.00 %	0.88
Convenience	9	0.51 (0.23-0.78)		0.99 (0.97-1.00)			
Operator							
Emergency physician	Ŋ	0.54(0.24-0.84)	0.96	0.99 (0.97-1.00)	0.07	0.00 %	0.99
Other physician	4	0.55 (0.22-0.87)		0.99 (0.96-1.00)			
Sample size							
<100	33	$0.69\ (0.38-1.00)$	0.32	0.94(0.81-1.00)	0.38	0.00 %	0.99
≥ 100	9	0.46 (0.21-0.72)		0.99(0.99-1.00)			
*, P value < 0.1 was considere	ed as significant for h	leterogeneity; CI: Confide	nce interv	/al.			



This open-access article distributed under the terms of the Creative Commons Attribution NonCommercial 3.0 License (CC BY-NC 3.0). Copyright © 2016 Shahid Beheshti University of Medical Sciences. All rights reserved. Downloaded from: www.jemerg.com W.S.D. in to evaluation of patients with greater volumes of free fluids that increases their chance of detection via radiography (43).

#### - Limitations

Since all included studies were observational precise evaluation of causal relationships was not possible. The skills of the operator in performing ultrasonography were not considered in any of these surveys and the effect of this bias is not clear in the present study. Finally a significant heterogeneity was found between the surveys whose effects were tried to be minimized by application of mixed random effect model and subgroup analysis.

#### **Conclusion:**

The results of this study showed that although the sensitivity of ultrasonography in detection of hemothorax is relatively higher than radiography, but it is still at a moderate level (0.67%). The specificity of both imaging modalities were found to be at an excellent level in this regard. The screening characteristics of ultrasonography was found to be influenced of the operator and frequency of transducer.

Acknowledgments: None Conflict of interest: None Funding support:

This research has been supported by Tehran University of Medical Sciences & health Services grant number: 93-02-38-25618.

#### Authors' contributions:

All authors passed four criteria for authorship contribution based on recommendations of the International

Committee of Medical Journal Editors.

## **References:**

1. Hill A, Fowler R, Pinto R, Nathens A. Epidemiology of major trauma: a Canadian perspective. Can J Surg. 2011;54(3):S45.

2. Søreide K. Epidemiology of major trauma. Br J Surg. 2009;96(7):697-8.

3. Heron M. Deaths: leading causes for 2008. Natl Vital Stat Rep. 2012;60(6):1-94.

4. Omert L, Yeaney WW, Protetch J. Efficacy of thoracic computerized tomography in blunt chest trauma. Am Surg. 2001;67(7):660.

5. Tocino I, Miller MH. Computed tomography in blunt chest trauma. J Thorac Imaging. 1987;2(3):45-59.

6. Trupka A, Waydhas C, Hallfeldt KK, Nast-Kolb D, Pfeifer KJ, Schweiberer L. Value of thoracic computed tomography in the first assessment of severely injured patients with blunt chest trauma: results of a prospective study. J Trauma. 1997;43(3):405-11.

7. Brenner DJ. Medical imaging in the 21st century—getting the best bang for the rad. The New England journal of medicine. 2010;362(10):943-5.

8. Brenner DJ, Hall EJ. Computed tomography -an increasing source of radiation exposure. N Engl J Med. 2007;357:2277-84.

9. Lee J, Kirschner J, Pawa S, Wiener DE, Newman DH, Shah K. Computed tomography use in the adult emergency department of an academic urban hospital from 2001 to 2007. Ann Emerg Med. 2010;56(6):591-6.

10. Holmes JF, Wisner DH, McGahan JP, Mower WR, Kuppermann N. Clinical prediction rules for identifying adults at very low risk for intra-abdominal injuries after blunt trauma. Ann Emerg Med. 2009;54(4):575-84.

11. Exadaktylos AK, Sclabas G, Schmid SW, Schaller B, Zimmermann H. Do we really need routine computed tomographic scanning in the primary evaluation of blunt chest trauma in patients with "normal" chest radiograph? Journal of Trauma-Injury, Infection, and Critical Care. 2001;51(6):1173-6.

12. Sears BW, Luchette FA, Esposito TJ, et al. Old fashion clinical judgment in the era of protocols: is mandatory chest X-ray necessary in injured patients? J Trauma. 2005;59(2):324-32.

13. Rodriguez RM, Hendey GW, Marek G, Dery RA, Bjoring A. A pilot study to derive clinical variables for selective chest radiography in blunt trauma patients. Ann Emerg Med. 2006;47(5):415-8.

14. Bokhari F, Brakenridge S, Nagy K, et al. Prospective evaluation of the sensitivity of physical examination in chest trauma. Journal of Trauma-Injury, Infection, and Critical Care. 2002;53(6):1135-8.

15. Forouzanfar MM, Safari S, Niazazari M, et al. Clinical decision rule to prevent unnecessary chest X-ray in patients with blunt multiple traumas. Emerg Med Australas. 2014;26(6):561-6.

16. Rodriguez RM, Anglin D, Langdorf MI, et al. NEXUS chest: validation of a decision instrument for selective chest imaging in blunt trauma. JAMA Surg. 2013;148(10):940-6.

17. Michalke JA, Rocovich C, Patel T, et al. An overview of emergency ultrasound in the United States. World. 2012;3(2):85-90.

18. Brooks A, Davies B, Smethhurst M, Connolly J. Emergency ultrasound in the acute assessment of haemothorax. Emerg Med J. 2004;21(1):44-6.

19. Hyacinthe A-C, Broux C, Francony G, et al. Diagnostic Accuracy of Ultrasonography in the Acute Assessment of Common Thoracic Lesions After TraumaUltrasonography in Thoracic Trauma. Chest. 2012;141(5):1177-83.

20. Soldati G, Testa A, Sher S, Pignataro G, La Sala M, Silveri NG. Occult Traumatic PneumothoraxDiagnostic Accuracy of Lung Ultrasonography in the Emergency Department. Chest. 2008;133(1):204-11.

21. Soldati G, Testa A, Silva FR, Carbone L, Portale G, Silveri NG. Chest ultrasonography in lung contusion. Chest. 2006;130(2):533-8.

22. Zhang M, Liu Z-H, Yang J-X, et al. Rapid detection of pneumothorax by ultrasonography in patients with multiple trauma. Critical Care. 2006;10(4):R112.

23. McGahan JP, Wang L, Richards JR. From the RSNA Refresher Courses Focused Abdominal US for Trauma. Radiographics. 2001;21(suppl 1):S191-S9.

24. Poletti PA, Kinkel K, Vermeulen B, Irmay F, Unger PF, Terrier F. Blunt Abdominal Trauma: Should US Be Used to Detect Both Free Fluid and Organ Injuries? . Radiology. 2003;227(1):95-103.

25. Poletti PA, Mirvis SE, Shanmuganathan K, et al. Blunt abdominal trauma patients: can organ injury be excluded



This open-access article distributed under the terms of the Creative Commons Attribution NonCommercial 3.0 License (CC BY-NC 3.0). Copyright © 2016 Shahid Beheshti University of Medical Sciences. All rights reserved. Downloaded from: www.jemerg.com/w.SID.u/

without performing computed tomography? J Trauma Acute Care Surg. 2004;57(5):1072-81.

26. Meyer DM. Hemothorax related to trauma. Thorac Surg Clin. 2007;17(1):47-55.

27. Ma OJ, Mateer JR. Trauma ultrasound examination versus chest radiography in the detection of hemothorax. Ann Emerg Med. 1997;29(3):312-6.

28. Abboud P-AC, Kendall J. Emergency department ultrasound for hemothorax after blunt traumatic injury. J Emerg Med. 2003;25(2):181-4.

29. Ebrahimi A, Yousefifard M, Mohammad Kazemi H, et al. Diagnostic Accuracy of Chest Ultrasonography versus Chest Radiography for Identification of Pneumothorax: A Systematic Review and Meta-Analysis. Tanaffos. 2014;13(4):29-40.

30. Alrajab S, Youssef AM, Akkus NI, Caldito G. Pleural ultrasonography versus chest radiography for the diagnosis of pneumothorax: review of the literature and meta-analysis. Critical Care. 2013;17(5):R208.

31. Traub M, Stevenson M, McEvoy S, et al. The use of chest computed tomography versus chest X-ray in patients with major blunt trauma. Injury. 2007;38(1):43-7.

32. suárez Poveda T, Uribe CHM, Loaiza JR, Hurtado ÉHO, Rodríguez JPL, valencia Delgado AM. Chest Ultrasonography versus Chest CT for Diagnosis of posttraumatic residual hemothorax. Rev Colomb Radiol. 2012;23(2):3464-8.

33. Błasińska-Przerwa K, Pacho R, Bestry I. Pneumonol Alergol Pol. Pneumonologia Alergologia Polska. 2013;81(6):518-26.

34. Uz I, Yuruktumen A, Boydak B, et al. Impact of the practice of "Extended Focused Assessment with Sonography for Trauma" (e-FAST) on clinical decision in the emergency department. Turk J Trauma Emerg Surg. 2013;19(4):327-32. 35. Chardoli M, Hasan-Ghaliaee T, Akbari H, Rahimi-Movaghar V. Accuracy of chest radiography versus chest computed

rclu

tomography in hemodynamically stable patients with blunt chest trauma. Chin J Traumatol. 2013;16(6):351-4.

36. Leblanc D, Bouvet C, Degiovanni F, et al. Early lung ultrasonography predicts the occurrence of acute respiratory distress syndrome in blunt trauma patients. Intensive Care Med. 2014;40(10):1468-74.

37. Langdorf MI, Medak AJ, Hendey GW, et al. Prevalence and Clinical Import of Thoracic Injury Identified by Chest Computed Tomography but Not Chest Radiography in Blunt Trauma: Multicenter Prospective Cohort Study. Ann Emerg Med. 2015.

38. Vafaei A, Hatamabadi HR, Heidary K, Alimohammadi H, Tarbiat M. Diagnostic Accuracy of Ultrasonography and Radiography in Initial Evaluation of Chest Trauma Patients. Emergency. 2015;3:[In press].

39. Sistrom CL, Mergo PJ. A simple method for obtaining original data from published graphs and plots. Am J Roentgenol. 2000;174(5):1241-4.

40. Whiting PF, Rutjes AW, Westwood ME, et al. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. Ann Intern Med. 2011;155(8):529-36.

41. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. BMJ. 2003;327(7414):557.

42. Majumdar S, Kumar PS, Pandit A. Effect of liquid-phase properties on ultrasound intensity and cavitational activity. Ultrason Sonochem. 1998;5(3):113-8.

43. Deeks JJ, Macaskill P, Irwig L. The performance of tests of publication bias and other sample size effects in systematic reviews of diagnostic test accuracy was assessed. J Clin Epidemiol. 2005;58(9):882-93.





Figure 2: Deeks' funnel plot asymmetry test for assessment of publication bias. P values < 0.05 were considered as significant. Ultrasonography (A); Radiography (B). ESS: Effective sample sizes.



Figure 3: Summary receiver operative curves (SROC) for ultrasound (A) and chest radiography (B) in detection of hemothorax. AUC: Area under the curve; SENS: Sensitivity; SPEC: Specificity.





Figure 4: Forest plot of screening performance characteristics of chest ultrasonography in detection of hemothorax. Sensitivity and specificity (A); Diagnostic likelihood ratio (DLR) (B). CI: Confidence interval.





Figure 5: Forest plot of screening performance characteristics of chest radiography in detection of hemothorax. Sensitivity and specificity (A); Diagnostic likelihood ratio (DLR) (B). CI: Confidence interval.

