

EVIDENCE BASE MEDICINE

Evolution of Fagan's Nomogram; a Commentary

Abdelrahman Ibrahim Abushouk^{1*}

1. Faculty of Medicine, Ain Shams University, Cairo, Egypt.

Received: february 2016; Accepted: February 2016

Cite this article as: Ibrahim Abushouk A. Evolution of Fagan's Nomogram; a Commentary. Emergency. 2016; NN(I):114-115.

Dear Editor

I read with interest your paper entitled "Pre and post-test probabilities and Fagan's nomogram" (1). I would like to add a note concerning an update on Fagan's Nomogram. Generally, the basic idea of most nomograms is having the scales of 3 variables in a manner that if you draw a straight line between 2 values, the 3rd value is found where the line intersects the 3rd scale (2). They were initially developed in the 1980s by Maurice D'Ocange. Nomograms remained popular in medical practice until the invention of pocket calculators and computers. Their use increased again with the introduction of evidence-based medicine in clinical practice. In a letter to the New England Journal of Medicine in July 1975, Dr Terry Fagan displayed a test characterization tool that went on to carry his name as the Fagan's nomogram (3). This nomogram is a simple application of the Bayes' theorem, which establishes a rule to calculate the post-test probability of a disease. However, Fagan's nomogram had a set of drawbacks that limited its use in clinical practice. These drawbacks included: 1- The original Bayes' theorem is designed to deal with odds ratios, not probabilities, so algebraic conversion is needed to calculate probability. 2- Most diagnostic tests are characterized in terms of sensitivity & specificity in the literature, which need special equations to be converted into likelihood ratios (4). Noticing these difficulties, in 2011, a group of researchers published a modern version of the nomogram that they named "Bayes' theorem nomogram". The new nomogram targeted the former problems using: A- Parallel lines for probability and odds on each side of the nomogram figure. B- The inner lip along the entire circle contains values for sensitivity and specificity that can be connected to calculate the likelihood ratio of a certain diagnostic test (5). As illustrated in figure (A), a pretest probability of 18% and a likelihood ratio (LR+) of 2.8 for a diagnostic test would give a posttest probability of 38%. Further advantages of the modern nomogram include: - In Rare




Figure 1: The modern Bayes' theorem nomogram with an example of probability calculation as shown by the green line.

disorders, having a very low pretest probability implies performing a diagnostic test with a fairly high LR (+). In Fagan's nomogram, the high values of LR are compressed in a tight portion over that scale (4), while in this model; a more spaced representation of high LR is feasible. - Replacing the linear form with a circular one works better for complex diagnostic protocols where addition of multiple arrows for different diagnostic tests may be required (5). Considering the mentioned superiorities of the Bayes' theorem nomogram over the conventional Fagan's nomogram, it is highly recommended for clinicians to use it in conducting diagnostic protocols and formulating therapeutic plans.

References

1. Safari S, Baratloo A, Elfil M, Negida A. Part 4: Pre and Post Test Probabilities and Fagan's Nomogram. EMERGENCY-An Academic Emergency Medicine Journal. 2016;4.
2. Thimbleby H, Williams D, editors. Using nomograms to reduce harm from clinical calculations. Healthcare Infor-

*Corresponding Author: Abdelrahman Ibrahim Abushouk; Ain Shams University, Cairo, Egypt. E-mail: abdelrahman.abushouk@med.asu.edu.eg.

- maths (ICHI), 2013 IEEE International Conference on; 2013: IEEE.
3. Fagan TJ. Letter: nomogram for Bayes theorem. The New England journal of medicine. 1975;293(5):257-.
 4. Miettinen OS, Caro JJ. Foundations of medical diagnosis: what actually are the parameters involved in Bayes' theorem? Statistics in medicine. 1994;13(3):201-9.
 5. Marasco J, Doerfler R, Roschier L. Doc, what are my chances. UMAP Journal. 2011;32:279-98.

Archive of SID

