

ORIGINAL RESEARCH

Characteristics of Traumatic Urogenital Injuries in Emergency Department; a 10-Year Cross-sectional Study

Babak Javanmard¹, Morteza Fallah-karkan^{1,2}, Mohammadreza Razzaghi^{2*}, Anahita Ansari Djafari¹, Saleh Ghiasy¹, Behzad Lotfi¹, Reza Vafaei³

1. Urology Department, Shohada e Tajrish Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran.
2. Laser Application in Medical Science Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.
3. Proteomics Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Received: July 2019; Accepted: September 2019; Published online: 9 November 2019

Abstract: **Introduction:** Urogenital system injuries (UGIs) are seen in 10% of adult cases with multiple trauma. Although UGIs are rarely life threatening, they can cause major long-term morbidities. This study aimed to evaluate the characteristics of traumatic UGIs in patients who were referred to emergency department following multiple traumas. **Methods:** This retrospective cross-sectional study was conducted on multiple trauma patients who were presented to emergency department during a 10-year period (2008-2017). All patients with kidney, ureter, bladder, urethra, or external genitalia injuries were studied. The patients' data were extracted from their clinical profiles. **Results:** Out of the 13598 admitted patients in our trauma center, UGIs were seen in 267 (1.9%) cases. The mean age of patients with UGIs was 27.3 ± 6.1 years (74.15% male). The highest incidence of UGI was seen in those aged between 21 and 30 years (39.7%) and motorcycle accidents (49%) was the most frequent cause of trauma. 221 patients had an unstable situation and were emergently transferred to operation room (13.57% with traumatic kidney injury). The most common injured sites of urogenital system were kidney with 155 (58%) cases, followed by external genitalia with 91 (34.1%) cases. 77.5% of cases were managed conservatively and the rest (22.5%) underwent surgical procedures. **Conclusion:** UGIs comprise a low percentage (2%) of traumatic injuries, which are mostly caused by blunt trauma due to road traffic accidents. Kidney is the most common injured organ and UGIs mostly happen in young ages.

Keywords: Urogenital system; acute kidney injury; wounds and injuries; multiple trauma; epidemiology; mortality

Cite this article as: Javanmard B, Fallah-karkan M, Razzaghi M, Ansari Djafari A, Ghiasy S, Lotfi B, Vafaei R. Characteristics of Traumatic Urogenital Injuries in Emergency Department; a 10-Year Cross-sectional Study. Arch Acad Emerg Med. 2019; 7(1): e63.

1. Introduction

With economic development, incidence of traffic and industry-related accidents, iatrogenic injury, intoxication and also psychological self-mutilation have been increasing (1-8). Morbidity due to trauma has been one of the most serious public health concerns all over the world (9). In most reports, urogenital system injuries (UGIs) are seen in approximately 10% of adult traumatic patients and in less than 3% of children with multiple/severe trauma of lower abdomen or pelvis (10-12). Kidney is the most common injured organ in

genitourinary system while ureteral traumas are rare (1, 13). Although UGIs are rarely life threatening, they can cause significant long-term morbidities such as renal function impairment, sexual dysfunction and urethral stricture (14-16). Finding an ideal management in patients with UGIs requires comprehensive epidemiological information in different populations and times (14, 17). Therefore, this study aimed to evaluate the characteristics of traumatic UGIs in patients who were referred to emergency department following multiple traumas.

2. Methods

2.1. Study design and setting

This retrospective cross-sectional study was conducted on multiple trauma patients who were presented to emergency department of Shohada-e-Tajrish Hospital, Tehran, Iran, dur-

* **Corresponding Author:** Mohammadreza Razzaghi; Laser Application in Medical Science Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran. Tel: +98 9121482592, Fax: +98 2122712234, Email: M-razzaghi@sbmu.ac.ir.



ing a 10-year period (2008-2017). This hospital is the main referral trauma center in north and northeast of Tehran. The Ethics committee of Shohada-e-Tajrish Hospital approved this study and gave us permission to review patients' medical data. Considering the retrospective design of the study, the ethical committee ceded the necessity to have patients provide consent for collecting their medical data. The patients' data were collected confidentially and de-identified.

2.2. Participants

All admitted patients for more than 6 hours with any kind of traumatic UGIs such as kidney, ureter, bladder, urethra, external genitalia and adrenal gland injuries, were included using a census sampling method. Patients with incomplete medical profiles or unclear evidence regarding the urogenital tract were excluded. There was not any age or gender limitation.

2.3. Data gathering

We reviewed all information regarding demographic characteristics (age, gender), history, physical examination, mechanism of injury, type and severity of UGIs, associated injuries, done procedures, and also type of management (conservative or surgical managements) by referring to the medical records of the cases and filling out a checklist designed for the investigation. The type and mechanism of the injury were adapted according to the International Classification of Diseases, 10th revision (ICD-10)(18). Two senior urology residents (MFK and SG) were responsible for data gathering.

2.4. Statistical Analysis

Statistical analyses were performed by the Statistical Package of Social Science System software, version 20. Findings were presented using descriptive statistics as mean ± standard deviation or frequency (%).

3. Results

3.1. Baseline characteristics of studied patients

Out of 13598 admitted patients in our trauma center during the study period, UGIs were seen in 267 (1.9%) patients (92.5% blunt and 7.5% penetrating trauma). Table 1 shows the baseline characteristics of studied patients. The mean age of patients with UGIs was 27.3 ± 6.1 (3 - 82) years (74.15% male). The highest incidence of UGI was seen in those aged between 21 and 30 years (39.7%) followed by 11-20 years (25.07%) and 3 -10 years (19.1%) age groups. Motorcycle accidents (49%) followed by falling down (26%) were the most frequent causes of trauma. 221 patients had unstable situation and were emergently transferred to operation room (13.57% with traumatic kidney injury). 7 (2.8%) cases died due to severity of associated injuries (5 patients had renal

Table 1: Baseline characteristics of studied patients

Variables	Number (%)
Gender	
Male	198 (74.15)
Female	69 (25.8)
Age (year)	
3 – 10	51 (19.1)
11 –20	67 (25.07)
21 –30	106 (39.7)
30 – 60	31 (11.4)
>60	12 (4.73)
Trauma mechanism	
Motorcycle accident	131(49)
Falling down	69 (26)
Car accident	58 (21.6)
Sports accident	9 (3.3)
Hemodynamics	
Stable	238 (89.2)
Unstable	29 (10.8)
Associated injuries	
Abdominopelvic	104 (38.8)
Extremities	85 (31.9)
Thoracic	46 (17.1)
Head and neck	32(12.2)
Presenting sign and symptom	
Hematuria	169 (63.2)
Flank pain	50 (19.1)
Flank or genital abrasion	19 (7.1)
Obstruction	15 (5.6)
Other	14 (5.2)

and 2 cases had bladder injuries). The most common injured sites of urogenital system were kidney with 155 (58%) cases, followed by external genitalia with 91 (34.1%) cases (table 2). The common associated injuries were extremity (52.8%) and abdominopelvic (41.9%) injuries.

3.2. Management strategies

Table 3 summarizes different types of treatment strategies used for management of injured patients.

Upper urinary tract

Conservative management was applied for 51.8% of patients with renal injury and 2 patients with grade IV renal injury underwent angiographic embolization. Two patients with adrenal trauma were managed conservatively.

Lower urinary tract

Nineteen cases out of 27 patients with extra peritoneal bladder rupture were managed conservatively, by foley catheter insertion for two weeks. In one case with extra peritoneal bladder rupture, during orthopedic surgery bladder was injured intra peritoneally so cystoraphy was done intraoperatively. Fourteen of 25 patients with testis trauma had intact tunica albuginea and underwent conservative management. Three patients needed dermal flap to reform scrota because

Table 2: Distribution of urogenital injuries among the studied subjects

Injured organ	Number (%)
Upper urinary tract	
Renal	155 (58.1)
Ureter	3 (1.1)
Adrenal	2 (0.7)
Lower urinary tract	
Scrotum	43 (16.1)
Testis	25(9.3)
Urethra	15(5.6)
Bladder	34 (12.7)
Genitalia	25 (9.9)
Severity of renal injury	
Grade 1	103 (38.5)
Grade 2	34 (12.7)
Grade 3	10 (3.7)
Grade 4	5 (1.8)
Grade 5	3 (1.1)
Type of bladder injury	
Intra peritoneal injury	7 (2.6)
Extra peritoneal injury	27 (10.1)
Type of ureteral injuries	
Partial rupture of urethra	10(3.7)
Complete rupture of urethra	5(1.8)

Table 3: Management strategies of studied patients

Variables	Number (%)
Type of management	
Conservative management	207(77.5)
Operative management	52(19.4)
Type of procedures	
Nephrectomy	7(2.6)
Bladder repair	8(2.9)
Immediate endoscopic realignment	10(3.7)
Renoraphy	10(3.7)
Ureter repair	2(0.7)
Partial orchiectomy	5(1.8)
Male genital organ repair	8(2.9)
Female genital organ repair	2(0.7)

of extensive external genitalia injuries. Nine men and one girl with external genitalia injuries were managed conservatively. Ten of fifteen patients with urethral injuries underwent primary realignment (seven cases had complete urethral injuries). For five patients cystostomy was performed and they became candidates for delayed treatment.

4. Discussion

According to our results, the frequency of UGIs in this series of cases was 1.9%. The most common injured sites of urogenital system were kidney with 155 (58%) cases, followed by external genitalia with 91 (34.1%) cases. 77.5% of cases were managed conservatively. Appropriate data are needed to

assess clinical presentation, therapeutic procedure, trauma systems issues, and quality of care for this health problem (19). According to burden of disease and injuries, 28% of years of life lost in Iran is attributed to accidents (20). This issue may point out that Emergency Medical Service is not able to provide trauma patients with efficient and perfect pre-hospital care (21). The condition of trauma possibly depends on the space from hospital to high ways, socioeconomic status, grade of hospital as trauma center, and public health infrastructure mainly in prehospital trauma care (22, 23). Abundance of construction projects may be the reason that falling down is the second most common cause of injuries.

According to the obtained results from this study, UGIs comprise a low proportion of injuries in trauma patients (267 of 24742 patients, about 2 %). This is similar to the findings of other studies (10, 14, 24-26). Presentations of injured patients were similar to those in the literature. Regarding gender, the males enrolled were four times the females, and mainly prevalent in the 3rd decade of life (14, 17, 27-29). This may owe to men working in more risky jobs, being more involved in violent behavior and also number of male drivers being more than females (30). Most traffic accidents involved middle-aged victims, while the pedestrian accidents involved older age patients.

Road traffic accidents are the cause of the majority of UGIs; trauma had a blunt nature in 92.5% of patients, which is almost similar to other studies (13, 31-34). Moreover, kidney was the most common injured organ and kidney repair was the most common surgical treatment in this study. The haemodynamic condition is the point of reference for the diagnostic and therapeutic approach in trauma patients. Haemodynamically unstable patients need urgent laparotomy (35).

Initial treatment of patients with posterior urethral injuries (PUI) is one of the most challenging and technically difficult problems in urology (36). The controversy in treatment of PUI is surrounding two basically different approaches; one is the insertion of suprapubic cystostomy followed by a delayed urethroplasty; and the other approach is concomitant one stage suprapubic cystostomy and urethral realignment (37, 38). Advances in endoscopic techniques have facilitated early realignment and transurethral catheterization as a new treatment approach. In the present study, two thirds of the urethral injury cases underwent primary realignment and the rest became delayed urethroplasty candidates after cystostomy. Skin and tissue defects are another problem in patients with trauma. Grafts can be utilized for extended skin and tissue defects (39, 40). Complete avulsion of the scrotum and penis skin, creates a very serious life-threatening problem. Also, there is a serious problem in saving the testes, covering the penis and rehabilitation of area function.



There are two types of bladder ruptures: intraperitoneal and extraperitoneal. The extraperitoneal ruptures are often associated with pelvic fractures. Intraperitoneal ruptures often happen as a result of frontal impact, with unexpected deceleration while having a full bladder (24). The two most common symptoms of bladder injuries are macroscopic hematuria and abdominal tenderness. Other findings consist of inability to void, bruising of the supra-pubic region, and contrast extravasations in computed tomography (CT) urography. Pelvic fractures, which are associated with multiple organ injuries, including bladder, have a considerable mortality rate (20–40 %) (11, 41). According to our findings, due to the low rate of positive imaging findings in stable trauma patients with microscopic hematuria, it seems that there is no need to routinely perform abdominopelvic CT scan with intravenous contrast in these patients.

Since most cases with urogenital trauma have multiple injuries, they need a multidisciplinary approach; disregarding these injuries may cause severe complications. We propose establishing an integrated trauma system in Iran to improve the quality of trauma care. Also, preventive programs aimed at behavioral modifications in the society should be performed to decrease the health and economic burdens imposed by accidents.

5. Limitation

The first limitation of this investigation is the retrospective nature of the study, which caused some missing medical history and physical examination data. The second limitation is the lack of follow up assessments of disease adjusted life years.

6. Conclusion

UGIs mostly result from blunt trauma due to road traffic accidents. Renal injuries are the most common; and UGI mostly develops in youth. Conservative management can be the treatment of choice for renal trauma patients who are hemodynamically stable.

7. Appendix

7.1. Acknowledgements

We would like to show our appreciation towards urology and emergency ward staff for their invaluable help throughout this study. We gratefully appreciate medical record department staff of Shohada e Tajrish hospital for their kind cooperation. We would like to express our sincere gratitude to Dr. Saeed Montazeri for his valuable assistance in English editing of the manuscript.

7.2. Author contribution

Study concept and design: All Authors.

Acquisition of data: Dr. Fallah-karkan, Dr. Ghiasi

Statistical analysis: Dr. Ghiasi, Dr. Vafae.

Drafting of the manuscript: Dr. Fallah-karkan, Dr. Ghiasi and Dr. Ansari.

Critical revision of the manuscript: All Authors.

Authors ORCIDs

Babak Javanmard: 0000-0002-0327-8548

Morteza Fallah-karkan: 0000-0001-8788-2760

Mohammadreza Razzaghi: 0000-0003-1429-1171

Anahita Ansari Djafari: 0000-0001-6392-1323

Saleh ghiasy: 0000-0002-1835-7441

Behzad Lotfi: 0000-0001-8835-8151

Reza Vafae: 0000-0003-0677-4755

7.3. Funding/Support

None.

7.4. Conflict of interest

The authors declare that there is no conflict of interest.

References

1. Zou Q, Fu Q. Diagnosis and treatment of acute urogenital and genitalia tract traumas: 10-year clinical experience. *Pakistan journal of medical sciences*. 2015;31(4):925.
2. Javanmard B. Recurrent hematuria in renal angio-venous malformation, delay diagnosis and endovascular treatment, a case report. *Urology case reports*. 2018;16:46.
3. Razzaghi MR, Rezaei A, Mazloomfard MM, Javanmard B, Mohammadhosseini M, Rezaei I. Successful macro-surgical reimplantation of an amputated penis. *Urology journal*. 2009;6(4):306-8.
4. Rahbar Taramsari M, Badsar A, Naghipour M, Dvam F, Shahrani H, Saeidin A, et al. Assessment of pre-hospital emergency performance in missions of 115 emergency bases of Rasht, Iran. *Annals of Biological Research*. 2013;4(12):75-81.
5. Javanmard B, Yousefi MR, Fadavi B, Karkan MF. Retained Surgical Gauze Presenting With Gross Hematuria: A Case Report. *Urology journal*. 2017;14(5):5027-9.
6. Aghajanzadeh M, Dehnadi A, Ebrahimi H, Karkan MF, Jahromi SK, Maafi AA, et al. Classification and management of subcutaneous emphysema: a 10-year experience. *Indian Journal of Surgery*. 2015;77(2):673-7.
7. Behboudi H, Moghadam RS, Tiefef N, Karkan MF. Vision disorders in drivers involved in traffic accidents. *Journal of ophthalmic & vision research*. 2017;12(4):451.
8. Taramsari MR, Badsar A, Hamrangnejad M, Saeidin A, Karkan MF. Alcohol Intoxication in Gilan Province

- Hospital Center of Toxicology, Iran: Demographic, Patterns and Outcome. Archives of Advanced Sciences. 2017;1(1):10-4.
9. Smith GS, Barss P. Unintentional injuries in developing countries: the epidemiology of a neglected problem. Epidemiologic reviews. 1991;13(1):228-66.
 10. Salimi J, Nikoobakht MR, Zareei M. Epidemiologic study of 284 patients with urogenital trauma in three trauma center in Tehran. Urology journal. 2009;1(2):117-20.
 11. Pereira BMT, De Campos CCC, Calderan TRA, Reis LO, Fraga GP. Bladder injuries after external trauma: 20 years experience report in a population-based cross-sectional view. World journal of urology. 2013;31(4):913-7.
 12. McAninch JW. Genitourinary trauma. Urologic Clinics. 2006;33(1):xiii.
 13. Laal M, Khodadadi M, Zarei M-R. Renal Trauma Management in 8 Cities of Iran. Acta Medica Iranica. 2009;47(1):61-4.
 14. Salimi J, Nikoobakht MR, Khaji A. Epidemiology of urogenital trauma: results of the Iranian national trauma project. Urology journal. 2009;3(3):171-4.
 15. Mirzazadeh M, Fallahkarkan M, Hosseini J. Penile fracture epidemiology, diagnosis and management in Iran: a narrative review. Translational andrology and urology. 2017;6(2):158.
 16. Allameh F, Najafi S, Karkan MF, Haghhighatkah H, Nekuie S. Chromophobe Renal Cell Carcinoma Presented with Wunderlich Syndrome. International Journal of Cancer Management. 2017;10(11).
 17. Sarvestani AS, Zamiri M. Assessment of genitourinary trauma in Southeastern Iran. Trauma monthly. 2013;18(3):113.
 18. Moore L, Clark DE. The value of trauma registries. Injury. 2008;39(6):686-95.
 19. Wynn A, Wise M, Wright MJ, Rafaat A, Wang Y-Z, Steeb G, et al. Accuracy of administrative and trauma registry databases. Journal of Trauma and Acute Care Surgery. 2001;51(3):464-8.
 20. Zargar M, Kalantar Motamedi SMR, Karbakhsh M, Ghodsi SM, Rahimi-Movaghar V, Panahi F, et al. Trauma care system in Iran. Chinese Journal of Traumatology. 2011;14(3):131-6.
 21. Khashayar P, Amoli HA, Tavakoli H, Panahi F. Efficacy of pre-hospital care in trauma patients in Iran. Emergency Medicine Journal. 2010;27(6):430-2.
 22. Zafarhandi M-R, Modagheh M-HS, Roudsari BS. Preventable trauma death in Tehran: an estimate of trauma care quality in teaching hospitals. Journal of Trauma and Acute Care Surgery. 2003;55(3):459-65.
 23. Sehat M, Naieni KH, Asadi-Lari M, Foroushani AR, Malek-Afzali H. Socioeconomic status and incidence of traffic accidents in metropolitan Tehran: a population-based study. International journal of preventive medicine. 2012;3(3):181.
 24. Paparel P, N'DIAYE A, Laumon B, CAILLOT JL, Perrin P, Ruffion A. The epidemiology of trauma of the genitourinary system after traffic accidents: analysis of a register of over 43 000 victims. BJU international. 2006;97(2):338-41.
 25. McGeady JB, Breyer BN. Current epidemiology of genitourinary trauma. The Urologic clinics of North America. 2013;40(3):323.
 26. Schmid CW, Schmid DM. Traumatic Injury of the Urogenital System. Head, Thoracic, Abdominal, and Vascular Injuries: Springer; 2011. p. 351-68.
 27. Bagga HS, Tasian GE, Fisher PB, McCulloch CE, McAninch JW, Breyer BN. Product related adult genitourinary injuries treated at emergency departments in the United States from 2002 to 2010. The Journal of urology. 2013;189(4):1362-8.
 28. Terrier J-E, Paparel P, Gadegbeku B, Ruffion A, Jenkins LC, N'diaye A. Genitourinary injuries after traffic accidents: Analysis of a registry of 162,690 victims. Journal of trauma and acute care surgery. 2017;82(6):1087-93.
 29. Roudsari BS, Sharzei K, Zargar M. Sex and age distribution in transport-related injuries in Tehran. Accident Analysis & Prevention. 2004;36(3):391-8.
 30. Stiell IG, Clement CM, Rowe BH, Schull MJ, Brison R, Cass D, et al. Comparison of the Canadian CT Head Rule and the New Orleans Criteria in patients with minor head injury. Jama. 2005;294(12):1511-8.
 31. Kuo RL, Eachempati SR, Makhuli MJ, Reed Li RL. Factors affecting management and outcome in blunt renal injury. World journal of surgery. 2002;26(4):416-9.
 32. Santucci R, Wessells H, Bartsch G, Descotes J, Heyns C, McAninch J, et al. Evaluation and management of renal injuries: consensus statement of the renal trauma subcommittee. BJU international. 2004;93(7):937-54.
 33. Kuan JK, Wright JL, Nathens AB, Rivara FP, Wessells H. American Association for the Surgery of Trauma Organ Injury Scale for kidney injuries predicts nephrectomy, dialysis, and death in patients with blunt injury and nephrectomy for penetrating injuries. Journal of Trauma and Acute Care Surgery. 2006;60(2):351-6.
 34. Bariol S, Stewart G, Smith R, McKeown D, Tolley D. An analysis of urinary tract trauma in Scotland: impact on management and resource needs. The Surgeon. 2005;3(1):27-30.
 35. Aragona F, Pepe P, Patane D, Malfa P, D'arrigo L, Pennisi M. Management of severe blunt renal trauma in adult patients: a 10-year retrospective review from an emergency hospital. BJU international. 2012;110(5):744-8.
 36. Chapple C, Barbagli G, Jordan G, Mundy A, Rodrigues-Netto N, Pansadoro V, et al. Consensus statement on ure-



- thral trauma. *BJU international*. 2004;93(9):1195-202.
37. Hadjizacharia P, Inaba K, Teixeira PG, Kokorowski P, Demetriades D, Best C. Evaluation of immediate endoscopic realignment as a treatment modality for traumatic urethral injuries. *Journal of Trauma and Acute Care Surgery*. 2008;64(6):1443-50.
 38. Balkan E, Kilic N, Dogruyol H. The effectiveness of early primary realignment in children with posterior urethral injury. *International journal of urology*. 2005;12(1):62-6.
 39. Brown JB, Fryer MP. Peno-Scrotal Skin Losses, Repaired by Implantation and Free Skin Grafting: Report of Known Normal Offspring:(Preliminary Report on Total and Deep Losses). *Annals of surgery*. 1957;145(5):656.
 40. Karian LS, Chung SY, Lee ES. Reconstruction of defects after Fournier gangrene: a systematic review. *Eplasty*. 2015;15.
 41. Wirth GJ, Peter R, Poletti PA, Iselin CE. Advances in the management of blunt traumatic bladder rupture: experience with 36 cases. *BJU international*. 2010;106(9):1344-9.