



## What's New in Treatment of *Helicobacter pylori*?

Masoud Mardani<sup>1,\*</sup>

<sup>1</sup>Infectious Diseases and Tropical Medicine Research Center, Shahid Beheshti University of Medical Sciences, Tehran, IR Iran

\*Corresponding author: Masoud Mardani, Infectious Diseases and Tropical Medicine Research Center, Shahid Beheshti University of Medical Sciences, Tehran, IR Iran. E-mail: drmasoudmardani@yahoo.com

Received 2017 November 11; Accepted 2017 December 22.

**Keywords:** Treatment, Eradication, *Helicobacter pylori*

*Helicobacter pylori* are Gram-negative, microaerophilic bacteria, and ubiquitous organisms, which are present in about 50% of the global population (1). Metaplastic changes in the stomach is a result of chronic infection with *H. pylori* infection, and can lead to peptic ulcer. The most common route of *H. pylori* infection is oral-to-oral or fecal-to-oral contact (2-4).

Generally, *H. pylori* is an asymptomatic disease with no specific clinical signs; possible symptoms, includes nausea, vomiting, abdominal pain, heartburn, diarrhea, hunger in the morning, and halitosis (5). Also, recent studies proved the role of *H. pylori* in the development of endocrinopathies (6). In the new guideline on the treatment of *Helicobacter pylori* by the Canadian association of gastroenterology and the Canadian *Helicobacter* study group, it is suggested that all *H. pylori* eradication regimens should be given for 14 days to replace the 10-day treatment (7).

Patient suspected of *H. pylori*, following laboratory study, should be done: first of all, fecal antigen test of *H. pylori* with 98% specificity and 94% sensitivity was performed, and in the next step, positive results obtained in the early stage of disease should be used in order to detect post-treatment eradication (7-9). Secondly, the Carbon 13 urea breath test should be done and when urease is present in the stomach, concentration of labeled carbon is high and will react with *H. pylori* infection (7, 9). Thirdly, although serology of *H. pylori* is not a good test for follow-up of treated patients, it has high specificity and sensitivity (> 90%), thus is useful for detecting new infected patients (10). Finally, anti-biogram is useful in geographic areas with a high resistance rate against metronidazole and clarithromycin (7).

Recommended first-line strategy include concomitant non-bismuth quadruple therapy (Proton Pump Inhibitor [PPI] + Amoxicillin + Metronidazole + Clarithromycin

[PAMC]) and traditional bismuth quadruple therapy (PPI + Bismuth + Metronidazole + Tetracycline [PBMT]) (7).

Also, it is recommended that PPI triple therapy (PPI + clarithromycin + either amoxicillin or metronidazole) is restricted to areas with known low Clarithromycin resistance (< 15%) or high eradication success with these regimens (> 85%), and rescue therapies include PBMT and levofloxacin-containing therapy (PPI + amoxicillin + levofloxacin). Rifabutin regimes should be restricted to patients, who have failed to respond to at least three prior options (7).

In Iran, due to increasing demand of therapy of *H. pylori* with short course treatment, it is highly recommended for all *H. pylori* eradication regimens to now last for 14 days.

### Acknowledgments

The author would like to thank Bita Pourkaveh for her contribution in collecting the data.

### References

- Sanders KM, Stern TA, O'Gara PT, Field TS, Rauch SL, Lipson RE, et al. Delirium during intra-aortic balloon pump therapy. Incidence and management. *Psychosomatics*. 1992;**33**(1):35-44. doi: 10.1016/S0033-3182(92)72019-2. [PubMed: 1539101].
- Testerman TL, Morris J. Beyond the stomach: an updated view of *Helicobacter pylori* pathogenesis, diagnosis, and treatment. *World J Gastroenterol*. 2014;**20**(36):12781-808. doi: 10.3748/wjg.v20.i36.12781. [PubMed: 25278678].
- Shahabimehr M, Alebouyeh M, Farzi N, Mahboubi A, Taslimi R, Zali MR. Resistance Rate and Minimum Inhibitory Concentration of Metronidazole Among *Helicobacter pylori* Strains in Tehran, Iran. *Arch Clin Infect Dis*. 2016;**11**(2).
- Fayaz F, Gachkar L, Roodsari S, Pourkaveh B. Prevalence of *Helicobacter pylori* infection in patients with Down syndrome and mental retardation. *Arch Clin Infect Dis*. 2014;**9**(4).
- Frias Ordóñez JS, Otero Regino W. [Practical issues in diagnostic tests for *Helicobacter pylori* infection: a narrative review]. *Rev Gastroenterol Peru*. 2017;**37**(3):246-53. [PubMed: 29093589].

6. Kiani AH, Asadbeik E, Bibalan M, Sedighi M, Eshaghi M, Gholami M, et al. Serological diagnosis of *Helicobacter pylori* infection in patients with a polycystic ovary syndrome. *Arch Clin Infect Dis*. 2015;**10**(2).
7. Fallone CA, Chiba N, van Zanten SV, Fischbach L, Gisbert JP, Hunt RH, et al. The Toronto Consensus for the Treatment of *Helicobacter pylori* Infection in Adults. *Gastroenterology*. 2016;**151**(1):51-69 e14. doi: [10.1053/j.gastro.2016.04.006](https://doi.org/10.1053/j.gastro.2016.04.006). [PubMed: [27102658](https://pubmed.ncbi.nlm.nih.gov/27102658/)].
8. da Silva-ETTO JMK, Mattar R, Villares-Lopes CA, Marques SB, Carilho FJ. Evaluation of diagnostic accuracy of two rapid stool antigen tests using an immunochromatographic assay to detect *Helicobacter pylori*. *Clin Biochem*. 2017;**50**(16-17):959-62. doi: [10.1016/j.clinbiochem.2017.05.005](https://doi.org/10.1016/j.clinbiochem.2017.05.005). [PubMed: [28483407](https://pubmed.ncbi.nlm.nih.gov/28483407/)].
9. Biranjia-Hurdoyal SD, Seetulsingh-Goorah SP. Performances of Four *Helicobacter pylori* Serological Detection Kits Using Stool Antigen Test as Gold Standard. *PLoS One*. 2016;**11**(10). e0163834. doi: [10.1371/journal.pone.0163834](https://doi.org/10.1371/journal.pone.0163834). [PubMed: [27736910](https://pubmed.ncbi.nlm.nih.gov/27736910/)].
10. Kim JH, Lee SY, Kwon HG, Lee SP, Kim JH, Sung IK, et al. [Serological Changes after Equivocal *Helicobacter pylori*-serology Test Findings Dependent on Gastric Secreting Ability]. *Korean J Gastroenterol*. 2017;**70**(4):181-9. doi: [10.4166/kjg.2017.70.4.181](https://doi.org/10.4166/kjg.2017.70.4.181). [PubMed: [29060956](https://pubmed.ncbi.nlm.nih.gov/29060956/)].