Published online 2019 August 19.

Research Article

Association Between Smoking, Alcohol Consumption, and Dietary Habits in Northern Thai Adolescents in Thailand, 2016 - 2017

Nonglak Boonchooduang¹, Narueporn Likhitweerawong¹ and Orawan Louthrenoo ¹,*

¹Department of Pediatrics, Faculty of Medicine, Chiang Mai University, Chiang Mai, Thailand

* Corresponding author: Department of Pediatrics, Chiang Mai University, 50200, Chiang Mai, Thailand. Tel: +66-53935412, Fax: +66-53936461, Email: orawan.l@cmu.ac.th Received 2018 March 06; Revised 2019 June 22; Accepted 2019 July 01.

Abstract

Background: Smoking and alcohol consumption have been common among adolescents in recent years, both of which are risk factors of cardiovascular disease. They may be associated with unhealthy dietary habits that can add to cardiovascular risks in adults. **Objectives:** This study aimed to evaluate the association between smoking status, alcohol consumption, and dietary habits in adolescents.

Methods: In this study, 4,372 adolescents (2,223 males and 2,149 females) aged 13 to 18 years were enrolled from randomly selected schools in Northern Thailand. They were asked to complete the Thai Youth Risk Behavior Survey regarding smoking, alcohol use, and dietary habits.

Results: The mean age of the participants was 15.4 (1.7) years and 15.0 (1.6) years in males and females, respectively. The prevalence of smokers and alcohol drinkers was 21% and 28.3%, respectively. Among alcohol drinkers, 52.4% were also smokers. Besides, 11% of the participants were current smokers. The proportions of adolescents with healthy dietary habits, defined as drinking 100% fruit juice, drinking milk, fruit/vegetable consumption, and having regular breakfast, were significantly higher in the non-smoking/non-drinking group ($P \le 0.05$ to $P \le 0.001$).

Conclusions: Adolescents who were smokers and alcohol drinkers were less likely to have healthy dietary habits. This can affect cardiovascular risks and associated diseases in the future. The proper advice for dietary and lifestyle modification among these high-risk groups may be needed from an early age.

Keywords: Smoking, Alcohol Drinking, Feeding Behavior, Adolescent, Cardiovascular Diseases

1. Background

Smoking and alcohol consumption are major health problems in adolescents in many countries. The use of tobacco and alcohol has been on the rise in adolescents worldwide. The youth risk behavior surveillance system (YRBSS) in the United States in 2015 indicated that during 30 days before the survey, 10.8% of high school students had smoked cigarettes and 32.8% had drunk alcohol (1). Among the youth, the short-term health consequences of smoking include respiratory effects and risk of other drug use (2). Individual, environmental, and family factors including family members currently smoking synergistically act to motivate adolescents toward smoking (3, 4). While alcohol consumption among adolescents has probably slightly declined over the past decade, the pattern of drinking too much and too early in adolescents has increased public concern (5). Worldwide, 38.3% of the populations aged 15 years or older have a history of alcohol consumption in the past 12 months (6).

Tobacco use and alcohol use are both strong predictors of cardiovascular risks leading to increased morbidity and mortality in adults. Cigarette smoking has an impact on hypertension, diabetes mellitus, dyslipidemia, and stroke in adults (2, 7, 8). Studies have shown that the early signs of these diseases can be found in adolescents who smoke (2, 9). In the same way, several studies have reported the association of alcohol consumption, even in moderate ranges, with cardiovascular diseases including hypertension and coronary artery disease (5, 10, 11). Unhealthy dietary behavior can also be associated with an overall unhealthy lifestyle, including time spent watching television (12), to contribute to overweight and obesity. Consequently, it has added to the risks of cardiovascular diseases. On the other hand, healthy dietary habits are promoted such as 100% fruit juice drinking, fruit/vegetable consumption, not consuming soft drinks and sports drinks, and having regular breakfast (13-15).

Few studies have reported some associations between

Copyright © 2019, Journal of Comprehensive Pediatrics. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited. tobacco use, alcohol consumption, and unhealthy dietary habits in adolescents and adults. Adolescent smokers were found to consume more fast food and soft drinks while they skipped several regular meals than non-smokers (16). Similarly, smokers more frequently consume fewer fruits and vegetables (17). Poor lifestyles including smoking status, alcohol drinking, and unhealthy dietary habits leading to cardiovascular risks can occur at an early age.

2. Objectives

This study aimed to study the association between smoking status, alcohol consumption, and dietary habits among Northern Thai adolescents so that appropriate early preventive programs could be provided to this population.

3. Methods

3.1. Participants

A cross-sectional survey was conducted in secondary and vocational schools randomly chosen across Chiang Mai, a city in Northern Thailand. We invited 5,639 students to participate in this study. A self-reported web-based questionnaire was administered to students in grades 7 to 12 (aged 13 - 18 years). School teachers obtained parental consent verbally and provided information regarding the study to students after a regular computer class period. All students participating in this study were assured that their responses were anonymous and voluntary. The participants were also asked to indicate their consent on the first page of the web-based questionnaire. The study was approved by the Research Ethics Committee of the Faculty of Medicine of Chiang Mai University.

3.2. Measurements

Information regarding smoking status, alcohol use, and dietary habits was obtained though the Thai youth risk behavior survey (Thai-YRBS), a school-based survey that monitors major risk-taking behaviors. The Thai-YRBS was derived from the 2015 youth risk behavior survey (YRBS) developed by the center for disease control and prevention (CDC) (18).

3.3. Definition

Smoking status and alcohol drinking were determined by the following questions:

"Have you ever tried cigarette smoking, even one or two puffs?" Participants who answered "yes" to this question were defined as ever smoking and those who answered "no" were defined as non-smoking. "During the past 30 days, on how many days did you smoke cigarettes?" Current smoking was defined as smoking a cigarette on any of the days in the past 30 days.

"During your life, on how many days have you had at least one drink of alcohol?" Ever drinking was defined as alcohol drinking on any occasion in life.

"During the past 30 days, on how many days did you have at least one drink of alcohol?" Current drinking was defined as alcohol drinking on any of the days in the past 30 days.

Dietary habits were determined by the following questions:

"During the past seven days, how many times did you drink 100% fruit juices"? Drinking 100% fruit juice was defined as drinking 100% fruit juice at least one time during the past seven days.

Similar questions were applied to fruit/vegetable consumption, soft drinks/sport drinks intake, drinking plain water, drinking milk, and having regular breakfast.

3.4. Statistical Analysis

Data were collected to evaluate the association between smoking status, alcohol consumption, and dietary habits in adolescents. The SPSS version 22.0 program (IBM Corp., Armonk, NY) was used to perform statistical analyses. The percentages, means, and standard deviations were calculated. Differences between study groups were analyzed using the chi-square test for qualitative parameters. A P value of less than 0.05 was considered statistically significant.

4. Results

In total, 2,223 male and 2,149 female students participated in this study, providing a response rate of 77%. The prevalence of smokers and alcohol drinkers was 21% and 28.3%, respectively. Among alcohol drinkers, 52.4% were also smokers. Moreover, 11.1% of the participants reported that they were current smokers whereas 18.6% reported current alcohol drinking. The characteristics of the study population are shown in Table 1.

Regarding dietary habits, non-smokers comprised a significantly higher proportion of students consuming healthy products and having dietary habits than smokers, except for drinking plain water. Likewise, non-alcohol drinkers consumed healthy foods and had healthy dietary habits more often than alcohol drinkers, as presented in Table 2.

Additionally, participants were classified into four groups including non-smoking/non-alcohol drinking, smoking only, alcohol drinking only, and both smoking

	Males, N = 2223	Females, N = 2149	
Age, mean \pm SD	15.4 ± 1.7	15.0 ± 1.6	
Age, y			
13 -15	1097 (49.3)	1294 (60.2)	
16 - 18	1126 (50.7)	855 (39.8)	
Type of school			
Public schools	396 (17.8)	689 (32.1)	
Private schools	1020 (45.9)	1301 (60.5)	
Vocational schools	807 (36.3)	159 (7.4)	
Smoking			
Current smoking	407 (18.3)	79 (3.7)	
Ever smoking, not current	295 (13.3)	137(6.4)	
Non-smoking	1521 (68.4)	1933 (89.9)	
Alcohol drinking			
Current drinking	567 (25.5)	248 (11.6)	
Ever drinking, not current	244 (11.0)	179 (8.3)	
Non-drinking	1412 (63.5)	1722 (80.1)	

Table 1. Characteristics of Participants^a

^aValues are expressed as No. (%) unless otherwise indicated.

and alcohol drinking groups. The prevalence of nonsmoking/non-alcohol drinking adolescents was 65.5%, whereas that of smoking only, alcohol drinking only, and both smoking and alcohol drinking was 6.2%, 13.5%, and 14.8%, respectively. The proportion of adolescents with healthy dietary habits was significantly higher in nonsmoking/non-drinking group (P \leq 0.05 to P \leq 0.001), as shown in Figure 1.

5. Discussion

In this study, the prevalence of smokers and alcohol drinkers was 21% and 28.3% among Northern Thai adolescents, respectively, that is consistent with the prevalence of cigarette smoking (22.3%) and alcohol use (27.9%) reported by the 2013 survey in Thailand (19). In the same way, the prevalence of high school students with current use of tobacco product (20.2%) did not significantly change in the US during 2011 - 2016 (20). In this study, 11% of the participants reported current smoking. Smoking was three times more common in males than in females whereas a previous report showed that smoking was 10 times higher in males than in females (21). This finding indicates more recent female smokers. Among Southeast Asian countries, the age of first smoking of a whole cigarette was 14.6-yearsold in Thailand whereas it was 13.9-years-old in Malaysia (22).

Most alcohol drinkers had their first-time drinking at the age of 13-14 years, which was near the age of first smoking (23). Males are the gender at risk of health problems due to more episodes and more consumption of alcohol than females. However, similar to smoking, alcohol consumption among females reported by the National Statistical Office in Thailand gradually increased from 8.8% in 2007 to 11.8% in 2013 (24). Alcohol consumption was associated with recognized risks for non-communicable diseases. After adjustment for variables, the odds ratios of outcomes (high cholesterol, hypertension, and liver disease) were significantly greater among both occasional heavy drinkers (OR = 1.2 - 1.5) and regular heavy drinkers (OR = 1.5 - 2.0) compared to non-drinkers (25).

Concurrent use of tobacco and alcohol was 14.8% in this study. The prevalence of alcohol and cigarettes use was found to be 19% - 22% elsewhere (26). Smoking adolescents were reported to be three times more likely than nonsmokers to use alcohol (2). Based on previous studies, smoking and alcohol consumption are the most significant health risks at any age. Mortality from cardiovascular diseases due to coronary heart disease, hypertension, and diabetes is the most well-known consequence of tobacco and alcohol use (27-29). This mortality plays an important role in overall mortality of Thai population (38%smoking, 16%-hypertension, and 10%-diabetes) in a previous study (29). Accordingly, smoking and alcohol use may be considered to affect both vasculature and cardiovascular risks.

Previous studies indicated that adolescents usually were less likely to engage in healthy behaviors. In the 2015 YRBSS, the findings indicated that 5.2% of high school students had not eaten fruit or drunk 100% fruit juice and 6.7% had not eaten vegetables during the seven days before the survey (1). Similarly, 46% and 60% of the adolescents did not eat fruit and vegetables daily, respectively (30). The international data collected in five Southeast Asian countries showed that 76.3% of the youth aged 13 to 15 years had inadequate fruits and vegetables consumptions (fewer than five servings per day) (31). The lack of protective factors and being physically inactive were associated with inadequate fruit/vegetable consumption and sedentary behavior (31). Unhealthy dietary habits are associated with overweight and obesity and consequently, they add to the risks of cardiovascular disease.

This study showed that the proportion of adolescents with healthy dietary habits was significantly higher in the non-smoking/non-drinking group. Our findings are in line with other studies that evaluated smoking frequency and unhealthy lifestyles and reported the association of smoking frequency with having skipped meals and less healthy food. The smoking frequency was also directly related

Table 2. The Proportions of Dietary Habits According to Smoking and Alcohol Drinking Status										
Dietary Habits	Smoking (N = 918)	Non-Smoking (N = 3452)	OR (95%CI)	P Value	Alcohol Drinking (N = 1238)	Non-Alcohol Drinking (N = 3134)	OR (95%CI)	P Value		
100% fruit juice	136	616	0.84 (0.71 - 0.99)	0.03	186	566	0.85 (0.74 - 0.97)	0.02		
Fruit consumption	207	1022	0.74 (0.65 - 0.86)	< 0.001	289	940	0.78 (0.69 - 0.87)	< 0.001		
Vegetable consumption	301	1276	0.86 (0.76 - 0.98)	0.02	408	1169	0.87(0.79-0.96)	0.007		
No soft drinks intake	141	1040	0.49 (0.41 - 0.58)	< 0.001	232	949	0.62 (0.55 - 0.71)	< 0.001		
No sport drinks intake	416	2205	0.55 (0.49 - 0.62)	< 0.001	594	2027	0.62 (0.56 - 0.68)	< 0.001		
Plain water drinking	807	3094	0.87 (0.73 - 1.04)	0.13	1081	2820	0.83 (0.72 - 0.95)	0.01		
Milk drinking	383	1731	0.76 (0.68 - 0.86)	< 0.001	501	1613	0.73 (0.66 - 0.80)	< 0.001		
Regular breakfast	467	2154	0.68 (0.61 - 0.77)	< 0.001	646	1975	0.73 (0.66 - 0.80)	< 0.001		



Figure 1. Dietary habits and proportions of smoking and alcohol consumption status in Northern Thai adolescents. *P \leq 0.001.

to the frequency of fast-food and soft drink consumption (16). Adolescents who smoked were less likely to consume fruits, vegetables, and milk/dairy products and they ate significantly more fast-food than non-smokers (32-34). Daily smokers reported more consumption of food rich in sugar and fat than non-smoking adolescents (35). Adolescents who skipped breakfast were found to smoke more than breakfast consumers (14). Furthermore, another study added to previous findings that heavy smokers were significantly less likely to follow healthy dietary restrictions than non-smokers (36). In the same way, adolescents who drank alcohol were found to be less likely to consume healthy food than non-alcohol users (32), which is consistent with the current study. The findings of this study are attributed to the fact that smokers and alcohol drinkers are less likely to consider lifestyle and health consequences. A study in adults showed that eating behavior was associated with BMI, besides current smoking, ex-smoking status, and alcohol consumption (37).

This study demonstrated a relationship between smoking, alcohol consumption, and dietary habits among Northern Thai adolescents. The recruitment from multiple school centers yielded a large and diverse sample that brought an adequate effect size to identify the association between smoking, alcohol consumption, and dietary habits. Additionally, the design of questionnaires conferred several methodological advantages for evaluating associations as mentioned above.

For limitations, our sample included randomly selected schools with more than 1,000 students indirectly implying a relative socioeconomically advantage especially regarding food habits, so that the results may not represent all Thai adolescents.. Furthermore, the results apply only to the youth who attend schools and are not representative of all adolescents in this age group. Finally, we might not be definitely assuring that non-smokers/nondrinkers were associated with healthy dietary habits. A longer study with more depth would provide more information.

5.1. Conclusions

This study showed many adolescents in Northern Thailand with recent cigarette and alcohol use. Cardiovascular diseases seem to be growing at the early young age associated with smoking, alcohol drinking, and having less healthy lifestyles. Adolescents who were smokers and alcohol drinkers were less likely to have healthy dietary habits. This may affect cardiovascular risks and their sequential diseases in the future. Health-risks surveillance should be monitored for adolescent health. The proper advice for healthy dietary habits and lifestyle modification in these high-risk groups is needed from an early age.

Footnotes

Authors' Contribution: Nonglak Boonchooduang contributed to the study design, developed the data collection instruments, carried out the analysis, and reviewed the manuscript. Narueporn Likhitweerawong contributed to data analysis and first drafting of the manuscript. Orawan Louthrenoo contributed to study conception and design, supported data analysis and interpretation, and wrote the manuscript. All authors read and approved the final manuscript.

Conflict of Interests: The authors declare no conflicts of interest with respect to the research, authorship, and/or publication of this article.

Ethical Approval: The study was approved by the Ethics Research Committee of the Faculty of Medicine, Chiang Mai University.

Funding/Support: This study was supported by the Faculty of Medicine, Chiang Mai University (code number 453/2515).

References

- Kann L, McManus T, Harris WA, Shanklin SL, Flint KH, Hawkins J, et al. Youth risk behavior surveillance - United States, 2015. *MMWR Surveill Summ*. 2016;65(6):1-174. doi: 10.15585/mmwr.ss6506a1. [PubMed: 27280474].
- World Health Organization. *Tobacco free initiative (TFI)*. 2017, [cited 11 November]. Available from: http://www.who.int/tobacco/research/ youth/health_effects/en/.
- 3. Hedman L, Bjerg A, Perzanowski M, Sundberg S, Ronmark E. Factors related to tobacco use among teenagers. *Respir Med*. 2007;**101**(3):496–502. doi: 10.1016/j.rmed.2006.07.001. [PubMed: 16914300].
- Hossain S, Hossain S, Ahmed F, Islam R, Sikder T, Rahman A. Prevalence of tobacco smoking and factors associated with the initiation of smoking among university students in Dhaka, Bangladesh. *Cent Asian J Glob Health.* 2017;6(1):244. doi: 10.5195/cajgh.2017.244. [PubMed: 29138736]. [PubMed Central: PMC5675390].
- Adger H Jr, Saha S. Alcohol use disorders in adolescents. *Pediatr Rev.* 2013;34(3):103–13. quiz 114. doi: 10.1542/pir.34-3-103. [PubMed: 23457197]. [PubMed Central: PMC4530292].
- World Health Organization. Management of substance abuse: alcohol. 2017, [cited 11 November]. Available from: http://www.who.int/ substance_abuse/facts/alcohol/en/.
- Katsiki N, Papadopoulou SK, Fachantidou AI, Mikhailidis DP. Smoking and vascular risk: Are all forms of smoking harmful to all types of vascular disease? *Public Health.* 2013;127(5):435–41. doi: 10.1016/j.puhe.2012.12.021. [PubMed: 23453194].
- Athyros VG, Tziomalos K, Katsiki N, Gossios TD, Giouleme O, Anagnostis P, et al. The impact of smoking on cardiovascular outcomes and comorbidities in statin-treated patients with coronary artery disease: A post hoc analysis of the GREACE study. *Curr Vasc Pharmacol.* 2013;11(5):779–84. doi: 10.2174/1570161111311050016. [PubMed: 23140546].

- [No authors listed]. Preventing tobacco use among young people. A report of the Surgeon General. Executive summary. MMWR Recomm Rep. 1994;43(RR-4):1-10. [PubMed: 8183225].
- Chen L, Smith GD, Harbord RM, Lewis SJ. Alcohol intake and blood pressure: A systematic review implementing a Mendelian randomization approach. *PLoS Med.* 2008;5(3). e52. doi: 10.1371/journal.pmed.0050052. [PubMed: 18318597]. [PubMed Central: PMC2265305].
- Djousse L, Lee IM, Buring JE, Gaziano JM. Alcohol consumption and risk of cardiovascular disease and death in women: Potential mediating mechanisms. *Circulation*. 2009;**120**(3):237–44. doi: 10.1161/CIR-CULATIONAHA.108.832360. [PubMed: 19597054]. [PubMed Central: PMC2745640].
- Kourlaba G, Panagiotakos DB, Mihas K, Alevizos A, Marayiannis K, Mariolis A, et al. Dietary patterns in relation to socio-economic and lifestyle characteristics among Greek adolescents: A multivariate analysis. *Public Health Nutr.* 2009;**12**(9):1366–72. doi: 10.1017/S1368980008004060. [PubMed: 19063765].
- Rasmussen M, Krolner R, Klepp KI, Lytle L, Brug J, Bere E, et al. Determinants of fruit and vegetable consumption among children and adolescents: A review of the literature. Part I: Quantitative studies. *Int J Behav Nutr Phys Act.* 2006;3:22. doi: 10.1186/1479-5868-3-22. [PubMed: 16904006]. [PubMed Central: PMC1564033].
- Kapantais E, Chala E, Kaklamanou D, Lanaras L, Kaklamanou M, Tzotzas T. Breakfast skipping and its relation to BMI and healthcompromising behaviours among Greek adolescents. *Public Health Nutr.* 2011;14(1):101-8. doi: 10.1017/S1368980010000765. [PubMed: 20529399].
- Wong JE, Skidmore PM, Williams SM, Parnell WR. Healthy dietary habits score as an indicator of diet quality in New Zealand adolescents. *J Nutr.* 2014;**144**(6):937-42. doi: 10.3945/jn.113.188375. [PubMed: 24744308].
- Larson NI, Story M, Perry CL, Neumark-Sztainer D, Hannan PJ. Are diet and physical activity patterns related to cigarette smoking in adolescents? Findings from Project EAT. *Prev Chronic Dis.* 2007;4(3). A51. [PubMed: 17572955]. [PubMed Central: PMC1955390].
- Gisle L, Van Oyen H, Demarest S, Tafforeau J. Food consumption patterns among adolescent and young adult smokers. *Public Health*. 2009;**123**(9):632–4. doi: 10.1016/j.puhe.2009.07.016. [PubMed: 19747704].
- Centers for Disease Control and Prevention. 2015 National Youth Risk Behavior Survey. 2015, [cited 12 November]. Available from: ftp://ftp. cdc.gov/pub/data/yrbs/2015/2015_xxh_questionnaire.pdf.
- Sirirassamee T, Sirirassamee B. Health risk behavior among Thai youth: National survey 2013. Asia Pac J Public Health. 2015;27(1):76–84. doi: 10.1177/1010539514548759. [PubMed: 25183211].
- Jamal A, Gentzke A, Hu SS, Cullen KA, Apelberg BJ, Homa DM, et al. Tobacco use among middle and high school students - United States, 2011-2016. MMWR Morb Mortal Wkly Rep. 2017;66(23):597–603. doi: 10.15585/mmwr.mm6623a1. [PubMed: 28617771]. [PubMed Central: PMC5657845].
- Sirirassamee T, Sirirassamee B. Trends in tobacco use among Thai adolescents. J Med Assoc Thai. 2013;96 Suppl 1:S78-84. [PubMed: 23724460].
- Al-Sadat N, Misau AY, Zarihah Z, Maznah D, Tin Tin S. Adolescent tobacco use and health in Southeast Asia. Asia Pac J Public Health. 2010;22(3 Suppl):175S–80S. doi: 10.1177/1010539510372835. [PubMed: 20566551].
- Chamratrithirong A, Miller BA, Byrnes HF, Rhucharoenpornpanich O, Cupp PK, Rosati MJ, et al. Spirituality within the family and the prevention of health risk behavior among adolescents in Bangkok, Thailand. Soc Sci Med. 2010;71(10):1855–63. doi: 10.1016/j.socscimed.2010.08.020. [PubMed: 20926170]. [PubMed Central: PMC3001397].
- 24. National Statistical Office. *Social indicators* 2013. 2013, [cited 23 January]. Available from: http://web.nso.go.th/en/survey/env/data_env/ 580611_indSoc2013.pdf.

- Wakabayashi M, McKetin R, Banwell C, Yiengprugsawan V, Kelly M, Seubsman SA, et al. Alcohol consumption patterns in Thailand and their relationship with non-communicable disease. *BMC Public Health*. 2015;15:1297. doi: 10.1186/s12889-015-2662-9. [PubMed: 26704520]. [PubMed Central: PMC4690366].
- Hoffman JH, Welte JW, Barnes GM. Co-occurrence of alcohol and cigarette use among adolescents. *Addict Behav*. 2001;26(1):63–78. doi: 10.1016/S0306-4603(00)00089-7. [PubMed: 11196293].
- Liang W, Chikritzhs T. Alcohol consumption during adolescence and risk of diabetes in young adulthood. *Biomed Res Int.* 2014;2014:795741. doi: 10.1155/2014/795741. [PubMed: 24757678]. [PubMed Central: PMC3976876].
- Thompson RI, Pyke S, Scott EA, Thompson SG, Wood DA. Cigarette smoking, polyunsaturated fats, and coronary heart disease. *Ann* N Y Acad Sci. 1993;686:130–8. discussion 138-9. doi: 10.1111/j.1749-6632.1993.tb39167.x. [PubMed: 8512244].
- Zhao J, Kelly M, Bain C, Seubsman SA, Sleigh A, Thai Cohort Study T. Risk factors for cardiovascular disease mortality among 86866 members of the Thai Cohort Study, 2005-2010. *Glob J Health Sci.* 2015;7(1):107-14. doi: 10.5539/gjhs.v7n1p107. [PubMed: 25651607]. [PubMed Central: PMC4796435].
- Paulus D, Saint-Remy A, Jeanjean M. Dietary habits during adolescence-results of the Belgian Adolux Study. *Eur J Clin Nutr.* 2001;55(2):130–6. doi: 10.1038/sj.ejcn.1601132. [PubMed: 11305626].
- 31. Peltzer K, Pengpid S. Fruits and vegetables consumption and associated factors among in-school adolescents in five Southeast

Asian countries. *Int J Environ Res Public Health*. 2012;9(10):3575-87. doi: 10.3390/ijerph9103575. [PubMed: 23202763]. [PubMed Central: PMC3509472].

- Papadopoulou SK, Hassapidou MN, Katsiki N, Fachantidis P, Fachantidou AI, Daskalou E, et al. Relationships between alcohol consumption, smoking status and food habits in Greek adolescents. Vascular implications for the future. *Curr Vasc Pharmacol.* 2017;**15**(2):167–73. doi: 10.2174/1570161114666161024123357. [PubMed: 27781959].
- Lee B, Yi Y. Smoking, Physical Activity, and Eating Habits Among Adolescents. West J Nurs Res. 2016;38(1):27-42. doi: 10.1177/0193945914544335. [PubMed: 25082709].
- Padrao P, Lunet N, Santos AC, Barros H. Smoking, alcohol, and dietary choices: Evidence from the Portuguese National Health Survey. *BMC Public Health*. 2007;7:138. doi: 10.1186/1471-2458-7-138. [PubMed: 17608935]. [PubMed Central: PMC1939992].
- Andersen LF, Nes M, Sandstad B, Bjorneboe GE, Drevon CA. Dietary intake among Norwegian adolescents. *Eur J Clin Nutr.* 1995;49(8):555–64. [PubMed: 7588506].
- Cavallo DA, Smith AE, Schepis TS, Desai R, Potenza MN, Krishnan-Sarin S. Smoking expectancies, weight concerns, and dietary behaviors in adolescence. *Pediatrics*. 2010;**126**(1):e66–72. doi: 10.1542/peds.2009-2381. [PubMed: 20547649]. [PubMed Central: PMC3662026].
- Worsley A, Wang WC, Hunter W. The relationships between eating habits, smoking and alcohol consumption, and body mass index among baby boomers. *Appetite*. 2012;**58**(1):74–80. doi: 10.1016/j.appet.2011.09.003. [PubMed: 21986185].