



# The Efficacy of Nebulized Soluble Mannitol and Comparison with 5% Hypertonic Saline on Pulmonary Function of Children with Cystic Fibrosis

Saeed Sadr<sup>1</sup>, Masood Kiani<sup>2\*</sup>, Mohammad Rezaei<sup>3</sup>, Ghamartaj Khanbabaee<sup>1</sup>, Seyed Ahmad Tabatabaee<sup>1</sup> and Amirhossein Hosseini<sup>4</sup> 

<sup>1</sup>Department of Pediatric Pulmonology, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>2</sup>Non-Communicable Pediatric Diseases Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, IR Iran

<sup>3</sup>Shahid Rajaei Heart Center, Tehran University of Medical Sciences, Tehran, Iran

<sup>4</sup>Pediatric Gastroenterology, Hepatology and Nutrition Research Center, Research Institute for Children's Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran

\*Corresponding author: Assistant Professor of Pediatric Pulmonology, Non-Communicable Pediatric Diseases Research Center, No 19, Amirkola Children's Hospital, Amirkola, Babol, Mazandaran Province, 47317-41151, IR Iran. Tel/Fax: +98-1132346963, Email: kianimasood@yahoo.com

Received 2018 October 21; Revised 2019 May 15; Accepted 2019 May 20.

## Abstract

**Background:** One of the main aspects of the treatment in patients with cystic fibrosis (CF) is the effective clearing of airway secretions. Inhaled hypertonic saline (HTS) and mannitol are among drugs used for this purpose. There are two forms of mannitol, including dry-powder and soluble form. We used soluble mannitol in this study.

**Objectives:** The aim of this study was to evaluate the effect of inhaled soluble mannitol on the pulmonary function of patients with CF as well as its comparison with 5% HTS.

**Methods:** This clinical trial study was performed on 30 children with CF  $\geq$  5 years of age who referred to the CF clinic in Mofid Children's Hospital. Patients were divided into two groups, including soluble mannitol and 5% HTS group. At baseline and two weeks after the treatment, a spirometry test was done and forced expiratory flow in 1 second (FEV<sub>1</sub>), forced vital capacity (FVC), and Forced expiratory flow between 25% and 75% of forced vital capacity (FEF<sub>25-75%</sub>) were measured.

**Results:** The mean of FEV<sub>1</sub>, FVC, and FEF<sub>25-75%</sub> in the mannitol group before the treatment were  $96 \pm 30.5\%$ ,  $99 \pm 27\%$ , and  $76.3 \pm 36.3\%$ , and after the treatment were  $100.5 \pm 29\%$ ,  $103.5 \pm 23.7\%$ , and  $79.9 \pm 37.3\%$ , respectively. The difference of FEV<sub>1</sub> before and after mannitol was statistically significant ( $P = 0.031$ ). The mean of FEV<sub>1</sub>, FVC, and FEF<sub>25-75%</sub> in 5% HTS group were  $96.2 \pm 25.5\%$ ,  $99 \pm 19.8\%$  and  $77.8 \pm 35.6\%$  before the treatment and  $95.2 \pm 20.1\%$ ,  $99.3 \pm 19.2\%$ ,  $74.2 \pm 29.5\%$  after the treatment. There was no significant difference in 5% HTS group.

**Conclusions:** Inhaled soluble mannitol improved pulmonary function of patients with CF and its effect was better than 5% HTS.

**Keywords:** Mannitol, Hypertonic Saline, Cystic Fibrosis, Children, Pulmonary Function

## 1. Background

Cystic fibrosis (CF) is the most common inherited life-limiting disease of Caucasians. Most of the CF patients' morbidity and mortality are caused by the deterioration of pulmonary function (1). The pathogenesis of CF results from reduced hydration of the airway surface, which impairs the clearance of the mucociliary system, and finally, leads to chronic airway obstruction and infection. Thus, the patients with CF mostly experience intermittent pulmonary exacerbations and hospital admissions (2-4).

Numerous treatments are available in order to decrease the inflammatory process and thus the complications, exacerbation periods, and hospitalizations. These

therapies include antibiotics, anti-inflammatory drugs, and mucolytics and maneuvers that enhance the airway clearance (4-6). A number of mucolytics have been used in patients with CF. These include inhaled rhDNase, N-acetylcysteine, hypertonic saline, and dry powder mannitol (1, 4).

Hypertonic saline (HTS) and mannitol are hyperosmolar agents that increase airway surface liquid (ASL) and improve the underlying hydration defects in patients with CF. The osmotic gradient by hyperosmolar agents induce transepithelial rehydration and optimize the mucosal viscoelasticity necessary for intact mucociliary clearance function (7-10). Clinical trials have confirmed the efficacy and safety of hypertonic saline as a treatment to improve mucociliary clearance and lung function in patients

with CF by improving ASL hydration, but this fact has been shown as a short-term effect (11).

Mannitol is sugar-based alcohol that creates an osmotic gradient and moves the water into the lumen and improves the mucociliary clearance (4, 12). It is thought that dry powder mannitol inhalation increases the volume of airway surface liquid and so it aims for mucosal clearance (13, 14). It has been confirmed that inhaled mannitol had advantages for patients with bronchiectasis and CF (13, 15, 16). A triple-phase study had been done by Bilton et al. in which the efficacy and safety of mannitol were proven and also demonstrated that mannitol had clinically significant effects on pulmonary function in patients with CF (13).

There are two forms of mannitol, including dry powder and soluble form. There are several studies conducted on the use of dry powder mannitol in patients with CF. Also, *in vitro* studies showed that soluble mannitol can be delivered to the lung with respirable particle size using mesh nebulizer device (10). Owing to the lack of mannitol in dry powder in Iran, our study was conducted using soluble mannitol.

## 2. Objectives

In this study, the effect of nebulized soluble mannitol on the pulmonary function of patients with CF, as well as its comparison with 5% hypertonic saline, were investigated.

## 3. Methods

Our research is a clinical trial study on 30 patients with CF older than 5 years because of the ability to perform spirometry who referred to the CF clinic of Mofid Children's Hospital as a referral medical center in Tehran, Iran from January 2016 to June 2016. The trial was approved by the Ethics Committee of Shahid Beheshti University of Medical Sciences (IR.SBMU.SM.REC.1394.62) and was registered in the Iranian Registry of Clinical Trials (IRCT20180307038994N1). The CF was diagnosed in these patients based on clinical manifestations and abnormal results of a sweat chloride test. The patients were entered to study by random sampling and participation in the study was voluntary. After complete explanations and awareness of our research aims and goals, written informed consent was obtained from the parents before each case entered the study.

A basic spirometry was performed for each patient and spirometric results, including forced expiratory flow in 1 second (FEV1), forced vital capacity (FVC), and Forced expiratory flow between 25% and 75% of forced vital capacity

(FEF25 - 75%), were recorded. The patients were divided into two groups by random sampling method. The first group underwent the treatment with soluble mannitol 4 cc twice a day by nebulizer device. We used soluble mannitol because of unavailability of dry-powder mannitol in Iran. Soluble mannitol was prepared in 4 cc vials from a combination of 20% mannitol (3 cc) and 0.9% saline (1 cc). This composition contains 150 mg/mL mannitol. The other group underwent the treatment with hypertonic saline (5%) 5 cc four times a day by nebulizer device. Duration of the treatment was two weeks in both groups. The patients were treated at home with once daily 5% HTS and chest physiotherapy. The patients were stable during the study and did not develop pulmonary exacerbation. Other previous medications used by patients have continued throughout the study period. In this study, medications were nebulized by the "eFlow Rapid Mesh Nebulizer" made by PARI company of Germany. According to the possibility of bronchospasm occurrence after nebulization of mannitol or hypertonic saline, we used a bronchodilator before each time of nebulization. Cough, pharyngitis, bronchospasm, and wheezing are the possible complications of mannitol. During the treatment time, the patients were completely monitored and observed from the point of view of drug complications and adverse effects via follow-up in the clinic or via telephone contact. The second spirometry was done after 2 weeks of treatment.

All the patients' data were analyzed by SPSS statistics software V. 21.0 and reviewed by an expert statistics specialist unaware of the patient group and medications.

## 4. Results

Thirty patients with CF, including 17 (57%) males and 13 (43%) females, participated in this study. The demographic characteristics of the patients are presented in Table 1. The mean of FEV1, FVC, and FEF25 - 75% before and after the treatment in two groups are presented in Table 2.

In the mannitol group, the mean of FEV1 at baseline was  $96 \pm 30.5\%$  and after the treatment was  $100.5 \pm 29\%$ . The difference of FEV1 before and after the mannitol treatment was 4.5% ( $P = 0.031$ ), which was statistically significant. In

**Table 1.** Demographic Characteristics of the Patients in the Mannitol and 5% HTS Groups<sup>a</sup>

Variable	Mannitol	5% HTS
<b>Gender</b>		
Male	9 (60)	8 (53)
Female	6 (40)	7 (47)
<b>Age</b>	$10.4 \pm 4.3$	$11 \pm 4.9$

Abbreviation: HTS, hypertonic saline.

<sup>a</sup>Values are expressed as mean  $\pm$  SD or No. (%).

**Table 2.** Comparison of FEV1, FVC, and FEF25 - 75% Before and After the Treatment in the Two Groups of Mannitol and 5% HTS<sup>a</sup>

Drug, PFT Parameters	Before Treatment	After Treatment	P Value
<b>Mannitol</b>			
FEV1	96 ± 30.5	100.5 ± 29	0.031
FVC	99 ± 27	103.5 ± 23.7	0.066
FEF25 - 75%	76.3 ± 36.3	79.9 ± 37.3	0.426
<b>5% HTS</b>			
FEV1	96.2 ± 25.5	95.2 ± 20.1	0.699
FVC	99 ± 19.8	99.3 ± 19.2	0.929
FEF25 - 75%	77.8 ± 35.6	74.2 ± 29.5	0.352

Abbreviations: FEF25 - 75%, forced expiratory flow between 25% and 75% of forced vital capacity; FEV1, forced expiratory volume in 1 second; FVC, forced vital capacity; HTS, hypertonic saline.

<sup>a</sup>Values are expressed as mean ± SD.

the 5% HTS group, the mean of FEV1 at baseline was  $96.2 \pm 25.5\%$  and after treatment was  $95.2 \pm 20.1\%$ . No significant statistical difference was seen after HTS nebulization.

The mean increase in FVC from the baseline after the treatment with mannitol was 4.5%, which was not statistically significant ( $P = 0.066$ ). At baseline, the mean of FVC in the HTS group was  $99 \pm 19.8\%$  and after the treatment was  $99.3 \pm 19.2\%$  in which no significant difference was observed ( $P = 0.929$ ).

In the mannitol group, the mean of FEF25 - 75% at baseline was  $76.3 \pm 36.3\%$  and after the treatment was  $79.9 \pm 37.3\%$ , which had no statistically significant difference ( $P = 0.426$ ). In the 5% HTS group, the mean of FEF25 - 75% at the beginning of the study was  $77.8 \pm 35.6\%$  and at the end of the study was  $74.2 \pm 29.5\%$ , which showed no statistically significant difference (0.352).

## 5. Discussion

Our study showed that inhaling soluble mannitol with mesh nebulizer device could improve pulmonary function. The improvement in pulmonary function following the use of inhaled mannitol is probably due to an increase in mucociliary clearance in central and small airways. In our study, pulmonary function, which was determined based on FEV1, FVC, and FEF25 - 75% measurements before and after receiving mannitol, showed considerable improvement.

The increase in FEV1 after two weeks of mannitol treatment was 4.5%, which was statistically significant ( $P = 0.031$ ). The FVC and FEF25 - 75% also increased in the mannitol group (FVC: 4.5% and FEF25 - 75%: 3.6%), but despite increasing from baseline, was not statistically significant ( $P = 0.66$  and  $P = 0.42$ , respectively).

Several studies on pulmonary function of the patients with CF indicated the effect of mannitol in the form of

dry-powder, as an osmotic agent, led to hydration of airway surface liquid (ASL) and increased mucociliary clearance. In these studies, the improvement in the pulmonary function was demonstrated based on pulmonary function test values. For example, in the Aitken et al. study, which was performed on 318 patients with CF, the increase in FEV1 was 8.2% from baseline in the mannitol group and this study showed that inhaled mannitol improved lung function and this improvement maintained for 26 weeks after the treatment (4). In another study by Bilton et al. on 324 patients with CF, the increase in FEV1 in the mannitol group was 6.5% and in this study, the safety of inhaled mannitol and an improvement in lung function were shown (13). In Jaques et al. study, which was performed on 39 patients with CF, inhaled mannitol treatment over a period of 2 weeks improved lung function (15).

In a Cochrane review by Nolan et al. in 2015, an improvement in pulmonary function and a decrease in pulmonary exacerbation in patients receiving inhaled mannitol were demonstrated (17). In our study, pulmonary function (FEV1, FVC, and FEF25 - 75%) after the treatment indicated no significant difference from baseline in 5% HTS group. There are studies that represent an improvement in pulmonary function using 7% hypertonic saline (11, 18), while in our study 5% HTS was used. According to some studies, with increasing saline concentration, the mucociliary clearance increases (19) and this may, to some extent, be the reason for the inconsistency of our study with studies using 7% HTS; however, a small study in India showed that 3% HTS is as effective as 7% HTS (20). The main difference of our study with other studies using inhaled mannitol is that in other studies, mannitol was used in the form of dry-powder, while in our study mannitol was used in the form of a solution with nebulizer device.

The present study was designed, based on the study of Chan et al., which was performed *in vitro* in 2011, and showed that soluble mannitol in combination with different saline concentrations using an eFlow rapid mesh nebulizer could be distributed at a particle size of 5 - 6 micrometer (10), and particles with this size could reach the lungs, and was able to show some improvement in pulmonary function. Mannitol complications consisted of cough, hemoptysis, and bronchospasm. In this study, the most common complication in patients receiving mannitol was cough; however, the induced cough could also be one of the therapeutic effects of mannitol and led to an increase in mucociliary clearance.

Hemoptysis was detected in no patient, and bronchospasm was prevented by giving the bronchodilator before using mannitol. Adherence to the treatment is one of the important aspects of treating patients with CF. In the current study, adherence in the mannitol group was bet-

ter than 5% HTS group, and this can be due to the use of mannitol twice a day instead of 5% HTS four times a day.

### 5.1. Conclusions

Our study demonstrated that inhaling soluble mannitol using the eFlow rapid mesh nebulizer could improve the pulmonary function of patients with CF, and its effects on pulmonary function of these patients were better than 5% HTS. This study also revealed that inhaling the soluble mannitol is safe and tolerable for the patients. Meanwhile, the use of soluble mannitol with nebulizer allows the drug to be used in younger children who cannot use the mannitol in dry-powder form. However, more extensive studies are needed to recommend the use of soluble mannitol as well as its safety and efficacy.

### Acknowledgments

The authors would like to acknowledge the Clinical Research Development Committee of Amirkola Children's Hospital and generous contributions of Pediatric Gastroenterology, Hepatology and Nutrition Research Center, Research Institute for Children's Health.

### Footnotes

**Authors' Contribution:** Patient gathering: Saeed Sadr, Masood Kiani, Seyed Ahmad Tabatabaee, and Gamartaj Khanbabae; supervisor: Saeed Sadr and Masood Kiani; data collection and analysis: Mohamad Rezaei; editing manuscript: Amirhossein Hosseini.

**Clinical Trial Registration Code:** This trial was registered in the Iranian Registry of Clinical Trials (IRCT20180307038994N1).

**Conflict of Interests:** There is no conflict of interest.

**Ethical Approval:** This study was approved by the Ethics Committee of Shahid Beheshti University of Medical Sciences (IR.SBMU.SM.REC.1394.62).

**Funding/Support:** There is no funding/support for this study.

### References

1. Verkman AS, Song Y, Thiagarajah JR. Role of airway surface liquid and submucosal glands in cystic fibrosis lung disease. *Am J Physiol Cell Physiol*. 2003;**284**(1):C2-15. doi: [10.1152/ajpcell.00417.2002](https://doi.org/10.1152/ajpcell.00417.2002). [PubMed: [12475759](https://pubmed.ncbi.nlm.nih.gov/12475759/)].
2. Puchelle E, Bajolet O, Abely M. Airway mucus in cystic fibrosis. *Paediatr Respir Rev*. 2002;**3**(2):115-9. [PubMed: [12297057](https://pubmed.ncbi.nlm.nih.gov/12297057/)].
3. Donaldson SH, Boucher RC. Sodium channels and cystic fibrosis. *Chest*. 2007;**132**(5):1631-6. doi: [10.1378/chest.07-0288](https://doi.org/10.1378/chest.07-0288). [PubMed: [17998363](https://pubmed.ncbi.nlm.nih.gov/17998363/)].
4. Aitken ML, Bellon G, De Boeck K, Flume PA, Fox HG, Geller DE, et al. Long-term inhaled dry powder mannitol in cystic fibrosis: an international randomized study. *Am J Respir Crit Care Med*. 2012;**185**(6):645-52. doi: [10.1164/rccm.201109-1666OC](https://doi.org/10.1164/rccm.201109-1666OC). [PubMed: [22198974](https://pubmed.ncbi.nlm.nih.gov/22198974/)].
5. Ratjen FA. Cystic fibrosis: pathogenesis and future treatment strategies. *Respir Care*. 2009;**54**(5):595-605. [PubMed: [19393104](https://pubmed.ncbi.nlm.nih.gov/19393104/)].
6. Davies JC, Alton EW, Bush A. Cystic fibrosis. *BMJ*. 2007;**335**(7632):1255-9. doi: [10.1136/bmj.39391.713229.AD](https://doi.org/10.1136/bmj.39391.713229.AD). [PubMed: [18079549](https://pubmed.ncbi.nlm.nih.gov/18079549/)]. [PubMed Central: [PMC2137053](https://pubmed.ncbi.nlm.nih.gov/PMC2137053/)].
7. Goralski JL, Boucher RC, Button B. Osmolytes and ion transport modulators: New strategies for airway surface rehydration. *Curr Opin Pharmacol*. 2010;**10**(3):294-9. doi: [10.1016/j.coph.2010.04.003](https://doi.org/10.1016/j.coph.2010.04.003). [PubMed: [20439165](https://pubmed.ncbi.nlm.nih.gov/20439165/)].
8. Daviskas E, Anderson SD, Eberl S, Young IH. Beneficial effect of inhaled mannitol and cough in asthmatics with mucociliary dysfunction. *Respir Med*. 2010;**104**(11):1645-53. doi: [10.1016/j.rmed.2010.05.019](https://doi.org/10.1016/j.rmed.2010.05.019). [PubMed: [20576419](https://pubmed.ncbi.nlm.nih.gov/20576419/)].
9. Daviskas E, Anderson SD, Jaques A, Charlton B. Inhaled mannitol improves the hydration and surface properties of sputum in patients with cystic fibrosis. *Chest*. 2010;**137**(4):861-8. doi: [10.1378/chest.09-2017](https://doi.org/10.1378/chest.09-2017). [PubMed: [19880909](https://pubmed.ncbi.nlm.nih.gov/19880909/)].
10. Chan JG, Kwok PC, Young PM, Chan HK, Traini D. Mannitol delivery by vibrating mesh nebulisation for enhancing mucociliary clearance. *J Pharm Sci*. 2011;**100**(7):2693-702. doi: [10.1002/jps.22494](https://doi.org/10.1002/jps.22494). [PubMed: [21283985](https://pubmed.ncbi.nlm.nih.gov/21283985/)].
11. Elkins MR, Robinson M, Rose BR, Harbour C, Moriarty CP, Marks GB, et al. A controlled trial of long-term inhaled hypertonic saline in patients with cystic fibrosis. *N Engl J Med*. 2006;**354**(3):229-40. doi: [10.1056/NEJMoa043900](https://doi.org/10.1056/NEJMoa043900). [PubMed: [16421364](https://pubmed.ncbi.nlm.nih.gov/16421364/)].
12. Daviskas E, Anderson SD, Brannan JD, Chan HK, Eberl S, Bautovich G. Inhalation of dry-powder mannitol increases mucociliary clearance. *Eur Respir J*. 1997;**10**(11):2449-54. [PubMed: [9426077](https://pubmed.ncbi.nlm.nih.gov/9426077/)].
13. Bilton D, Robinson P, Cooper P, Gallagher CG, Kolbe J, Fox H, et al. Inhaled dry powder mannitol in cystic fibrosis: An efficacy and safety study. *Eur Respir J*. 2011;**38**(5):1071-80. doi: [10.1183/09031936.00187510](https://doi.org/10.1183/09031936.00187510). [PubMed: [21478216](https://pubmed.ncbi.nlm.nih.gov/21478216/)].
14. Brannan JD, Anderson SD, Perry CP, Freed-Martens R, Lassig AR, Charlton B, et al. The safety and efficacy of inhaled dry powder mannitol as a bronchial provocation test for airway hyperresponsiveness: A phase 3 comparison study with hypertonic (4.5%) saline. *Respir Res*. 2005;**6**:144. doi: [10.1186/1465-9921-6-144](https://doi.org/10.1186/1465-9921-6-144). [PubMed: [16336673](https://pubmed.ncbi.nlm.nih.gov/16336673/)]. [PubMed Central: [PMC1326200](https://pubmed.ncbi.nlm.nih.gov/PMC1326200/)].
15. Jaques A, Daviskas E, Turton JA, McKay K, Cooper P, Stirling RG, et al. Inhaled mannitol improves lung function in cystic fibrosis. *Chest*. 2008;**133**(6):1388-96. doi: [10.1378/chest.07-2294](https://doi.org/10.1378/chest.07-2294). [PubMed: [18339790](https://pubmed.ncbi.nlm.nih.gov/18339790/)].
16. Teper A, Jaques A, Charlton B. Inhaled mannitol in patients with cystic fibrosis: A randomised open-label dose response trial. *J Cyst Fibros*. 2011;**10**(1):1-8. doi: [10.1016/j.jcf.2010.08.020](https://doi.org/10.1016/j.jcf.2010.08.020). [PubMed: [20888307](https://pubmed.ncbi.nlm.nih.gov/20888307/)].
17. Nolan SJ, Thornton J, Murray CS, Dwyer T. Inhaled mannitol for cystic fibrosis. *Cochrane Database Syst Rev*. 2015;(10). CD008649. doi: [10.1002/14651858.CD008649.pub2](https://doi.org/10.1002/14651858.CD008649.pub2). [PubMed: [26451533](https://pubmed.ncbi.nlm.nih.gov/26451533/)].
18. Donaldson SH, Bennett WD, Zeman KL, Knowles MR, Tarran R, Boucher RC. Mucus clearance and lung function in cystic fibrosis with hypertonic saline. *N Engl J Med*. 2006;**354**(3):241-50. doi: [10.1056/NEJMoa043891](https://doi.org/10.1056/NEJMoa043891). [PubMed: [16421365](https://pubmed.ncbi.nlm.nih.gov/16421365/)].
19. Robinson M, Hemming AL, Regnis JA, Wong AG, Bailey DL, Bautovich GJ, et al. Effect of increasing doses of hypertonic saline on mucociliary clearance in patients with cystic fibrosis. *Thorax*. 1997;**52**(10):900-3. doi: [10.1136/thx.52.10.900](https://doi.org/10.1136/thx.52.10.900). [PubMed: [9404379](https://pubmed.ncbi.nlm.nih.gov/9404379/)]. [PubMed Central: [PMC1758438](https://pubmed.ncbi.nlm.nih.gov/PMC1758438/)].
20. Gupta S, Ahmed F, Lodha R, Gupta YK, Kabra SK. Comparison of effects of 3 and 7% hypertonic saline nebulization on lung function in children with cystic fibrosis: A double-blind randomized, controlled trial. *J Trop Pediatr*. 2012;**58**(5):375-81. doi: [10.1093/tropej/fms004](https://doi.org/10.1093/tropej/fms004). [PubMed: [22374985](https://pubmed.ncbi.nlm.nih.gov/22374985/)].