

## Review Article:

# Mortality and Morbidity Patterns of Very Low Birth Weight Newborns in Eastern Mediterranean Region: A Meta-Analysis Study



Fatemeh Nayeri<sup>1</sup>, Zahra Emami<sup>1</sup>, Yasamin Mohammadzadeh<sup>2</sup>, Mamak Shariat<sup>1</sup>, Setareh Sagheb<sup>2</sup>, Leyla Sahebi<sup>1\*</sup>

1. Maternal, Fetal, and Neonatal Research Center, Tehran University of Medical Sciences, Tehran, Iran.

2. Breast Feeding Research Center, Tehran University of Medical Sciences, Tehran, Iran.



**Citation:** Nayeri F, Emami Z, Mohammadzadeh Y, Shariat M, Sagheb S, Sahebi L. Mortality and Morbidity Patterns of Very Low Birth Weight Newborns in Eastern Mediterranean Region: A Meta-Analysis Study. J Pediatr Rev. 2019; 7(2):67-76. <http://dx.doi.org/10.32598/jpr.7.2.67>

**doi:** <http://dx.doi.org/10.32598/jpr.7.2.67>



**Funding:** See Page 74

#### Article info:

**Received:** 21 February 2018

**First Revision:** 15 March 2018

**Accepted:** 10 June 2018

**Published:** 01 April 2019

#### Key Words:

Very low birth weight infant, Premature birth, Mortality, Morbidity, Mediterranean region, Meta-analysis

## ABSTRACT

**Context:** Preterm and low-birth weight are of the most common causes of mortality and critical disabilities in newborns. Although advances in neonatal care improved survival and quality of life in the developed countries, mortality and morbidity due to prematurity are still high in the developing and underdeveloped countries such countries in Eastern Mediterranean Region (EMR).

**Objective:** The study aimed at evaluating mortality and morbidity among Very Low Birth Weight (VLBW) and Very Preterm (VP) neonates in EMR.

**Data Sources:** A systematic search was conducted in CINAHL, Cochrane Library, Current Contents, Database of Abstracts of Reviews of Effects (DARE), Google Scholar, PubMed, PsycINFO, Thomson Reuters, Scientific Information Database (SID), and Medical Library (MedLib) as well as regional WHO database to detect relevant studies published from 1996 to 2016.

**Study Selection:** English Language Literature From EMR countries, which reporting outcomes of VLBW and VP were considered. Finally, 32 full text articles were included.

**Data Extraction:** Data on VLBW and VP outcomes including mortality, and short- and long-term morbidities were extracted.

**Results:** Meta-analysis results of prevalence of mortality among VLBW newborns was obtained as 32.0%, which was different from Asian and African countries (23% vs. 55%). Pooled odds ratio of mortality among VLBW newborns was estimated as 2.41. The most prevalent comorbidity in VLBW and VP newborns were allocated to septicemia and Neurodevelopmental Delay (NDD), and the rarest belonged to Periventricular Leukomalacia (PVL) and Necrotizing Enterocolitis (NEC). The main limitation of the current study was lack of relevant studies and adequate sample size.

**Conclusions:** It seems that indicators in EMR should be evaluated independently in Asian and African countries. Septicemia, as relatively curable cause of morbidity, is the most prevalent comorbidity among VLBW infants.

\* Corresponding Author:

**Leyla Sahebi, PhD.**

**Address:** Maternal, Fetal and Neonatal Research Center, Tehran University of Medical Sciences, Tehran, Iran.

**Tel:** +98 (21) 61192631

**E-mail:** sahebileila@yahoo.com

## 1. Context

**N**eonatal mortality is one of the major global health concerns to such an extent that 45% of total mortalities in children less than five years occur in neonatal period (1). Very Preterm (VP) and Very Low Birth Weight (VLBW) are defined as gestational age <32 weeks and birth weight <1500 g, respectively (1). Preterm birth and Low-Birth Weight (LBW) are the most common causes of newborn mortality. In addition, survived preterm infants are compromised to lifelong disabilities, such as major neurodevelopmental disorders, intellectual disability, hydrocephaly, as well as visual and hearing impairments (2-7).

Most of the VP and VLBW infants need advanced neonatal cares to survive. Although advances in neonatal care have improved their survival and quality of life in the developed countries, VP and VLBW morbidity and mortality are still at their peak in the developing and underdeveloped countries (8). In high-income countries, 95% of preterm babies survive, of which 90% develop no impairments, but in many low-income countries, almost all of the VP births die from lack of basic cares (9). On the other hand, most of the middle-income countries suffer from long-term consequences and disabilities caused by preterm births, especially extremely preterm births (10).

Overall, limited studies have been conducted on the outcomes of VLBW in infants. In addition, they also assessed a few types of morbidity. According to the World Health Organization (WHO), Eastern Mediterranean Region (EMR) mostly includes the middle- and low-income countries, and in spite of the importance of the study subject, no review study was conducted in this region. Therefore, there is a paucity of comprehensive studies on the outcomes of VP and VLBW in EMR countries.

## 2. Objective

The current study aimed at reviewing the morbidity and mortality of prematurity among VP and VLBW infants in EMR countries. Also, the study compared morbidity and mortality rate among infants born weighing <1500 to  $\geq$ 1500 g. The authors believe that the current meta-analysis can help have an insight of the prematurity problem in this region and clarify life-threatening outcomes under similar situations.

## 2.1. Inclusion criteria

All the articles published in English on outcomes of VP and VLBW (morbidity and mortality) in EMR countries from January 1996 to December 2016 were entered into the study.

## 3. Data Sources

The current study used web-based citation index to cite manuscripts of these identified articles. All the articles published from January 1996 to December 2016 as well as their reference lists were searched in order to extract relevant studies. The authors made a direct contact with some other authors from various EMR countries. Searching was conducted from September to December 2016.

### 3.1. Search strategies

The authors searched CINAHL, Cochrane Library, Current Contents, Database of Abstracts of Reviews of Effects (DARE), Google Scholar, PubMed, PsycINFO, Thomson Reuters, Scientific Information Database (SID), Medical Library (MedLib), and regional WHO database to detect relevant studies.

Medical Subject Heading (MeSH) and Cumulative Index to Nursing and Allied Health Literature (CINAHL) database were also used for terms and keywords. The following MeSH terms were used in the current study as keywords: (Preterm OR very preterm OR premature OR very premature OR low-birth-weight OR very-low-birth-weight immature) AND (infants OR newborns OR neonates) AND (mortality OR survival OR morbidity OR outcome OR follow-up) in combination with EMR countries names, including Afghanistan, Bahrain, Djibouti, Egypt, Iran (Islamic Republic of), Iraq, Jordan, Kuwait, Lebanon, Libya, Morocco, Oman, Pakistan, Palestine, Qatar, Saudi Arabia, Somalia, Sudan, Syrian Arab Republic, Tunisia, United Arab Emirates, Yemen.

## 4. Study Selection

To find articles reporting mortality and or morbidity of VP and or VLBW in EMR countries, a literature search was performed on observational and analytical studies, published in peer-reviewed journals from 1996 to 2016. The search strategy was developed and studied by all authors.

Publications would be included in the study if 1. They were published in English language from January 1996 to December 2016; 2. Reported outcomes of VP/VLBW

including mortality, short- and long-term morbidities; and 3. Had reports statistics from EMR countries. Studies would be excluded if they contained no data on above-mentioned outcomes, not reported data from EMR countries, systematic reviews with no direct data on required information, and case studies with sample sizes less than 15 subjects. In addition, publications reporting overall neonatal mortality or morbidity rate, without subcategories of gestational age or birth weight were excluded. Eligible studies were reviewed to recheck relevancy and accordingly, related data were excluded.

## 5. Data Extraction

Two reviewers extracted data based on data extraction template of the Cochrane consumers and communication review group. The pilot study was conducted on eight randomly-selected articles. One review author extracted the related data from included studies and the second author checked the accuracy of extracted data. Extracted data elements comprised study year, study design, geographical location, sample size, participants, inclusion criteria, predicted outcomes, type of statistics, and analysis method. Disagreements were resolved by a third or fourth author. Studies with duplicate citations and those conducted on certain subpopulations were excluded.

### 5.1. Data items

The primary outcome was prevalence and odds ratio of mortality among VLBW/VP newborns. Secondary outcomes included prevalence and odds ratio of morbidity from Respiratory Distress Syndrome (RDS), Bronchopulmonary Dysplasia (BPD), pneumothorax, pulmonary hemorrhage, Retinopathy of Prematurity (ROP), Intraventricular Hemorrhage (IVH), Periventricular Leukomalacia (PVL), Necrotizing Enterocolitis (NEC), Patent ductus Arteriosus (PDA), asphyxia, congenital anomalies, sepsis and Neurodevelopmental Delay (NDD) or Psychomotor Development Retardation (PDR), speech delay, Attention-Deficit/Hyperactivity Disorder (ADHD), Obsessive-Compulsive Disorder (OCD), and possibly seizures.

### 5.2. Risk of bias in case studies

The reviewers evaluated the risk of bias in the current review study. Risk of bias was checked by the Cochrane collaboration tool.

### 5.3. Summary measures

Pooled ratio of mortality and morbidity from various disorders and receiving surfactant, mechanical venti-

lation, or CPAP (Continuous Positive Airway Pressure) among VLBW and VP newborns were calculated by random-effects models; pooled odds ratio also was calculated by meta-regression analysis. All analyses were conducted with STATA 14 (Stata Corp. 2014. Stata Statistical Software: Release 14. College Station, TX: Stata Corp LP).

## 5.4. Study analyses

Heterogeneity and meta-regression were investigated by subgroups of the study year (before 2006 vs. after 2007), type of design (cross sectional study vs. other designs), sample size (<200 and >201), and countries (Asian vs. African countries).

## 5.5. Risk of bias across studies

The bias findings for prevalence and odds ratio of mortality in VLBW neonates are shown in funnel plots (Figures 1, 2).

## 6. Results

Initial search identified 1339 titles. After removing duplications, the remaining 1307 titles and abstracts were screened. Overall, the full-text of 117 screened studies were selected and reviewed. Finally, 32 full-texts were reviewed and analyzed to extract data (Figure 3). Overall, 9 study had reported mortality among VLBW newborns (11-19) (Figure 1).

The basic characteristics of studies reviewing mortality and morbidities among VLBW/VP newborns are shown in Tables 1 and 2. Considering the prevalence of mortality among VLBW newborns, the highest and lowest levels in the Middle-East were 60.0% in Egypt (12) and 14.0% in EUA (16). Unfortunately, information about prevalence of VP infants was not accessible in the studies. Meta-analysis of prevalence of mortality among VLBW newborns were 32.0% (95%CI: 27.0-38.0;  $P_{\text{heterogeneity}}=0.00$ ) (Figure 4).

In the meta-regression analysis, the prevalence of mortality among VLBW newborns was not heterogeneous based on the study design (meta-analysis of prevalence of mortality in cross-sectional studies=40.0% [95%CI:28.0-51.0] vs. other designs=26.0% [95%CI: 17.0-35.0],  $P_{\text{heterogeneity}}=0.074$ ) and year of study (before 2006=46% [95%CI: 20-72] vs. after 2007=26% [95%CI: 19-33],  $P_{\text{heterogeneity}}=0.137$ ).




Figure 1. Publication bias for the prevalence of mortality among VLBW neonates

Journal of Pediatrics Review

Pooled prevalence of mortality among VLBW newborns was heterogeneous based on Asian or African countries (Asian countries=23% [95%CI:19-27] vs. African countries=55% [95%CI:48.0-27.0];  $I^2=92.07\%$ ,  $P=0.00$ ).

Pooled prevalence of VLBW newborns mortality in studies with sample sizes smaller than 200 subjects was lower than those of the studies with sample sizes larger than 201 subjects (26%; 95%CI:15.0-37.0 vs. 36.0%; 95%CI:27.0-45.0) ( $I^2=92.07\%$ ,  $P=0.00$ ). In logistic regression analysis, pooled odds ratio of mortality among VLBW newborns was 2.41 (95%CI: 1.887-3.069) ( $P_{\text{heterogeneity}}=0.15$ ).

The pooled mean Apgar scores (95%CI) at the first and fifth minutes were 4.67 (95%CI:1.91-7.42,  $P_{\text{heterogeneity}}=0.815$ ) and 7.40 (95%CI: 5.38-9.42,  $P_{\text{heterogeneity}}=0.987$ ), respectively. The meta-analysis of prevalence of VLBW newborns receiving surfactant was 62% (95%CI:52-72), mechanical ventilation=56% (95%CI:51.0-61.0), and CPAP=41.0% (95%CI:36-46) (16-18, 20). Also, the meta-analysis of prevalence of VP

newborns receiving surfactant was 32% (95%CI:29.0-36.0) and CPAP=45.0% (95%CI:41.0-49.0) (21, 22).

In the current study, prevalence of various comorbidities with VLBW or VP in neonates was also analyzed. Table 3 summarizes the results (15, 16, 18, 20-26). The most prevalent comorbidity in VLBW newborns allocated to sepsis (49.0%; 95%CI: 38.0-59.0) (15) and the least belonged to PVL (3.0%; 95%CI:2.0-5.0) (16, 18, 20). In VP group, the most prevalent and the least comorbidities were NDD (73.0%, 95%CI:57.0-0.85) and NEC (0.0), respectively (21, 22).

Summary of odds ratio of comorbidity in VLBW neonates are presented in Table 4 (10, 27-32). The univariate and multiple-regression analysis were not possible in VP newborns because of the studies limitations.

### 7. Discussion

The average global rate of neonatal mortality was 32 deaths per 1000 live births in 2015 (33) and about




Figure 2. Publication bias for the odds ratio of mortality among VLBW neonates

Journal of Pediatrics Review




Figure 3. Flowchart of the included studies

Journal of Pediatrics Review

two-thirds of neonatal deaths occur in the African and Southeast Asian regions, according to WHO (34).

The risk of death in LBW infants is 40 times greater than normal weight babies. In addition, short gestation and preterm birth are of the main reasons of LBW (35).

The global rate of preterm birth is about 11% (from 5% in some European regions to 18% in some African regions) (35), which is responsible for 22% of neonatal mortality rate in EMR (34). Prevalence of LBW in EMR countries is high (19.5%), and accounts for 60% to 80% of all neonatal deaths either directly or indirectly (34).

Table 1. Characteristics of selected studies conducted on mortality rate of VLBW/VP newborns in Eastern Mediation region

Author(s) (Published Year)	Location	Sample Size	Type of Study
Chedid, et al. (2008) (16)	United Arab Emirates	173	Retrospective cohort
Al Hazzani, et al. (2011) (18)	Saudi Arabia	186	Retrospective cohort
Hussein, et al. (2014) (17)	Iraq	150	Retrospective cohort
Dawodu, et al. (2005) (14)	United Arab Emirates	40408	Cross-sectional
Malek-mellouli, et al. (2013) (15)	Tunis	82	Retrospective cohort
Ziadeh, et al. (2000) (11)	Jordan	386	Cross-sectional
Fakhr, et al. (2005) (12)	Egypt	480	Cross-sectional
Al Azzawi (2011) (19)	Iraq	924	Cross-sectional
Obaid, et al. (2010) (13)	Iraq	600	Retrospective cohort

Journal of Pediatrics Review


**Table 2.** Characteristics of the selected studies conducted on VLBW/VP newborns in Eastern Meditation region

Morbidity	Author(s) (Published Year)	Location	Sample Size	Type of Study
NEC	Chedid, et al. (2008) (16)	United Arab Emirates	173	Retrospective cohort
	Al Hazzani, et al. (2011) (18)	Saudi Arabia	186	Retrospective cohort
	Al-Abdi, et al. (2011) (20)	Saudi Arabia	18123	Retrospective cohort
ROP	Chedid, et al. (2008) (16)	United Arab Emirates	173	Retrospective cohort
	Wani, et al. (2014)	Kuwait	207	Retrospective study
BPD	Al Hazzani, et al. (2011) (18)	Saudi Arabia	186	Retrospective cohort
	Al-Abdi, et al. (2011) (20)	Saudi Arabia		Retrospective cohort
Pneumothorax	Al Hazzani, et al. (2011) (18)	Saudi Arabia	186	Retrospective cohort
	Al-Abdi, et al. (2011) (20)	Saudi Arabia	18123	Retrospective cohort
PVL	Chedid, et al. (2008) (16)	United Arab Emirates	173	Retrospective cohort
	Al Hazzani, et al. (2011) (18)	Saudi Arabia	186	Retrospective cohort
	Al-Abdi, et al. (2011) (20)	Saudi Arabia	18123	Retrospective cohort
PDA	Al Hazzani, et al. (2011) (18)	Saudi Arabia	186	Retrospective cohort
NDD	Khan, et al. (2012) (21)	Pakistan	110	Prospective cohort
	Mansouri (2001)	Saudi Arabia	92	Retrospective cohort

NEC: Necrotizing Enterocolitis; IVH: Intraventricular Hemorrhage; ROP: Retinopathy of Prematurity; BPD: Bronchopulmonary Dysplasia; PVI: Periventricular Leukomalacia; PDA: Patent Ductus Arteriosus; and NDD: Neurodevelopmental Delay

The current meta-analysis study evaluated mortality and morbidity as well as odds ratio on VLBW /VP infants in EMR countries. The present study was the first meta-analysis in this regard. Although most studies

had reported mortality (19 studies) or survival (6 studies) rate in their studied populations, different patient categorizations restricted the authors to include all the extracted literature in the present meta-analysis. Thus,



**Figure 4.** Meta-analysis of prevalence of mortality among VLBW newborns



**Table 3.** Meta-analysis of prevalence of various disorders among VLBW and VP newborns in Eastern Mediterranean region

Variable	VLBW Newborn				Very Preterm Newborn			
	Number of Studies (Reference)	Meta-analysis of Prevalence (%)	95%CI	I <sup>2</sup> % (P)	Number of Study (Reference)	Meta-analysis of Prevalence (%)	95%CI	I <sup>2</sup> % (P)
NEC	3 (16-18, 20)	7.0	5.0, 9.0	0.0(0.77)	1 (22)	0.0	2.0, 4.0	-
IVH	3 (16-18, 20)	19.0	10.0, 28.0	88.8(0.00)	1 (22)	1.0	1.0, 3.0	-
ROP	2 (16, 23)	14.0	10.0, 17.0	0.0(0)	3 (22, 24, 43)	17.0	2.0, 32.0	0.89(0.00)
BPD	2 (18, 20)	15.0	12.0, 19.0	0.0(0)	-	-	-	-
PT	2 (18, 20)	8.0	5.0, 10.0	0.0(0)	-	-	-	-
PVL	3 (16, 18, 20)	3.0	2.0, 5.0	0.0(0.55)	1 (22)	1.0	0.0, 1.0	-
PDA	1 (18)	31.0	25.0, 36.0	-	1 (22)	3.0	2.0, 5.0	-
NDD	2 (21, 26)	7.0	4.0, 10.0	0.0(0)	1 (21)	73.00	57.0, 85.0	-
Sepsis	1 (15)	49.0	38.0, 59.0	-	--	-	-	-

Journal of Pediatrics Review

NEC: Necrotizing Enterocolitis; IVH: Intraventricular Hemorrhage; ROP: Retinopathy of Prematurity; BPD: Bronchopulmonary Dysplasia; PT: Pneumothorax; PVL: Periventricular leukomalacia; PDA: Patent Ductus Arteriosus; and NDD: Neurodevelopmental Delay

**Table 4.** Pooled odds ratio of comorbidity in VLBW neonates

Variable	Number of Studies (Reference)	Pooled Odds Ratio	95%CI	X <sup>2</sup> (P)
CM	3 (24, 28, 29)	0.11	0.07-0.17	28.88(0.000)
Asphyxia	2 (30, 31)	1.10	0.75-1.60	2.92(0.232)
Sepsis	3 (30-32)	1.73	1.19-2.51	155.65(0.000)
NDD	1 (10)	0.03	0.00-0.66	0.000(.)

CM: Congenital Malformation; NDD: Neurodevelopmental Delay

Journal of Pediatrics Review

only 9 studies entered into meta-analysis of prevalence of mortality in VLBW neonates.

In the current study, the pooled prevalence of mortality in VLBW was 32% (95%CI:27.0-38.0). According to a study conducted by the National Institute of Child Health and Human Development (NICHD) on 9575 VLBW and VP infants, from 2003 to 2007, the overall mortality rate was 28%, and the highest mortality rate occurred in the first 12 hours of life (36). Overall, in-hospital mortality rate among VLBW infants in the developed countries was 5% in New Zealand (2009) (37), 6.5% in Korea (2009) (38), and 12.9% in the United States of America (from 2007 to 2008) (39).

In the current review study, the pooled odds ratio of mortality in VLBW infant was 2.41 (95%CI:1.887-3.069); the finding was not heterogeneous. Due to lack of con-

sistent studies, multiple analyses were not possible in the current review. In the NICHD study, the adjusted risk of early death was significantly high for infants born at 22<sup>nd</sup> to 24<sup>th</sup> weeks of gestation compared with the ones born at the 28<sup>th</sup> week (36). In the current study, meta-analysis of prevalence of mortality in the studied countries showed heterogeneity of mortality between African and Asian countries (55% vs. 23%).

An estimated by WHO in 2013, about 6.3 million children under five years died globally, of which 2.9 million were in African regions. About one-quarter of these deaths occur in the first month of life, and over two-thirds in the first seven days (40). According to WHO EMRO reports, 99% of infant deaths occur in the low- and middle-income countries (34) such as Southeast Asian and Sub-Saharan African countries (41).

Pooled mortality prevalence of VLBW infants in the studies performed before 2006 was higher than those of conducted after 2006 (46% vs. 26%). However, the difference was non-significant. There was a significant progress in child survival in the Western Pacific region. The overall estimations indicated that the mortality rate reduced by 65% from 1990 to 2011 (34). Because of the small number of studies, two of the subgroups could not be analyzed simultaneously (year of study and country).

Prevalence of mortality was different based on the sample size (under 200 cases, 26% and above 201 cases, 36%), and other factors probably had synergistic effects on this heterogeneity too. However, there was not enough information to estimate these effects. The second objective of the current systematic review was to analyze comorbidities with VLBW and VP infants. Birth weight and gestational age are two of the most important factors that predict the quality of life of neonates (41). According to the included studies, there was a wide variation in comorbidities among study results. Even the same comorbidities were defined in different ways which made the analysis difficult. However, comorbidities were categorized according to the most prevalent definition in the included studies.

The rarest comorbidity in VLBW belonged to PVL (3.0%) (16, 18, 20) and the most prevalent comorbidity was sepsis (49.0%) (15). Also, meta-analysis of regression analysis showed a significant comorbidity between VLBW and sepsis (OR=1.73) (28-30). In the VP group, the most common and the rarest comorbidity were NDD (73.0%) and NEC (0.0) (21, 22).

In the NICHD study, Respiratory Distress Syndrome (RDS) was the most common disease among VLBW infants and other comorbidities were as follows: 36% late-onset sepsis, 16% severe intraventricular hemorrhage, and 11% necrotizing enterocolitis (36). Consistent with the current study, sepsis in less than 1000 g neonates was the most frequent morbidity in Ataly study (42). In the current study, BPD was reported 15.0%. However, there was only one case in Ataly study (42). In the current study, significant differences were observed between birth weight and Apgar scores at 1 minute and 5 minutes, as well as application of nasal CPAP, ventilatory therapy, and the need for surfactant therapy.

The most restrictive factor of the current study was the limitation in the eligible studies and insufficient sample sizes. For example, mortality pooled prevalence, and univariate and multiple regression analysis in VP newborns could not be estimated. Lack of in-

formation from some of EMR countries was another important limitation. Wide discrepancy in the study design, populations, mortality indicators, and neonatal definitions of morbidities created difficulty to make a perfect picture of prematurity and low birth weight and their subgroups of these regions.

## 8. Conclusions

In conclusion, there was a relatively huge discrepancy between the prevalence of mortality among VLBW newborns in the Asian and African countries. It seems that these statistics in EMR should be reported and evaluated independently in Asian and African countries. Septicemia caused a large proportion of deaths in VLBW newborns, therefore, it is clear that the greater proportion of neonatal mortality (caused by infections) is preventable by simple interventions and vice versa; negligence in treatment has serious consequences.

## Ethical Considerations

### Compliance with ethical guidelines

Ethical code number of this study is: No. 27833.

### Funding

The current meta-analysis was financially supported by Tehran University of Medical Sciences.

### Authors contributions

The authors contributions is as follows: Conceptualization: Fatemeh Nayeri, Mamak Shariat, Setareh Sagheb; Methodology: Leyla Sahebi, Mamak Shariat, Fatemeh Nayeri; Software: Leyla Sahebi, Zahra Emami; Validation: Fatemeh Nayeri, Setareh Sagheb, Mamak Shariat, Leyla Sahebi; Formal Analysis: Leyla Sahebi; Investigation: All authors; Resources: Leyla Sahebi, Zahra Emami, Yasamin Mohammadzadeh; Data curation: Leyla Sahebi, Zahra Emami, Yasamin Mohammadzadeh; Writing—original draft preparation: Leyla Sahebi; Writing—review & editing: Fatemeh Nayeri, Mamak Shariat, Leyla Sahebi; Visualization: Fatemeh Nayeri, Mamak Shariat, Setareh Sagheb; Supervision: Fatemeh Nayeri, Mamak Shariat; and Project administration: Fatemeh Nayeri.

### Conflict of interest

The authors declare no conflict of interest.



## References

1. World Health Organization. Children reducing mortality [Internet]. 2014 [cited 2017 July 13]. Available from: <http://www.who.int/mediacentre/factsheets/fs178/en/>
2. Blencowe H, Cousens S, Oestergaard MZ, Chou D, Moller AB, Narwal R, et al. National, regional, and worldwide estimates of preterm birth rates in the year 2010 with time trends since 1990 for selected countries: A systematic analysis and implications. *The Lancet*. 2012; 379(9832):2162-72. [DOI:10.1016/S0140-6736(12)60820-4]
3. Horbar JD, Carpenter JH, Badger GJ, Kenny MJ, Soll RF, Morrow KA, et al. Mortality and neonatal morbidity among infants 501 to 1500 Grams From 2000 to 2009. *Pediatrics*. 2012; 129(6):1019-26.
4. Markestad T. Early death, morbidity, and need of treatment among extremely premature infants. *Pediatrics*. 2005; 115(5):1289-98.
5. Stoll BJ, Hansen NI, Bell EF, Walsh MC, Carlo WA, Shankaran S, et al. Trends in care practices, morbidity, and mortality of extremely preterm Neonates, 1993-2012. *Journal of the American Medical Association*. 2015; 314(10):1039-51. [DOI:10.1001/jama.2015.10244] [PMID] [PMCID]
6. Qiu X, Lodha A, Shah PS, Sankaran K, Seshia MMK, Yee W, et al. Neonatal outcomes of small for gestational age preterm infants in Canada. *American Journal of Perinatology*. 2012; 29(2):87-93. [DOI:10.1055/s-0031-1295647] [PMID]
7. Soleimani F, Zaheri F, Abdi F. Long-term neurodevelopmental outcomes after preterm birth. *Iranian Red Crescent Medical Journal*. 2014; 16(6):e17965. [DOI:10.5812/ircmj.17965]
8. Saigal S, Doyle LW. An overview of mortality and sequelae of preterm birth from infancy to adulthood. *The Lancet*. 2008; 371(9608):261-9. [DOI:10.1016/S0140-6736(08)60136-1]
9. Lawn JE, Davidge R, Paul VK, von Xylander S, de Graft Johnson J, Costello A, et al. Born too soon: Care for the preterm baby. *Reproductive Health*. 2013; 10(1):S5. [DOI:10.1186/1742-4755-10-S1-S5] [PMID] [PMCID]
10. Blencowe H, Cousens S, Chou D, Oestergaard M, Say L, Moller AB, et al. Born Too Soon: The global epidemiology of 15 million preterm births. *Reproductive Health*. 2013; 10(1):S2. [DOI:10.1186/1742-4755-10-S1-S2]
11. Ziadeh SM. Obstetrical outcomes amongst preterm singleton births. *Saudi Medical Journal*. 2001; 22(4):342-6. [PMID]
12. Fakher M, Shaaban W, Abdel Monein A, Hassan Z, Moustafa Fikry M. Statistical study of preterm infants admitted to NICU in Fawzy Moaz Hospital for children. *Alexandria Journal of Pediatrics*. 2005; 19(1):155-8.
13. Obaid KA, Al Azzawi DSH. Outcome of low birth weight infants in Diyala province of Iraq. *Journal of Tropical Pediatrics*. 2011; 57(4):280-2. [DOI:10.1093/tropej/fmq085] [PMID]
14. Dawodu A, Várady E, Nath KNR, Rajan TV. Neonatal outcome in the United Arab Emirates: The effect of changes in resources and practices. *Eastern Mediterranean Health Journal*. 2005; 11(4):673-9. [PMID]
15. Malek Mellouli M, Ben Amara F, Gallouz N, Gada H, Klilia M, Reziga H. Does the mode of delivery affect neonatal morbidity and mortality in very low-birth-weight infants? *Journal Medical Tunisie*. 2013; 91(3):183-7.
16. Chedid F, Shanteer S, Haddad H, Musharraf I, Shihab Z, Imran A, et al. Short-term outcome of very low birth weight infants in a developing country: Comparison with the Vermont Oxford network. *Journal of Tropical Pediatrics*. 2009; 55(1):15-9. [DOI:10.1093/tropej/fmn064] [PMID]
17. Hossein HJ, Kadhum NM, Muneeb A, Salih LM. Relationship between very low birth weight neonates and increased neonatal mortality for age 0 to 7 days. *Al-Kindy College Medical Journal*. 2014; 10(1):62-7.
18. Al Hazzani F, Al Alaiyan S, Hassanein J, Khadawardi E. Short-term outcome of very low-birth-weight infants in a tertiary care hospital in Saudi Arabia. *Annals of Saudi Medicine*. 2011; 31(6):581-5.
19. Alazzawi DS. Mortality rate among low birth weight infants in Al-Battool Teaching Hospital, Diyala Province, Iraq. *Iraqi Academic Scientific Journal*. 2011; 10(4):499-503.
20. Al-Abdi S, Abou Mehrem A, Dabelah K, Al Aghbari M, Al Aamri M. The effect of low to moderate patient volume on very low-birth-weight outcomes in a level III-B neonatal intensive care unit. *American Journal of Perinatology*. 2011; 28(3):219-26. [DOI:10.1055/s-0030-1268236] [PMID]
21. Khan MR, Maheshwari PK, Shamim H, Saleem AF, Ahmed S, Ali SR, et al. Neurodevelopmental outcomes of premature infants at a tertiary care center in Pakistan. *Pediatric Neurology*. 2012; 47(2):109-13. [DOI:10.1016/j.pediatrneurol.2012.05.010] [PMID]
22. Parappil H, Rahman S, Salama H, Rifai HAI, Parambil NK, Ansari WEI. Outcomes of 28+1 to 32+0 weeks gestation babies in the State of Qatar: Finding facility-based cost effective options for improving the survival of preterm neonates in low income countries. *International Journal of Environmental Research and Public Health*. 2010; 7(6):2526-42. [DOI:10.3390/ijerph7062526] [PMID] [PMCID]
23. Wani V, Al Kandari J, Al Naqeeb N, Al Serafi M, Gani M, Kazem M, et al. Type I retinopathy of prematurity in infants with birth weight less than 1251 g: Incidence and risk factors for its development in a nursery in Kuwait. *Middle East African Journal of Ophthalmology*. 2013; 20(1):66-71.
24. Arjumand S, Tikmani SS, Khan IA, Atiq H, Akhtar ASM, Kumar P, et al. Frequency of retinopathy of prematurity in premature neonates with a birth weight below 1500 grams and a gestational age less than 32 weeks: A study from a tertiary care hospital in a lower-middle income country. *PLoS One*. 2014; 9(7):e100785.

25. Nair PMC, Ganesh A, Mitra S, Ganguly SS. Retinopathy of prematurity in VLBW and extreme LBW babies. *Indian Journal of Pediatrics*. 2003; 70(4):303-6.
26. Mansouri HA. Perinatal factors and neonatal outcome of very low birth weight and extremely premature babies at KAUH. *Bahrain Medical Bulletin*. 2001; 23(2):1-10.
27. Sohaila A, Tikmani SS, Khan IA, Atiq H, Akhtar ASM, Kumar P, et al. Frequency of retinopathy of prematurity in premature neonates with a birth weight below 1500 grams and a gestational age less than 32 weeks: A study from a tertiary care hospital in a lower-middle income country. *PLoS One*. 2014; 9(7):7-10. [DOI:10.1371/journal.pone.0100785] [PMID] [PMCID]
28. Manzar N, Manzar B, Yaqoob A, Ahmed M, Kumar J. The study of etiological and demographic characteristics of neonatal mortality and morbidity-a consecutive case series study from Pakistan. *BMC Pediatrics*. 2012; 12:131. [DOI:10.1186/1471-2431-12-131] [PMID] [PMCID]
29. Sobaih B, Hadid A, Fariss A, Banoo R, AlKharfi T, AlFaleh K. Mortality and short term outcome of Very Low Birth Weight (VLBW) infants at a tertiary care center in Saudi Arabia: 9 years' data. *Kuwait Medical Journal*. 2014; 46(3):233-6.
30. SA Y. Early outcome of pre-term neonates delivered at Soba University Hospital. *Sudanese Journal of Paediatrics*. 2013; 13(2):37-44.
31. Rahman A, Abdellatif M, Sharef SW, Fazalullah M, Al Senaidi K, Khan AA, et al. Changing survival rate of infants born before 26 gestational weeks: Single-centre study. *Sultan Qaboos University Medical Journal*. 2015; 15(3):e351.
32. Nassar MM. Screening for retinopathy of prematurity: A report from upper Egypt. *International Journal of Ophthalmology*. 2016; 9(2):262-5.
33. World Health Organization. Global Health Observatory (GHO) data: Infant mortality, Situation and trends [Internet]. 2016 [Updated 2016 August 14] Available from: [http://www.who.int/gho/child\\_health/mortality/neonatal\\_text/en/](http://www.who.int/gho/child_health/mortality/neonatal_text/en/)
34. World Health Organization. Regional office for the eastern Mediterranean. Geneva: World Health Organization; 2007.
35. Mandy GT. Incidence and mortality of the preterm infant [Internet]. 2017 [cited 2017 July 13]. Available from: <http://www.uptodate.com/contents/incidence-and-mortality-of-the-preterm-infant>
36. Stoll BJ, Hansen NI, Bell EF, Shankaran S, Laptook AR, Walsh MC, et al. Neonatal outcomes of extremely preterm infants from the NICHD. *Pediatrics*. 2010; 126(3):443-56. [DOI:10.1542/peds.2009-2959] [PMID] [PMCID]
37. Battin MR, Knight DB, Kuschel CA, Howie RN. Improvement in mortality of very low birthweight infants and the changing pattern of neonatal mortality: The 50-year experience of one perinatal centre. *Journal of Paediatrics and Child Health*. 2012; 48(7):596-9. [DOI:10.1111/j.1440-1754.2012.02425.x]
38. Shim JW, Kim MJ, Kim EK, Park HK, Song ES, Lee SM, et al. The impact of neonatal care resources on regional variation in neonatal mortality among very low birthweight infants in Korea. *Paediatric and Perinatal Epidemiology*. 2013; 27(2):216-25. [DOI:10.1111/ppe.12033]
39. Staiger D, Staiger D, Horbar J, Cheung R, Kenny MJ, Patrick T, et al. Association between hospital recognition for nursing excellence and outcomes of very low-birth-weight infants. *JAMA*. 2012; 307(16):1709-16. [DOI:10.1001/jama.2012.504] [PMID] [PMCID]
40. Lobach KS. Child and Adolescent Health. *Journal of Urban Health*. 2007; 84(2):139-41.
41. The World Bank. Low income | Data [Internet]. 2017 [cited 2018 February 26]. Available from: <https://data.worldbank.org/income-level/low-income>
42. Atalay D, Salihoglu O, Can E, Beskardes A, Hatipoglu S. Short-term outcomes of very low birth weight infants born at a tertiary care hospital, Istanbul, Turkey. *Iranian Journal of Pediatrics*. 2013; 23(2):205-11. [PMID] [PMCID]
43. Nair PMC, Ganesh A, Mitra S, Ganguly SS. Retinopathy of prematurity in VLBW and extreme LBW babies. *Indian Journal of Pediatrics*. 2003; 70(4):303-6. [DOI:10.1007/BF02723585] [PMID]