# Original Article

## Relation between Mortality Rate and air pollutant concentrations in Mashhad, 2007-2009

\*Hossein Alidadi<sup>1</sup> Mohammad Taghi Shakeri<sup>2</sup> Ahmad Hossein zadeh<sup>3</sup> Anoosheh Noorani<sup>4</sup>

1-Department of Environmental Health Engineering, Health Sciences Research Center, Mashhad University of Medical Sciences, Mashhad, Iran.

2-Department of Community Medicine, Medical School, Mashhad University of Medical Sciences, Mashhad, Iran.

3-Department of Environmental Health Engineering, School of Health, Mashhad University of Medical Sciences, Mashhad, Iran.

4-Medical School, Mashhad University of Medical Sciences, Mashhad, Iran

#### \* alidadih@mums.ac.ir

### Abstract

**Background and purpose:** Air pollution has been known as one of the most effective factors on cardiovascular, pulmonary and infectious diseases. This study aims to determine the relation between mortality rate and air pollution in Mashhad during 2007-2009.

*Materials and Methods:* Data of air pollution, mortality rate and geographic factors were collected during 2007-2009 and analyzed. Mortality rate, the average PSI of individual pollutants, temperature and moisture were weekly measured and the diagrams were then drawn according to the methods of statistical correlation and regression analysis of air pollution and mortality rate.

**Results:** According to the findings, there was no significant relation between CO and mortality rate, but the other pollutants played a significant role in this regard. The maximum correlation was obtained for  $SO_2$  and  $O_3$  in the concentrations of 0.936 and 0.154 ppm, respectively, but in the final model for all pollutants and weather variables, only PSI for  $O_3$ , season and humidity showed significant change.

*Conclusion:*  $O_3$  has increased during three years of research and has been the leading cause of death among the pollutants. Most deaths occurred in cold seasons. Among all pollutants, the maximum concentration of  $SO_2$  was in winter and the maximum concentration of  $O_3$  was in spring and summer 2009. [\*Alidadi II. Shakeri M.T. Hossein zadeh II. Noorani A. The Relation between Mortality Rate and Air Pollutant concentrations in Mashhad, 2007-2009. [JHS 2013; 1(2):68-74] http://jhs.mazums.ac.ir

Key words: Air pollution, Mortality Rate, Pollutants.

#### 1. Introduction

Air pollution is a major environmental risk for human health, and its impacts on economic activities continue to be a worldwide concern. Air pollution is the presence of one or more pollutants or the combination of them in atmosphere for a certain period of time, which can be harmful to humans, plants or animal life (1,2). The major sources of air pollution are vehicles, industrial factories, smokes, fire, pollens, sand storms, storms, dust, volcanic activities and forest fires. Five air pollutants criteria including particulate matter (PM), sulfur dioxide  $(SO_2)$ , nitrogen dioxide  $(NO_2)$ , ozone  $(O_3)$  and carbon monoxide (CO) are regularly measured around the world.  $PM_{10}$ or particles which are less than 10 micron in diameter, or toxic gases such as SO<sub>2</sub> are considered to be the most hazardous because when inhaled, these particulate matters or gas can reach deep into the lungs and interfere with internal gas exchange(2,3,4,5).

In recent years, more attention has been given to the effects of air pollution on human health. In the late 20th century, episodes of extremely high air pollution have been strongly correlated with mortality (6,7). Epidemiological research over the last two decades has determined that there is a direct relationship between poor air quality and a decline in overall human health. The EPA estimates that mobile sources (car. truck and bus) of air toxics account for as much as half of all cancers attributed to outdoor sources of air toxics (8,9,10,11). In a study on the relationship between air pollution and mortality rate, it was revealed that a significant relationship exists between these two variables. As WHO mentioned, the estimated annual air pollution is responsible for 3 million premature deaths across the world. These contaminants can cause various diseases and can be influential in mortality rate. Nowadays, the role of air pollution in urban communities as an effective factor in various diseases and the has been confirmed mortality rate (4,8,10,11). Reducing pollution and thereby mortality rate seems to be important and necessary. Although several studies on the relationship between air pollution-related health complications and mortality rate have been done, there is not an accurate and appropriate methodology for this problem in our country. Most studies have been performed in the United States and Europe and such studies have been done less frequently in Asia (9, 13, 14, 18). In this study. the relationship between the frequencies of deaths associated with air pollution has been studied in Mashhad.

### 2. Materials and Methods

This longitudinal study was conducted on the relationship between mortality rate due to cardiovascular, pulmonary and infectious diseases and pollutant concentrations in the air in Mashhad during 2007 - 2009. In order to determine the levels of air pollution parameters such as carbon monoxide, sulfur dioxide, oxides of nitrogen, hydrocarbons and particles in different times of the year and deaths due to cardiovascular, pulmonary and infectious diseases in Mashhad during the year, all data were taken collected from Environmental Protection Agency, in Mashhad, Khorasan, also data of deaths due to air pollution such as cardiovascular,

pulmonary and infectious diseases in Mashhad during 2007-2009, were taken collected Emam Reza cemeteries of municipality. Information about daily concentrations of pollutants was taken from the site of the Environmental Protection Agency. The average concentration of about 24 micro pollutants smaller than 10 micrometers  $(PM_{10})$  in microgram per cubic  $(\mu g/m^3)$ 24-hour average meter concentration of Nitrogen dioxide (No<sub>2</sub>), maximum-hour concentration of ozone  $(O_3)$ . maximum 8-hour concentrations of carbon monoxide (CO) and 24-hour mean concentration of sulfur dioxide (SO2) (all in PPM) have been reported.

All data was gathered from Vahdat Station which measures the concentrations of pollutants in Mashhad, located at the nearest station to Imam Reza Holy Shrine in the city center which is the most polluted part of city. The pollutants standard index, PSI, which is an index for reporting daily air quality and usually five of the pollutants including CO, O3, NO2, SO2 and PM were used in this report to specify which of the pollutants have more adverse effects. To calculate the PSI value, the concentrations of the pollutants were transformed to a numerical scale between 0 and 500. As it measures the concentration of the pollutants based on a single standard scale, comparison between indicators becomes possible.

The equation below is used for converting the pollutants concentration to PSI. Equation:

I max-I min/Cmax-Cmin(Cmax-Cmin) +I min

In this relation  $I_{Max}$  and  $I_{min}$  are the maximum standard range and minimum standards of the desired interval, and  $C_{Max}$  and  $C_{min}$  are maximum concentration ranges and the minimum concentration intervals of the desired interval.

### 2.1. Statistical Analysis

At first, Matlab software was used for converting the daily concentrations of pollutants to standard index, then, the average standard of each pollutant was calculated for five weeks. The frequencies of mortality due to three factors were calculated in each week. The relationships between the variables were described in the graphics. The correlation coefficient of individual pollutants with the number of deaths was calculated and its significance was set. ANOVA test was used to compare the emissions in different seasons. The linear regression method was used in order to fit statistical models to determine the frequency of death and pollution.



### 3. Results

Season/Year	2007	2008	2009
Spring	512	592	784
Summer	464	746	656
Autumn	505	944	779
Winter	614	975	942
Total	2095	3227	3161

**Table1.** Prevalence of seasonal mortality (person) due to air pollution in Mashhad

Results shows that the most deaths occur in 2008 and the mortality rate in winter was more than other seasons. The table 2 shows mean and standard deviation Psi of carbon monoxide in different seasons 2007 to 2009.

Table2. Mean and standard deviation Psi of carbon monoxide at different seasons in Mashhad

Season/Year	2007	2008	2009
Spring	26.02±11.19	25.76±9.60	19.11±6.69
Summer	$25.03 \pm 7.647$	22.21±10.81	21.12±8.98
Autumn	33.038±15.99	29.80±19.03	25.94±6.54
Winter	28.79±12.02	31.30±8.86	37.29±14.56
P <sub>value</sub>	P=0.231	P=0.267	P<0.001

Results showed that high pollution and low pollution result of Psi carbon monoxide occurs in winter and spring 2008. There was significant relationship between Psi CO and seasons (P value<0.001). The table 3 shows mean and standard deviation Psi of Nitrogen dioxide in different seasons 2007 to 2009.

Table3. Mean and standard deviation Psi of Nitrogen dioxide at different seasons in Mas	hhad
---	------

Season/Year	2007	2008	2009
Spring	$17.30 \pm 1.60$	11.90±1.58	6.41±2.61
Summer	$20.62 \pm 6.30$	10.68±1.26	$6.27 \pm 2.63$
Autumn	$14.10 \pm 17.87$	13.66±3.69	$8.59 \pm 2.52$
Winter	$17.02 \pm 28.66$	$19.92 \pm 4.31$	$3.63 \pm 3.00$
P <sub>value</sub>	P=0.793	P<0.001	P<0.001

Results showed that between 2007 to 2009 Psi Nitrogen dioxide decreased and There was significant relationship between Psi CO and seasons in 2008 and 2009(P<sub>value</sub><0.001).

The table 4 shows mean and standard deviation Psi of ozone in different seasons 2007 to 2009.

Table4. Mean and standard deviation Psi of ozone at different seasons in M	Aashhad
--	---------

Season/Year	2007	2008	2009
Spring	15.16±3.12	52.71±3.72	51.10±8.44
Summer	7.02±3.86	38.60±3.67	62.51±12.61
Autumn	29.64±7.38	40.86±6.39	32.23±15.19
Winter	$42.98 \pm 8.45$	42.71±20.18	32.11±29.17
P <sub>value</sub>	P<0.001	P>0.001	P<0.001



Results showed that between 2007 to 2009 Psi Nitrogen dioxide increased and high pollution and low pollution result of Psi  $O_3$ occurs in spring, summer and autumn. There was significant relationship between Psi  $O_3$  and seasons in 2007 to 2009(P  $_{value}$ <0.001). The table 5 shows mean and standard deviation Psi of ozone in different seasons 2007 to 2009.

Table 5. Mean and standard	d deviation Psi of sulfu	r dioxide at different so	easons in Mashhad
----------------------------	--------------------------	---------------------------	-------------------

Season/Year	2007	2008	2009	
Spring	5.81±1.07	10.75±2.62	8.51±1.92	-
Summer	$5.80 \pm 0.82$	$8.62 \pm 2.08$	$11.95 \pm 2.64$	
Autumn	$15.30 \pm 3.78$	$17.09 \pm 8.49$	17.97±3.20	
Winter	$20.75\pm5.78$	14.19±3.84	$15.69 \pm 5.28$	
P <sub>value</sub>	P<0.001	P<0.001	P<0.001	

Results showed that high pollution and low pollution result of Psi SO2 occurs in autumn, winter and spring, summer. There significant was relationship between Psi SO2 and seasons in 2007 to 2009(P value<0.001). Results showed that there was no significant relationship between CO and mortality. however. other pollutants showed a significant role in the mortality. The highest correlation was for the gases SO2 and O3 in the concentrations of 0.936 and 0.154 ppm, Linear regression model respectively. was used for the role of pollutants in

mortality changes. In this model, the dependent variable was the weekly number of mortality, and the independent variables were average amount of pollutants over 3 years. This model was fitted with data every 3 years, the PSI for CO was not significant (Pvalue = 0.619). However, other pollutants showed a role significant in this model. Determining factors in this model such as regression coefficients R2=0.187 and significance of individual variables in this model is shown in the table below.

Table6. Regres	sion coefficien	ts ( <b>β)</b> of pollut	ants in Mashhad
----------------	-----------------	--------------------------	-----------------

Variable	β Coefficient	<b>P</b> <sub>value</sub>
Constant	50.158	P<0.001
$PSI(SO_2)$	0.936	P<0.001
$PSI(O_3)$	0.154	P=0.014
$PSI(PM_{10})$	0.129	P = 0.014
PSI(NO <sub>2</sub> )	0.222	P = 0.045

There was a close relationship between mortality, and air pollution and cardiovascular, infectious and pulmonary diseases. This means that increasing concentration of pollutants in the three years is associated with increasing the frequency of morbidity and mortality through the three mentioned categories of disease.

### 4. Discussion

Based on correlation coefficient during 2007-2009, the weekly average  $PM_{10}$ ,  $NO_2$ ,  $O_3$ , and  $SO_2$  showed a significant relationship between death and cardiovascular, pulmonary and infectious diseases, but CO showed no significant association with mortality. In the first statistical model, SO<sub>2</sub> and O<sub>3</sub> had the highest correlation with mortality. Other variables such as the level of socio-economic or smoking can be confounding variables and play an important role in this regard; it seems that it is one of the reasons responsible for the different results in different years of research by creating a fake relationship or by covering the real relationship. It can be due to the other causes such as environmental or statistical problems of data or different interpretations in different years. According to the study by Yang in Taiwan, NO<sub>2</sub> and CO had the greatest impact on mortality due to respiratory disease (5). General results indicated that ozone has increased during research process; it was the main cause of death among gases because of 15-20 kilometers destruction of ozone layer above the atmosphere.

Ozone depleting gases are mainly from refrigeration and cooling industries, the foam-building sponge and industries, systems of agricultural pests, air conditioning, fire capsules, solvent cleaners spray and electronics. Shelf-life of ozone depleting gases is from 50 to 150 years; to remove these gases from atmosphere, at least half a century is necessary. Kim et al., found that ozone has a strong seasonal component; it is very difficult to separate the main effect of ozone on mortality from the confounding effect of season and temperature (12).

Lee et al., reported that  $SO_2$  was significantly associated with daily mortality, but total suspended particulate was not (4). Xu et al., discovered that  $SO_2$  was significantly associated with increased mortality, but TSP was not significant (3.16).Correlation and regression relationships between climatic elements and the average weekly mortality due to cardiovascular, pulmonary and infectious diseases demonstrated that climatic elements such as temperature and humidity and the mortality are associated with a significant strong correlation. Facing decreasing temperature, the number of mortality has been increased as mortality rate increases .As long as cooling air, pressure, solar radiation and short day length reduce; the layer height increases inversion the concentration of pollutants and the density of dust in the limited space and surface. The minimum mortality rate was in spring and summer, with the relative rise of temperature and decreased pressure, number of thermal inversion. height atmospheric and concentration of pollutants. Weekly average correlation between the elements of climate suggests a strong relationship with mortality. Farajzadeh reported more mortality rate in cold season (December, January and February) (17, 19).  $SO_2$  and CO peak were in winter and maximum O<sub>3</sub> in spring and summer during 2009. Due to the increased mortality in winter in cardiopulmonary patients, it could be advised to them not to get out of their houses or to use mask.

### **Acknowledgements**

The authors would like to thank Vice Chancellor of Researches of Mashhad University of Medical Sciences for their financial support in this research.

IJHS 2013; 1(2): 73ID.ir

### References

1. Toren K, Bergdahl IA, Nilsson T. Occupational exposure to particulate air pollution and Mortality due to ischaemic heart disease and cerebrovascular disease. Occup Environ Med 2007; 64: 515–519.

2.Perez L, Medina-Ramón M, Künzli N. Size fractionate particulate matter, vehicle traffic, and case-specific daily mortality in Barcelona, Spain. Environ Sci Technol 2009; 43:4707–4714.

3. Xu X, Gao J, Dockery D. Air Pollution and Daily Mortality in Residential Areas of Beijing, China. Archives of environmental health 1994; 49:216-216.

4. Lee J, Son J, Cho Y. A comparison of mortality related to urban air particles between periods with Asian dust days and without Asian dust days in Seoul, Korea,2000-2004. Environmental Research 2007;105:409-413.

5.Yang Ch, Chang Ch, Chuang HY, Tsai ShSh, Wu TN, Ho Chk. Relationship between air pollution and daily mortality in a subtropical city: Taipei, Taiwan. Environ int 2004;30(4): 519-23.

6. Zhang P, Dong G, Sun B. Long-Term Exposure to Ambient Air Pollution and Mortality Due to Cardiovascular Disease and Cerebrovascular Disease in Shenyang, China. Journal pone. 2011;20:827.

7. Helen S .The relationship between ambient air pollution and heart rate variability differs for individuals with heart and pulmonary disease. Environmental Health Perspectives 2006.

8. Filleul L, Rondeau V, Vandentorren S. Twenty five year mortality and air pollution: results from the French PAARC survey. Occup Environ Med 2005;62:453–460.

9. Franklin M, Koutrakis P, Schwartz J. The role of particle composition on the association between PM <sub>2.5</sub> and mortality. Epidemiology 2008;19: 680–689.

10.Dominici F, Daniels M, Zeger S. Air Pollution and Mortality: Estimating Regional and National Dose–Response Relationships. Journal of the American Statistical Association. 2002;97:457

11.Murray Ch,Nelson J, Charles R. State space modeling of the relationship between air quality and mortality. J. of the Air and Waste Management Association 2000.

12.Kim S, Lee J, Hong Y. Determining the threshold effect of ozone on daily mortality: an analysis of ozone and mortality in Seoul, Korea,1995-1999.Environmental Research 2004; 94:113-119.

12.Lee J,Shin D,Chung Y, Air pollution and daily mortality in Seoul and Ulsan, Korea. Environmental Health Perspectives 1999;107:149-154.

13.Min J,Min K,Cho S. Lag effect of particulate air pollution on lung function in children. Pediatric pulmonology, 2008.

14.Welty L,Peng R,Zeger S, Dominici F. Bayesian Distributed Lag Models:Estimating Effects of Particulate Matter Air Pollution on Daily Mortality. Biometrics 2008.

15.Peng R,Dominici F, Louis T. Model choice in time series studies of air pollution and mortality. Journal of the Royal Statistical Society Series A 2006;169:179-203.

16.Kan H, London SJ, Chen G. Season, sex, age, and education as modifiers of the effects of outdoor air pollution on daily mortality in Shanghai, China: The Public Health and Air Pollution in Asia Study. Environ Health Perspect 2008;116: 1183–1188

17. Pope CA, Burnett RT, Thurston GD. Cardiovascular mortality and long term exposure to particulate air pollution: epidemiological evidence of general pathophysiological pathways of disease. Circulation 2004;109:71–7. 18.Zhang F, Krafft Th. Study on the Association between Ambient Air Pollution and Daily Cardiovascular and Respiratory Mortality in an Urban District of Beijing. Int J Environ Res Public Health 2011;8(6):2109–2123.

19.Faraj-Zadeh M. The effect of temperature on mortality in Tehran. Hakim Research Journal 2008;11(1):34-27.

IJHS 2013; 1(2); 74ID.ir