

# The Effects of the Synthetic Antioxidant, Tempol, on Serum Glucose and Lipid Profile of Diabetic and Non-Diabetic Rats

Siamak Shahidi,<sup>1\*</sup> Zahra Jabbarpour,<sup>2</sup> Masoud Saidijam,<sup>3</sup> Rasoul Esmaeili,<sup>2</sup> Alireza Komaki,<sup>1</sup> and Nasrin Hashemi Firouzi<sup>1</sup>

<sup>1</sup>Neurophysiology Research Center, Hamadan University of Medical Sciences, Hamadan, IR Iran

<sup>2</sup>Department of Physiology, School of Medicine, Hamadan University of Medical Sciences, Hamadan, IR Iran

<sup>3</sup>Research Center for Molecular Medicine, Hamadan University of Medical Sciences, Hamadan, IR Iran

\*Corresponding author: Siamak Shahidi, Neurophysiology Research Center, Hamadan University of Medical Sciences, Hamadan, IR Iran. Tel: +98-8138380462, Fax: +98-8138380208, E-mail: SiamakShahidi@yahoo.com

Received 2015 June 25; Revised 2015 July 20; Accepted 2015 July 27.

## Abstract

**Background:** Hyperlipidemia and low antioxidant levels is one the diabetes side effects. Some studies have indicated the possible effects of nutrients on the improvement of hyperlipidemia, by their antioxidants ingredients.

**Objectives:** The aim of the present study was to evaluate the effect of the synthetic antioxidant, tempol, on blood lipid profiles and glucose levels in healthy and diabetic rats.

**Materials and Methods:** Adult Wistar rats were randomly divided to four experimental groups including, healthy control, diabetic control, diabetic receiving tempol and healthy receiving tempol groups. Diabetes was induced by injection of streptozotocin (60 mg/kg, Intraperitoneally (IP)). The rats were then fed saline or tempol (30 mg/kg) by gavage for 60 days. Blood samples were collected by cardiac puncture. Next, glucose, high-density lipoprotein (HDL), low-density lipoprotein (LDL), very low-density lipoprotein (VLDL), cholesterol, triglyceride and HbA1c were measured by specific kits. Also, the coronary risk index was calculated.

**Results:** The blood glucose level increased following diabetes induction. The level of blood glucose in the diabetic receiving tempol group decreased compared to the control diabetic group. The comparison of LDL, VLDL, cholesterol, triglyceride, HbA1c and coronary risk index among experimental groups indicated the increase of these factors in the diabetic group. High-density lipoprotein in the diabetic groups was lower than the other groups.

**Conclusions:** It can be concluded that tempol can improve dyslipidemia and may decrease hyperglycemia in diabetes. It seems that antioxidants such as tempol can improve dyslipidemia and may decrease hyperglycemia in diabetes.

**Keywords:** Tempol, Lipid Profiles, Diabetes, Rat

## 1. Background

Diabetes mellitus is a major public health problem throughout the world, and is the leading cause of global mortality (1). Diabetic disease is characterized by hyperglycemia, which is the accumulation of free glucose in the blood. Hyperglycemia induces oxidative stress via glucose autoxidation and leads to generation of free radicals due to autoxidation of glucose and glycosylation of proteins (2-4) and has an important role in the development of diabetic complications (5-7). Diabetes is likely to increase the risk of developing various metabolic disorders, including hyperlipidemia, liver-kidney dysfunctions, and hypertension (8).

On the other hand, hyperlipidemia has been observed in diabetic patients (9-11) and experimental diabetic animal models (12-14). Triglyceride levels are enhanced under diabetic conditions (5-7). In experimental models of diabetes, high glucose induced oxidative stress (15), increases in oxidative stress related to lipid, and inhibition of the

synthesis of endogenous antioxidants (16, 17). Oxidative stress and triglyceride levels are enhanced in patients with diabetes (2).

In cure of diabetes it is important to prevent diabetes complications. The current evidence suggests that supplementation of antioxidant compounds may protect against diabetic complications (15, 18-21). There are evidences about the protection effect of antioxidant compounds against diabetic problems (15, 18-23).

Tempol (4-hydroxy-2,2,6,6-tetramethylpiperidine-1-oxyl) is a member of a family of nitroxide compounds that is an efficient scavenger of free radicals (24) and improves insulin responsiveness and dyslipidemia in models of diabetes mellitus (25). The anti-inflammatory, neuroprotective effects of tempol have been shown previously (26-30). Furthermore, it is an efficient scavenger of free radicals and improves diabetes-associated dyslipidemia (25) and cardiac fibrosis in rats (31).

## 2. Objectives

There is no direct study on the efficient activity of tempol on hyperlipidemia in diabetes. Therefore, this study aimed to test the hypothesis that chronic oral administration of tempol could ameliorate hyperlipidemia in a diabetes rat model.

## 3. Materials and Methods

### 3.1. Animals

Forty Wistar rats, weighting 200 - 300 g, were supplied by the breeding colony of the Iran Pasteur institute, Tehran. They were maintained at  $20 \pm 2^\circ\text{C}$  on a 12-hour light/dark cycle (lights on 07:00 am). Water and food were available ad libitum. All rats were acclimatized to the environment for one week prior to initiation of testing. All procedures for the treatment of animals were approved by the research committee of the Hamadan university of medical sciences.

### 3.2. Induction of Diabetes and Treatment

The animals were divided to the following groups; control (C), diabetic (D), diabetic tempol treatment (D + T), and control group receiving tempol (C + T). The model of type I diabetes was induced by a single dose of intraperitoneal (IP) injection of 60 mg/kg of streptozotocin (32-34). The control rats received IP injections of physiological saline. Blood samples were taken from the tail vein, and glucose levels were determined using a strip-operated blood glucose sensor (Accucheck; Roche, Mannheim, Germany). One week after streptozotocin injection, the rats with blood glucose levels exceeding 250 mg/dL were considered diabetic. Tempol (30 mg/kg; Sigma) was administered to D + T and C + T groups by the gavage process, every day for two months. The control and non-treated diabetes groups received physiological saline with the same volume.

At the end of the treatment period, the rats were anaesthetized with ketamine (100 mg/kg) and blood samples were collected by cardiac puncture. Next, glucose, high-density lipoprotein (HDL), low-density lipoprotein (LDL), cholesterol and triglyceride were measured by their specific kits (Biolabo, France). The HbA1c was assessed by its specific kit (Bionik, Iran) and the turbidimetry method. The concentration of very low density-lipoprotein (VLDL) was calculated as TG/5. Also, the coronary risk index was calculated (34-36).

### 3.3. Statistical Analysis

One-way analysis of variance (ANOVA) was used to determine the statistical significant differences between experimental groups, which were followed by Tukey's post hoc test. P values of  $<0.05$  were considered statistically significant. All data were represented as mean  $\pm$  standard error of the mean (SEM).


## 4. Results

Figure 1 shows the glucose levels of rat groups, after diabetes induction. One-way ANOVA proved that there were significant differences between the groups after induction of diabetes ( $P < 0.01$ ). Tukey's post hoc test revealed that blood glucose levels of diabetic induction groups were significantly higher than non-diabetic groups (Figure 1A,  $P < 0.01$ ). Moreover, one-way ANOVA showed that there were significant differences among the experimental groups of rats after tempol treatment. Tukey's post hoc test revealed that tempol-treated rats had a significant decrease in their plasma glucose compared to untreated diabetic rats at the end of the experiment (Figure 1B,  $P < 0.01$ ).

Figure 2 shows the LDL, VLDL, cholesterol and triglyceride levels in plasma. One-way ANOVA demonstrated that there is a significant difference between the groups in the level of LDL, VLDL, cholesterol and triglyceride in serum plasma. There was a significant increase in the level of LDL in plasma of non-treated diabetic rats in comparison with other groups ( $P < 0.05$ ; Figure 2A). Also, non-treated diabetic rats had a significant increase in their plasma levels of LDL, VLDL, cholesterol and triglyceride in comparison with other groups ( $P < 0.01$ ; Figure 2B - D, respectively). The diabetic rats that received tempol exhibited significantly lower LDL, VLDL, cholesterol and triglyceride compared to the non-treated diabetic group.

Figure 3 shows the HDL level in plasma of the rat groups. One-way ANOVA suggested that there was a significant difference between the groups. The level of plasma HDL in non-treated diabetic rats was significantly lower than other experimental groups ( $P < 0.01$ ).

Figure 4 illustrates the HbA1c and coronary risk factor in the experimental groups. One-way ANOVA approves that there was a significant difference between the groups ( $P < 0.05$ ). Tukey's post hoc test revealed that HbA1c and the coronary risk factor in the streptozotocin (STZ)-receiving rats were significantly higher than non-diabetic groups ( $P < 0.01$ ; Figure 4A and B, respectively). The STZ-induced diabetic rats were administrated tempol and showed lower HbA1c and coronary risk factor when compared to the non-treated diabetic rats ( $P < 0.05$ ).

**Figure 1.** Blood Glucose Levels in Experimental Groups

A, Blood glucose levels after induction of diabetes; B, Blood glucose levels after induction of diabetes at the end of the study. C, control group; C + T, control group receiving tempol; D, diabetic group; D + T, diabetic group that received tempol. \*\*: ( $P < 0.01$ ) as compared with the control. \*\*: ( $P < 0.01$ ), ##: ( $P < 0.01$ ) and \$\$: ( $P < 0.01$ ) as compared with the control, control group receiving tempol or diabetic group, respectively. The values represent means  $\pm$  standard error of the mean (SEM) ( $n = 10$  per group). Each symbol on a column compares the mentioned group with the specified group.

## 5. Discussion

The present findings demonstrated that administration of tempol for 60 days, improved the blood lipid profiles and hyperglycemia in diabetic rats.

Tempol treated diabetic rats had lower glucose and percentage of HbA1C levels than the diabetic rats. The HbA1c is evaluated in long-term control of diabetes. Glycated hemoglobin reflects the previous two to three months of glycemic control (37). These results confirm previous studies that tempol has hypoglycemic properties (25, 38, 39). Our results are similar to a previous study that demonstrated oral tempol treatment of diabetic mice during eight weeks, reduced plasma glucose (40). Ten weeks of oral tempol administration to rats, that were fed high fat diets, decreased plasma glucose (41), and improved insulin sensitivity in obese diabetic rats (42). Reactive oxygen species (ROS) are involved in many of the complications of

diabetes (43, 44), and generation of ROS is prevented with tempol in pancreatic islet cells (45) and in diabetic mice (40).

Tempol acts by several mechanisms such as by alleviation effects on insulin resistance (46), enhancing insulin secretion from rat cultured pancreatic islet cells (38), and increasing the membrane abundance of the glucose transporter-1 and enhanced glucose uptake (47). Oxygen-derived free radicals are easily produced in diabetic disease and have important roles in the development of diabetic complications (5-8, 44, 48). Tempol is a superoxide dismutase mimetic and efficient scavenger of free radicals (49).


In diabetes, alleviated blood lipid profile is due to increased absorption of cholesterol from the intestine by a carrier of cholesterol acetyltransferase (50, 51). Plasma cholesterol, triglycerides are raised and hyperlipidemia is distinguishable in diabetes (16, 17). Lack of insulin raises free fatty acid mobilization from adipose tissue, which is followed by production of cholesterol rich LDL particles and dyslipidemia (16, 52). On the other hand, production of oxygen free radicals increases in hypercholesterolemia (53).

Antioxidants are a normal defense mechanism of the cell and are involved in the termination of the lipid peroxidation process (4). Tempol decrease VLDL and total cholesterol and increase HDL (40, 54). Antioxidants protected the polyunsaturated fatty acids (a major component of cell membranes) from oxygen free radical attack in diabetes (55) and end the peroxidation events (4).

Another finding of the present study demonstrated that tempol has advantages for coronary risk factor. Increased reactive oxygen species cause oxidative myocardial injury and diabetic cardiomyopathy (56, 57). Evidence has demonstrated specific cardiomyopathy associated with diabetes such as cardiomyopathy, cardiac dysfunction and cardiovascular disease in humans and rats (58-61). The beneficial effect of tempol was shown on blood pressure (62), cardiac fibrosis (63) and amelioration cardiac dysfunction in diabetic rats (31). The potential antioxidant role of tempol was shown in decreasing reactive oxygen species and amelioration of cardiac dysfunction (31).


In conclusion, this study confirmed that tempol improved blood lipid profiles, hyperglycemia and coronary risk factor in the diabetic rats. It could prevent the development of diabetic complications. In order to more precisely determine the mechanism of the present findings, measurement of oxidative stress indexes is recommended for the future studies.

**Figure 2.** The Lipid Profile in Experimental Groups



Effect of 60 days of treatment on A, LDL level in serum; B, VLDL level in serum; C, cholesterol level in serum; D, Triglyceride level in serum. Groups are C (control group), C + T (control group that received temporal), D (diabetic group), D + T (diabetic group that received temporal). Groups are C (control group), C + T (control group that received temporal), D (diabetic group), D + T (diabetic group that received temporal). \*: (P < 0.05), and \*\*: (P < 0.01) as compared with non-treated diabetic group. The values represent the mean  $\pm$  SEM (n = 10 per group).

**Figure 3.** High-Density Lipoprotein Levels at the End of the Study




Groups are C (control group), C + T (control group that received temporal), D (diabetic group), D + T (diabetic group that received temporal). \*\*, (P < 0.01) as compared with non-treated diabetic group. The values represent mean  $\pm$  SEM (n = 10 per group).

### Footnotes

**Authors' Contribution:** Study concept and design, analysis and interpretation of data, and critical revision of the manuscript for important intellectual content: Siamak Shahidi, Zahra Jabbarpour, Masoud Saidijam, Rasoul Esmaeili, Alireza Komaki and Nasrin Hashemi-Firouzi; acquisition of data, statistical analysis, and drafting of the manuscript: Siamak Shahidi, Zahra Jabbarpour, Rasoul Esmaeili and Alireza Komaki; administrative, technical and material support: Siamak Shahidi, Zahra Jabbarpour, Masoud Saidijam, Rasoul Esmaeili and Alireza Komaki; study supervision: Siamak Shahidi.

**Funding/Support:** This work was supported by a grant from Hamadan University of Medical Sciences.

**Figure 4.** HbA1c and Coronary Risk Index in the Experimental Groups



A, effect of 60 days of treatment on percentage of HbA1c; B, effect of 60 days of treatment on percentage of coronary risk factor; Groups are C (control group), C + T (control group that received temporal), D (diabetic group), D + T (diabetic group that received temporal). \*\*: ( $P < 0.01$ ) as compared with non-treated diabetic group. #: ( $P < 0.05$ ) as compared to diabetic rats. The values represent mean  $\pm$  SEM ( $n = 10$  per group).

## References

- Herman WH, Zimmet P. Type 2 diabetes: an epidemic requiring global attention and urgent action. *Diabetes Care*. 2012;**35**(5):943-4. doi: [10.2337/dc12-0298](https://doi.org/10.2337/dc12-0298). [PubMed: [22517937](https://pubmed.ncbi.nlm.nih.gov/22517937/)].
- Ziegler D, Sohr CG, Nourooz-Zadeh J. Oxidative stress and antioxidant defense in relation to the severity of diabetic polyneuropathy and cardiovascular autonomic neuropathy. *Diabetes Care*. 2004;**27**(9):2178-83. [PubMed: [15333481](https://pubmed.ncbi.nlm.nih.gov/15333481/)].
- Wang WT, Lee P, Yeh HW, Smirnova IV, Choi IY. Effects of acute and chronic hyperglycemia on the neurochemical profiles in the rat brain with streptozotocin-induced diabetes detected using in vivo (1)H MR spectroscopy at 9.4 T. *J Neurochem*. 2012;**121**(3):407-17. doi: [10.1111/j.1471-4159.2012.07698.x](https://doi.org/10.1111/j.1471-4159.2012.07698.x). [PubMed: [22353009](https://pubmed.ncbi.nlm.nih.gov/22353009/)].
- Valko M, Leibfritz D, Moncol J, Cronin MT, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol*. 2007;**39**(1):44-84. doi: [10.1016/j.biocel.2006.07.001](https://doi.org/10.1016/j.biocel.2006.07.001). [PubMed: [16978905](https://pubmed.ncbi.nlm.nih.gov/16978905/)].
- West IC. Radicals and oxidative stress in diabetes. *Diabet Med*. 2000;**17**(3):171-80. [PubMed: [10784220](https://pubmed.ncbi.nlm.nih.gov/10784220/)].
- Jakus V. The role of free radicals, oxidative stress and antioxidant systems in diabetic vascular disease. *Bratisl Lek Listy*. 2000;**101**(10):541-51. [PubMed: [11218944](https://pubmed.ncbi.nlm.nih.gov/11218944/)].

- Gurler B, Vural H, Yilmaz N, Oguz H, Satici A, Aksoy N. The role of oxidative stress in diabetic retinopathy. *Eye (Lond)*. 2000;**14 Pt 5**:730-5. doi: [10.1038/eye.2000.193](https://doi.org/10.1038/eye.2000.193). [PubMed: [11116694](https://pubmed.ncbi.nlm.nih.gov/11116694/)].
- Liu PY, Lin LY, Lin HJ, Hsia CH, Hung YR, Yeh HI, et al. Pitavastatin and Atorvastatin double-blind randomized comparative study among high-risk patients, including those with Type 2 diabetes mellitus, in Taiwan (PAPAGO-T Study). *PLoS One*. 2013;**8**(10):e76298. doi: [10.1371/journal.pone.0076298](https://doi.org/10.1371/journal.pone.0076298). [PubMed: [24098467](https://pubmed.ncbi.nlm.nih.gov/24098467/)].
- Reaven GM, Greenfield MS. Diabetic hypertriglyceridemia: evidence for three clinical syndromes. *Diabetes*. 1981;**30**(Suppl 2):66-75. [PubMed: [7028540](https://pubmed.ncbi.nlm.nih.gov/7028540/)].
- Nikkila EA, Kekki M. Plasma triglyceride transport kinetics in diabetes mellitus. *Metabolism*. 1973;**22**(1):1-22. [PubMed: [4628966](https://pubmed.ncbi.nlm.nih.gov/4628966/)].
- Nikkila EA, Hormila P. Serum lipids and lipoproteins in insulin-treated diabetes. Demonstration of increased high density lipoprotein concentrations. *Diabetes*. 1978;**27**(11):1078-86. [PubMed: [214367](https://pubmed.ncbi.nlm.nih.gov/214367/)].
- Man ZW, Zhu M, Noma Y, Toide K, Sato T, Asahi Y, et al. Impaired beta-cell function and deposition of fat droplets in the pancreas as a consequence of hypertriglyceridemia in OLETF rat, a model of spontaneous NIDDM. *Diabetes*. 1997;**46**(11):1718-24. [PubMed: [9356017](https://pubmed.ncbi.nlm.nih.gov/9356017/)].
- Ito M, Kondo Y, Nakatani A, Hayashi K, Naruse A. Characterization of low dose streptozotocin-induced progressive diabetes in mice. *Environ Toxicol Pharmacol*. 2001;**9**(3):71-8. [PubMed: [11167151](https://pubmed.ncbi.nlm.nih.gov/11167151/)].
- Hirano T, Mamo JC, Takeuchi H, Nagano S, Takahashi T. Correlation of insulin deficiency and hypertriglyceridemia in diabetic rats. *Diabetes Res Clin Pract*. 1991;**12**(3):173-80. [PubMed: [1889346](https://pubmed.ncbi.nlm.nih.gov/1889346/)].
- Da Ros R, Assaloni R, Ceriello A. Antioxidant therapy in diabetic complications: what is new?. *Curr Vasc Pharmacol*. 2004;**2**(4):335-41. [PubMed: [15320813](https://pubmed.ncbi.nlm.nih.gov/15320813/)].
- Latha M, Pari L. Preventive effects of Cassia auriculata L. flowers on brain lipid peroxidation in rats treated with streptozotocin. *Mol Cell Biochem*. 2003;**243**(1-2):23-8. [PubMed: [12619885](https://pubmed.ncbi.nlm.nih.gov/12619885/)].
- Choi JS, Yokozawa T, Oura H. Improvement of hyperglycemia and hyperlipemia in streptozotocin-diabetic rats by a methanolic extract of Prunus davidiana stems and its main component, prunin. *Planta Med*. 1991;**57**(3):208-11. doi: [10.1055/s-2006-960075](https://doi.org/10.1055/s-2006-960075). [PubMed: [1896517](https://pubmed.ncbi.nlm.nih.gov/1896517/)].
- Sivitz WI, Yorek MA. Mitochondrial dysfunction in diabetes: from molecular mechanisms to functional significance and therapeutic opportunities. *Antioxid Redox Signal*. 2010;**12**(4):537-77. doi: [10.1089/ars.2009.2531](https://doi.org/10.1089/ars.2009.2531). [PubMed: [19650713](https://pubmed.ncbi.nlm.nih.gov/19650713/)].
- Juranek I, Horakova L, Rackova L, Stefek M. Antioxidants in treating pathologies involving oxidative damage: an update on medicinal chemistry and biological activity of stobadine and related pyridindoles. *Curr Med Chem*. 2010;**17**(6):552-70. [PubMed: [20015031](https://pubmed.ncbi.nlm.nih.gov/20015031/)].
- Cumaoglu A, Ozansoy G, Irat AM, Aricioglu A, Karasu C, Ari N. Effect of long term, non cholesterol lowering dose of fluvastatin treatment on oxidative stress in brain and peripheral tissues of streptozotocin-diabetic rats. *Eur J Pharmacol*. 2011;**654**(1):80-5. doi: [10.1016/j.ejphar.2010.11.035](https://doi.org/10.1016/j.ejphar.2010.11.035). [PubMed: [21172345](https://pubmed.ncbi.nlm.nih.gov/21172345/)].
- Ceriello A. Controlling oxidative stress as a novel molecular approach to protecting the vascular wall in diabetes. *Curr Opin Lipidol*. 2006;**17**(5):510-8. doi: [10.1097/01.mol.0000245256.17764.fb](https://doi.org/10.1097/01.mol.0000245256.17764.fb). [PubMed: [16960499](https://pubmed.ncbi.nlm.nih.gov/16960499/)].
- Maritim AC, Sanders RA, Watkins J3. Effects of alpha-lipoic acid on biomarkers of oxidative stress in streptozotocin-induced diabetic rats. *J Nutr Biochem*. 2003;**14**(5):288-94. [PubMed: [12832033](https://pubmed.ncbi.nlm.nih.gov/12832033/)].
- Maritim A, Dene BA, Sanders RA, Watkins JB. Effects of pycnogenol treatment on oxidative stress in streptozotocin-induced diabetic rats. *J Biochem Mol Toxicol*. 2003;**17**(3):193-9. doi: [10.1002/jbt.10078](https://doi.org/10.1002/jbt.10078). [PubMed: [12815616](https://pubmed.ncbi.nlm.nih.gov/12815616/)].
- Carroll RT, Galatsis P, Borosky S, Kopec KK, Kumar V, Althaus JS, et al. 4-Hydroxy-2,2,6,6-tetramethylpiperidine-1-oxyl (Tempol) inhibits peroxynitrite-mediated phenol nitration. *Chem Res Toxicol*. 2000;**13**(4):294-300. [PubMed: [10775330](https://pubmed.ncbi.nlm.nih.gov/10775330/)].



25. Wilcox CS. Effects of tempol and redox-cycling nitroxides in models of oxidative stress. *Pharmacol Ther.* 2010;**126**(2):119-45. doi: [10.1016/j.pharmthera.2010.01.003](https://doi.org/10.1016/j.pharmthera.2010.01.003). [PubMed: [20153367](https://pubmed.ncbi.nlm.nih.gov/20153367/)].
26. Wilcox CS, Pearlman A. Chemistry and antihypertensive effects of tempol and other nitroxides. *Pharmacol Rev.* 2008;**60**(4):418-69. doi: [10.1124/pr.108.000240](https://doi.org/10.1124/pr.108.000240). [PubMed: [19112152](https://pubmed.ncbi.nlm.nih.gov/19112152/)].
27. Volk T, Hensel M, Schuster H, Kox WJ. Secretion of MCP-1 and IL-6 by cytokine stimulated production of reactive oxygen species in endothelial cells. *Mol Cell Biochem.* 2000;**206**(1-2):105-12. [PubMed: [10839200](https://pubmed.ncbi.nlm.nih.gov/10839200/)].
28. Kurihara N, Yanagisawa H, Sato M, Tien CK, Wada O. Increased renal vascular resistance in zinc-deficient rats: role of nitric oxide and superoxide. *Clin Exp Pharmacol Physiol.* 2002;**29**(12):1096-104. [PubMed: [12390298](https://pubmed.ncbi.nlm.nih.gov/12390298/)].
29. Di Paola R, Mazzon E, Zito D, Maiera D, Britti D, Genovese T, et al. Effects of Tempol, a membrane-permeable radical scavenger, in a rodent model periodontitis. *J Clin Periodontol.* 2005;**32**(10):1062-8. doi: [10.1111/j.1600-051X.2005.00818.x](https://doi.org/10.1111/j.1600-051X.2005.00818.x). [PubMed: [16174269](https://pubmed.ncbi.nlm.nih.gov/16174269/)].
30. Cuzzocrea S, McDonald MC, Mota-Filipe H, Mazzon E, Costantino G, Britti D, et al. Beneficial effects of tempol, a membrane-permeable radical scavenger, in a rodent model of collagen-induced arthritis. *Arthritis Rheum.* 2000;**43**(2):320-8. doi: [10.1002/1529-0131\(200002\)43:320:aid-ar1131.3.0.co;2-9](https://doi.org/10.1002/1529-0131(200002)43:320:aid-ar1131.3.0.co;2-9). [PubMed: [10693871](https://pubmed.ncbi.nlm.nih.gov/10693871/)].
31. Taye A, Abouzied MM, Mohafez OM. Tempol ameliorates cardiac fibrosis in streptozotocin-induced diabetic rats: role of oxidative stress in diabetic cardiomyopathy. *Naunyn Schmiedeberg Arch Pharmacol.* 2013;**386**(12):1071-80. doi: [10.1007/s00210-013-0904-x](https://doi.org/10.1007/s00210-013-0904-x). [PubMed: [23949118](https://pubmed.ncbi.nlm.nih.gov/23949118/)].
32. Cheta D. Animal models of type I (insulin-dependent) diabetes mellitus. *J Pediatr Endocrinol Metab.* 1998;**11**(1):11-9. [PubMed: [9642624](https://pubmed.ncbi.nlm.nih.gov/9642624/)].
33. Hasanein P, Shahidi S. Effects of combined treatment with vitamins C and E on passive avoidance learning and memory in diabetic rats. *Neurobiol Learn Mem.* 2010;**93**(4):472-8. doi: [10.1016/j.nlm.2010.01.004](https://doi.org/10.1016/j.nlm.2010.01.004). [PubMed: [20085822](https://pubmed.ncbi.nlm.nih.gov/20085822/)].
34. Ishiguro T, Seki M, Yokota M, Uchiyama M, Yamamoto T. [Coronary risk index for pre-operative evaluation of ischemic heart disease]. *Ma-sui.* 1995;**44**(1):51-9. [PubMed: [7699824](https://pubmed.ncbi.nlm.nih.gov/7699824/)].
35. Adeneye AA. Methanol seed extract of *Citrus paradisi* Macfad lowers blood glucose, lipids and cardiovascular disease risk indices in normal Wistar rats. *Nig Q J Hosp Med.* 2008;**18**(1):16-20. [PubMed: [19062465](https://pubmed.ncbi.nlm.nih.gov/19062465/)].
36. Ojezele MO, Abatan OM. Hypoglycaemic and coronary risk index lowering effects of *Bauhinia thonningii* in alloxan induced diabetic rats. *Afr Health Sci.* 2011;**11**(1):85-9. [PubMed: [21572862](https://pubmed.ncbi.nlm.nih.gov/21572862/)].
37. Rahbar S. The discovery of glycated hemoglobin: A major event in the study of nonenzymatic chemistry in biological systems. *Ann NY Acad Sci.* 2005;**1043**:9-19. doi: [10.1196/annals.1333.002](https://doi.org/10.1196/annals.1333.002). [PubMed: [16037217](https://pubmed.ncbi.nlm.nih.gov/16037217/)].
38. Tang C, Han P, Oprescu AI, Lee SC, Gyulchandanyan AV, Chan GN, et al. Evidence for a role of superoxide generation in glucose-induced beta-cell dysfunction in vivo. *Diabetes.* 2007;**56**(11):2722-31. doi: [10.2337/db07-0279](https://doi.org/10.2337/db07-0279). [PubMed: [17682092](https://pubmed.ncbi.nlm.nih.gov/17682092/)].
39. Jabbarpour Z, Shahidi S, Saidijam M, Sarihi A, Hassanzadeh T, Esmaeili R. Effect of tempol on the passive avoidance and novel object recognition task in diabetic rats. *Brain Res Bull.* 2014;**101**:51-6. doi: [10.1016/j.brainresbull.2013.12.013](https://doi.org/10.1016/j.brainresbull.2013.12.013). [PubMed: [24412412](https://pubmed.ncbi.nlm.nih.gov/24412412/)].
40. San Martin A, Du P, Dikalova A, Lassegue B, Aleman M, Gongora MC, et al. Reactive oxygen species-selective regulation of aortic inflammation gene expression in Type 2 diabetes. *Am J Physiol Heart Circ Physiol.* 2007;**292**(5):H2073-82. doi: [10.1152/ajpheart.00943.2006](https://doi.org/10.1152/ajpheart.00943.2006). [PubMed: [17237245](https://pubmed.ncbi.nlm.nih.gov/17237245/)].
41. Ebenezer PJ, Mariappan N, Elks CM, Haque M, Francis J. Diet-induced renal changes in Zucker rats are ameliorated by the superoxide dismutase mimetic TEMPOL. *Obesity (Silver Spring).* 2009;**17**(11):1994-2002. doi: [10.1038/oby.2009.137](https://doi.org/10.1038/oby.2009.137). [PubMed: [19424163](https://pubmed.ncbi.nlm.nih.gov/19424163/)].
42. Rafikova O, Salah EM, Tofovic SP. Renal and metabolic effects of tempol in obese ZSF1 rats—distinct role for superoxide and hydrogen peroxide in diabetic renal injury. *Metabolism.* 2008;**57**(10):1434-44. doi: [10.1016/j.metabol.2008.05.014](https://doi.org/10.1016/j.metabol.2008.05.014). [PubMed: [18803950](https://pubmed.ncbi.nlm.nih.gov/18803950/)].
43. Simonsen U, Christensen FH, Buus NH. The effect of tempol on endothelium-dependent vasodilatation and blood pressure. *Pharmacol Ther.* 2009;**122**(2):109-24. doi: [10.1016/j.pharmthera.2009.02.002](https://doi.org/10.1016/j.pharmthera.2009.02.002). [PubMed: [19268689](https://pubmed.ncbi.nlm.nih.gov/19268689/)].
44. Brownlee M. The pathobiology of diabetic complications: a unifying mechanism. *Diabetes.* 2005;**54**(6):1615-25. [PubMed: [15919781](https://pubmed.ncbi.nlm.nih.gov/15919781/)].
45. Oprescu AI, Bikopoulos G, Naassan A, Allister EM, Tang C, Park E, et al. Free fatty acid-induced reduction in glucose-stimulated insulin secretion: evidence for a role of oxidative stress in vitro and in vivo. *Diabetes.* 2007;**56**(12):2927-37. doi: [10.2337/db07-0075](https://doi.org/10.2337/db07-0075). [PubMed: [17717282](https://pubmed.ncbi.nlm.nih.gov/17717282/)].
46. Banday AA, Marwaha A, Tallam LS, Lokhandwala MF. Tempol reduces oxidative stress, improves insulin sensitivity, decreases renal dopamine D1 receptor hyperphosphorylation, and restores D1 receptor-G-protein coupling and function in obese Zucker rats. *Diabetes.* 2005;**54**(7):2219-26. [PubMed: [15983225](https://pubmed.ncbi.nlm.nih.gov/15983225/)].
47. Alpert E, Altman H, Totary H, Gruzman A, Barnea D, Barash V, et al. 4-Hydroxy tempol-induced impairment of mitochondrial function and augmentation of glucose transport in vascular endothelial and smooth muscle cells. *Biochem Pharmacol.* 2004;**67**(10):1985-95. doi: [10.1016/j.bcp.2004.02.005](https://doi.org/10.1016/j.bcp.2004.02.005). [PubMed: [15130774](https://pubmed.ncbi.nlm.nih.gov/15130774/)].
48. Simmons RA. Developmental origins of diabetes: The role of oxidative stress. *Free Radic Biol Med.* 2006;**40**(6):917-22. doi: [10.1016/j.freeradbiomed.2005.12.018](https://doi.org/10.1016/j.freeradbiomed.2005.12.018). [PubMed: [16540386](https://pubmed.ncbi.nlm.nih.gov/16540386/)].
49. Nilsson UA, Olsson LI, Carlin G, Bylund-Fellenius AC. Inhibition of lipid peroxidation by spin labels. Relationships between structure and function. *J Biol Chem.* 1989;**264**(19):11131-5. [PubMed: [2738061](https://pubmed.ncbi.nlm.nih.gov/2738061/)].
50. Mochizuki H, Takido J, Oda H, Yokogoshi H. Improving effect of dietary taurine on marked hypercholesterolemia induced by a high-cholesterol diet in streptozotocin-induced diabetic rats. *Biosci Biotechnol Biochem.* 1999;**63**(11):1984-7. [PubMed: [10635563](https://pubmed.ncbi.nlm.nih.gov/10635563/)].
51. Kusunoki J, Aragane K, Kitamine T, Kozono H, Kano K, Fujinami K, et al. Postprandial hyperlipidemia in streptozotocin-induced diabetic rats is due to abnormal increase in intestinal acyl coenzyme A:cholesterol acyltransferase activity. *Arterioscler Thromb Vasc Biol.* 2000;**20**(1):171-8. [PubMed: [10634814](https://pubmed.ncbi.nlm.nih.gov/10634814/)].
52. Holvoet P, De Keyser D, Jacobs DJ. Oxidized LDL and the metabolic syndrome. *Future Lipidol.* 2008;**3**(6):637-49. doi: [10.2217/17460875.3.6.637](https://doi.org/10.2217/17460875.3.6.637). [PubMed: [19802339](https://pubmed.ncbi.nlm.nih.gov/19802339/)].
53. Muggle A, Brandes RP, Boger RH, Dwenger A, Bode-Boger S, Kienke S, et al. Vascular release of superoxide radicals is enhanced in hypercholesterolemic rabbits. *J Cardiovasc Pharmacol.* 1994;**24**(6):994-8. [PubMed: [7898085](https://pubmed.ncbi.nlm.nih.gov/7898085/)].
54. Gonzalez-Flecha B, Reides C, Cutrin JC, Llesuy SF, Boveris A. Oxidative stress produced by suprahepatic occlusion and reperfusion. *Hepatology.* 1993;**18**(4):881-9. [PubMed: [8406364](https://pubmed.ncbi.nlm.nih.gov/8406364/)].
55. Yoshida M, Kimura H, Kyuki K, Ito M. Combined effect of vitamin E and insulin on cataracts of diabetic rats fed a high cholesterol diet. *Biol Pharm Bull.* 2004;**27**(3):338-44. [PubMed: [14993799](https://pubmed.ncbi.nlm.nih.gov/14993799/)].
56. Yu W, Wu J, Cai F, Xiang J, Zha W, Fan D, et al. Curcumin alleviates diabetic cardiomyopathy in experimental diabetic rats. *PLoS One.* 2012;**7**(12):e52013. doi: [10.1371/journal.pone.0052013](https://doi.org/10.1371/journal.pone.0052013). [PubMed: [23251674](https://pubmed.ncbi.nlm.nih.gov/23251674/)].
57. Ma G, Al-Shabraway M, Johnson JA, Datar R, Tawfik HE, Guo D, et al. Protection against myocardial ischemia/reperfusion injury by short-term diabetes: enhancement of VEGF formation, capillary density, and activation of cell survival signaling. *Naunyn Schmiedeberg Arch Pharmacol.* 2006;**373**(6):415-27. doi: [10.1007/s00210-006-0102-1](https://doi.org/10.1007/s00210-006-0102-1). [PubMed: [16955284](https://pubmed.ncbi.nlm.nih.gov/16955284/)].
58. Tschöpe C, Walther T, Escher F, Spillmann F, Du J, Altmann C, et al. Transgenic activation of the kallikrein-kinin system inhibits intramyocardial inflammation, endothelial dysfunction and oxidative stress in experimental diabetic cardiomyopathy. *Faseb J.* 2005;**19**(14):2057-9. doi: [10.1096/fj.05-4095fj](https://doi.org/10.1096/fj.05-4095fj). [PubMed: [16129698](https://pubmed.ncbi.nlm.nih.gov/16129698/)].

59. Mihm MJ, Seifert JL, Coyle CM, Bauer JA. Diabetes related cardiomyopathy time dependent echocardiographic evaluation in an experimental rat model. *Life Sci.* 2001;**69**(5):527-42. [PubMed: [11510948](#)].
60. Jarrett RJ. Cardiovascular disease and hypertension in diabetes mellitus. *Diabetes Metab Rev.* 1989;**5**(7):547-58. [PubMed: [2689118](#)].
61. Frustaci A, Kajstura J, Chimenti C, Jakoniuk I, Leri A, Maseri A, et al. Myocardial cell death in human diabetes. *Circ Res.* 2000;**87**(12):1123-32. [PubMed: [11110769](#)].
62. Schnackenberg CG, Welch WJ, Wilcox CS. Normalization of blood pressure and renal vascular resistance in SHR with a membrane-permeable superoxide dismutase mimetic: role of nitric oxide. *Hypertension.* 1998;**32**(1):59-64. [PubMed: [9674638](#)].
63. Zhao W, Zhao T, Chen Y, Ahokas RA, Sun Y. Oxidative stress mediates cardiac fibrosis by enhancing transforming growth factor-beta1 in hypertensive rats. *Mol Cell Biochem.* 2008;**317**(1-2):43-50. doi: [10.1007/s11010-008-9803-8](#). [PubMed: [18581202](#)].