

Research Paper: Comparison of Perceived Musculoskeletal Discomfort Among Six Common Postures for Laptop Use in Female Students



Nahid Allafan^{1*}, Shahnaz Shahrbanian², Reza Rajabi¹, Hooman Minoonejad¹, Mahdi Bayati³

1. Department of Sports Medicine and Health, Faculty of Physical Education and Sport Sciences, University of Tehran, Tehran, Iran.

2. Department of Sport Science, Faculty of Humanities, Tarbiat Modares University, Tehran, Iran.

3. Department of Exercise Physiology, Sports Medicine Research Center, Sport Sciences Research Institute, Tehran, Iran.



Citation Allafan N, Shahrbanian Sh, Rajabi R, Minoonejad H, Bayati M. Comparison of Perceived Musculoskeletal Discomfort Among Six Common Postures for Laptop Use in Female Students. *Physical Treatments*. 2018; 8(1):37-44. <http://dx.doi.org/10.32598/ptj.8.1.37>

<http://dx.doi.org/10.32598/ptj.8.1.37>



Article info:

Received: 26 Oct 2017

Accepted: 05 Feb 2018

Available Online: 01 Apr 2018

Keywords:

Musculoskeletal abnormalities, Students, Female, Posture, Computers

ABSTRACT

Purpose: The present study aimed to compare Musculoskeletal Discomforts (MSDs) among six different common postures while working with laptop in female students of University of Tehran, Tehran City, Iran.

Methods: This was a crossover trial study. Eighteen female students voluntarily and purposefully participated in all stages of the study. The study participants were randomly assigned into groups to work for 10 minutes on different postures for laptop use during six continuous nights. MSDs was measured each night after 10 minutes of laptop use. For this purpose, Van der Grinten and Smith (1992) method was applied. The obtained data were analyzed by repeated measures one-way ANOVA at a significance level of $P \leq 0.05$, in SPSS.

Results: The obtained results suggested a significant difference among six working postures in MSDs ($P=0.005$). The results of Bonferroni post hoc analysis revealed that the highest level of MSDs was observed in a cross-legged sitting position on the floor. While, the lowest level of MSDs was found in the sitting on a chair posture. In addition, the study participants who used a desk in order to increase the height of the laptop, reported less levels of MSDs than laptop use in the cross-legged position on the floor ($P=0.039$) or sitting on the bed ($P=0.011$).

Conclusion: According to this study, in order to minimize MSDs during working with laptop, it is recommended to use desk and chairs instead of sitting cross-legged on the floor or bed.

* Corresponding Author:

Nahid Allafan, PhD.

Address: Department of Sports Medicine and Health, Faculty of Physical Education and Sport Sciences, University of Tehran, Tehran, Iran.

Phone: +98 (910) 8349691

E-mail: n.allafan@ut.ac.ir

Highlights

- All six working postures of working with laptop create musculoskeletal discomforts for users.
- Sitting on a chair has the lowest level of musculoskeletal discomfort.
- Cross-legged sitting position on the floor creates the highest level of musculoskeletal discomfort.

Plain Language Summary

In recent years, people, and especially students use laptops more than desktop computers. The present study aimed to compare Musculoskeletal Discomforts (MSDs) in six different common postures female university students use while working with laptops. MSDs were assessed after six continuous nights of working on the laptop (each night for 10 min). The obtained results showed significant differences among six working postures with regard to MSDs. The highest level of MSDs was observed in a cross-legged sitting position on the floor; while, the lowest level of MSDs was found in the sitting on a chair posture. In addition, participants who used a desk in order to increase the height of the laptop, reported lower level of MSDs than when they used laptop at the cross-legged position on the floor or sitting on the bed.

1. Introduction

Musculoskeletal Disorders (MSDs) are considered among the major problems in industrialized countries [1]. According to the Bureau of Labor Statistics of the United States of America (2014), MSDs accounts for 32% of all injuries and diseases [2]. MSDs are the main cause of work absence in 40% of cases [3]. According to the United Kingdom Labor Force Survey (2015), 9.5 million working days are lost due to musculoskeletal disorders, which is equivalent to 17 days per person [2]. These disorders have serious effects on the economy, reduced productivity, and early disabilities [4-6].

MSDs are the most expensive forms of labor disability [7] and impact many people [8]. MSDs occurs following damages to the muscle, bone, nerve, tendon, tendon sheath, or synovial tendon sheath lubrication [9]. MSDs is associated with tension, fatigue, soreness, heat or tremor in the affected area [10]. Short-term discomforts can predict musculoskeletal pain in the future [10-15]. Prior research reported that discomfort in the neck and shoulder regions are significantly associated with the prevalence of tendonitis in the upper limbs within a period of 4 to 5 years. Cumulative stress in the waist has also been reported to be associated with the future development of low back pain [10].

Women are more susceptible to MSDs due to lower muscle strength, higher flexibility, and less pain thresh-

old [16]. Computer users are significantly susceptible to MSDs, too [2]. Many recent studies have evaluated the effects of computer use on people's health [8]. These studies argue that the prevalence of MSDs is increasing in computer users [17] and it is even higher than some other occupations, like manual labor [18].

In recent years, laptops have been used more than desktop computers among people, and especially students. A study on 30616 students revealed that the number of laptop users has been increased from 66% in 2006 to 88% in 2009, while during this period, the number of desktop users have been decreased from 71% to 44% [19]. According to D'Silva (2016), 46.8% of US students used laptop in 2004, which reached 90% over the 10 years [20]. Laptop is among the electronic devices which can be used in any circumstances due to its portability; thus users can have a variety of postures while working with it [21]. In addition, the special design of laptops has led to increase in the frequency of bending the neck, applied torque, tissue stress, and physical discomfort [22].

Various studies demonstrated that the odds of developing MSDs is higher in working with laptop than using desktop computers [19, 23]. In addition, the comparison of MSDs among men and women has revealed that during work with laptops, women report more discomfort than men [24]. Researchers have also reported the lower level of pain threshold [25], more working hours with laptop [24, 26], and smaller skeletal systems in women, than men [24].

On average, each student spends 21.3 hours per week and 8.8% of students spend more than 40 hours per week on the internet [19]. Among which, women spend more time working with laptops per week [26] and spend more hours in sitting position for studying and working with laptops. Therefore, it can be expected that many of them suffer from MSDs due to inappropriate postures.

Smith et al. (2009) estimated that 2.52 million students suffer from upper limbs MSDs before being hired [19]. In addition, given the importance of preventing musculoskeletal disorders, also the expanded use of laptops among students, identification of postures with the least musculoskeletal injuries will greatly help to prevent such disorders, especially for the students, as the next generation of labor force. Therefore, the current study aimed to compare the MSDs in six frequent postures of using laptops for 10 minutes among female students of University of Tehran, Tehran, Iran.

2. Materials and Methods

This was a crossover trial study. The participants were randomly assigned into the groups of different postures for laptop use within a period of six continuous nights.

Participants

The study population consisted of the female students of University of Tehran. Available population was all female students who lived in dormitories of University

of Tehran. Of them, 30 female students voluntarily and purposefully participated in this study. They met the inclusion criteria and attended the evaluation session. Students who had at least 1-year experience of working with laptop, lacked any acute and chronic pain during the study [21, 27, 28], and were right handed attended the study [21, 29].

At first, the aim and procedure of the research were described to the study participants and a written consent was obtained from them for participation in the study. During the research process, 12 participants were excluded from the study due to working with laptops an hour before the time of assignment to the desired posture, or they were reluctant to continue attending the study. Eventually, 18 students participated in all stages of the study. The study participants were not allowed to use laptop accessories like cool pad and mouse during the testing process.

Data collection

The musculoskeletal discomfort was defined as any burning sensation, fatigue, stiffness, heat, or vibration in the affected area. Musculoskeletal discomfort was measured by Localized Musculoskeletal Discomfort (LMD). LMD is usually used for evaluating the severity and number of affected areas of musculoskeletal discomfort in most relevant studies [6, 19, 21, 30]. The LMD method was introduced by Van der Grinten and Smitt in



- 10= Extreme discomfort (almost maximum)
- 9=
- 8=
- 7= Very high discomfort
- 6=
- 5= High discomfort
- 4= Somewhat high discomfort
- 3= Moderate discomfort
- 2= Little discomfort
- 1= Very little discomfort
- 1/2= Extremely little discomfort
- 0= No discomfort

Figure 1. Body regions of discomfort

1992 [10]. The basis of this approach is the Berg Rating of Perceived Exertion (RPE) scale.

Participants should report any type of discomfort by selecting a number from 0 to 10 which represents the level of perceived musculoskeletal discomfort. Zero indicates lack of perceived discomfort, and 10 represents the highest level of discomfort. Participants were asked to identify the severity of discomfort in 13 areas of the body (Figure 1). The sum of 13 scales were considered as the total score for the musculoskeletal discomfort in the entire body.

Intervention

The researcher selected six frequent postures of working with laptops after interviewing 300 female students residing in the dormitory of University of Tehran and

asking about their common postures while working with laptop (Figure 2). Then, six tasks similar to working with laptop including a combination of clicking, dragging, reading and typing were designed by the researcher. The participants were randomly assigned into the groups of common postures of laptop use for six consecutive nights; 10 minutes per night [28]. More than that time, users may adjust their postures because of the limited condition. Also, in similar research studies, users reported discomfort after 10 minutes of working with computers or laptops [28, 31].

A map of human body was provided to each participant immediately after completing work with laptops, (Figure 1). Participants must first determine the letter representing the segment in which they felt discomfort from the 13 regions of human body map. Then, rate their level of musculoskeletal discomfort from 0-10. The designed



P1: (Posture) person sat on the floor. The knee was extended and laptop on the femur.



P2: person sat on the floor. The knee was extended and laptop on the pillow.



P3: person sat cross-legged on the floor and laptop on the floor.



P4: person sat cross-legged on the bed laptop on the chair.



P5: person sat on the chair. One knee flexed and laptop on the table.



P6: person sat on the chair and laptop on the table.

Figure 2. Six different common postures of students when using laptop

computer task consisted of a collection of tasks including typing, dragging, dropping, and drawing shapes which were identical for everyone.

Statistical analysis

Descriptive statistics were used to describe the participants. Shapiro-Wilk test of normality was used to examine the normal distribution of data. To determine differences among different postures, repeated measures Analysis of Variance (ANOVA) and Bonferroni post hoc test were used. All statistical methods were performed using SPSS at the significance level of $P \leq 0.05$.

3. Results

The demographic characteristics of the study participants are presented in Table 1. The obtained results suggested that the highest musculoskeletal discomfort was observed in the crossed leg posture on the floor. Also, the lowest discomfort was observed in the use of table while working with laptops. There was a significant difference between musculoskeletal disorder in six postures ($P=0.005$). Also, the results of Bonferroni post hoc test revealed a significant difference between P5 and P3 ($P=0.039$), as well as between P5 and P4 ($P=0.011$) postures (Table 2).

Table 1. Descriptive characteristics of the participants (Mean±SD)

| Variable | N=18 |
|------------------------------------|-------------|
| Age, y | 26.90±3.03 |
| Sanding height, cm | 164.16±4.79 |
| Body mass, kg | 55.26±7.21 |
| Body mass index, kg/m ² | 20.48±2.36 |

PHYSICAL TREATMENTS

Table 2. Results of inferential statistic

| Postures | Mean±SD | 95% CI | | P of ANOVA | Post Hoc ANOVA P |
|----------------|-------------|-------------|-------------|-----------------------------------|------------------|
| | | Lower Bound | Upper Bound | | |
| P ₁ | 15.05±10.82 | 9.68 | 20.43 | P ₁ vs. P ₂ | 1 |
| | | | | P ₁ vs. P ₃ | 1 |
| | | | | P ₁ vs. P ₄ | 1 |
| P ₂ | 12.72±10.41 | 7.55 | 17.90 | P ₁ vs. P ₅ | 0.105 |
| | | | | P ₁ vs. P ₆ | 0.107 |
| | | | | P ₂ vs. P ₃ | 1 |
| P ₃ | 18.78±15.53 | 11.05 | 26.50 | P ₂ vs. P ₄ | 1 |
| | | | | P ₂ vs. P ₅ | 1 |
| | | | | P ₂ vs. P ₆ | 1 |
| P ₄ | 15.11±8.11 | 11.08 | 19.14 | P ₂ vs. P ₅ | 1 |
| | | | | P ₂ vs. P ₆ | 1 |
| | | | | P ₃ vs. P ₄ | 1 |
| P ₅ | 9.33±7.24 | 5.73 | 12.94 | P ₃ vs. P ₅ | 0.039† |
| | | | | P ₃ vs. P ₆ | 0.076 |
| | | | | P ₄ vs. P ₅ | 0.011† |
| P ₆ | 9.77±9.61 | 5 | 14.55 | P ₅ vs. P ₆ | 1 |

PHYSICAL TREATMENTS

* Significantly different between the postures; † Significantly different compared with P₅ group ($P \leq 0.05$).

4. Discussion

The present study aimed to compare musculoskeletal discomfort among six common postures for laptop use after 10 minutes of using laptops in female students of University of Tehran. Results indicated that the differences in the perceived musculoskeletal discomfort between the positions that participants who used a table and a chair (P5) was significantly less than sitting in a cross legged position on the floor or bed (P3 and P4).

The obtained results of this study are consistent with the results obtained by Gold et al. Asundi et al. Moffet et al. and Fogleman et al. [21, 32-34]. The aforementioned investigations indicate that the perceived musculoskeletal discomfort has increased with decreased laptop height below the line of sight. Gold et al. (2012) evaluated kinematics and discomfort in 3 postures while working with laptop. They concluded that discomfort and non-neutral shoulders, elbows, wrists, and neck in the prone posture were more frequently reported than the posture of sitting on the couch in different states [21].

Fogleman et al. investigated the association between the screen height and keyboard dimensions on perceived discomfort in six areas of the body. They found that the lower height of the screen (below the line of sight) and the keyboard (below-elbow) was associated with increased perceived discomfort in the head/eye and shoulders/lower back, respectively [32].

Two important features of the laptops include the desktop screen connected to the keyboard and the effect of angle between the keyboard and screen on the screen light. This matter limits adjusting the position of keyboard, and screen that are the main determinants of the posture [27]. In the present study, by decreasing the laptop height (in the crossed-legs position) the study participants could increase the angle between the keyboard and screen, in order to increase the dominance over the laptop screen.

According to Villanuevaimbg et al. it requires the adoption of a forward head posture leading to increased stress on the extensor muscles of the neck [35]. In addition, in the postures that the height of the laptop is low, the keyboard is also lower in height than the elbow level. Thus, according to Simoneau et al. it leads to an increase in the extension of user's wrist [36]. Mene'ndez et al. stated that over 50% of the reported pain by students is due to working with computer [19]. Moreover, according to Chang et al. the use of more than 3 hours of computer per day doubles the odds of developing MSDs [19]. As

a result, any effective factor in preventing MSDs is important for health.

Some of the limitations of this study were the small sample size and the limited time of conducting the intervention. Therefore, longitudinal studies with larger sample sizes are required to attain more precise conclusions.

The present study compared six common postures of working with laptops after 10 minutes of using laptops among female students of University of Tehran. The obtained results recommend laptop use in a sitting position on the chair if possible. This will reduce the musculoskeletal discomfort during working with laptops. We also suggest placing and using the laptop on a table, instead of sitting on the floor or bed.

Ethical Considerations

Compliance with ethical guidelines

The study was approved by the Ethics Committee of the Sport Sciences Research Institute of Iran (IR.SSRI. REC.1397.227) and was in accordance with the Declaration of Helsinki.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authors contributions

Conceptualization and Methodology: Nahid Allafan, Reza Rajabi, Mahdi Bayati; Investigation: Nahid Allafan; Writing-Original Draft: Nahid Allafan, Mahdi Bayati, Shahnaz Shahrbanian; Writing-Review & Editing: All authors; Funding Acquisition: Mahdi Bayati; Resources: All authors; Supervision: Reza Rajabi, Shahnaz Shahrbanian.

Conflict of interest

The authors declared no conflict of interest.

References

- [1] Batham C, Yasobant S. A risk assessment study on work-related musculoskeletal disorders among dentists in Bhopal, India. *Indian Journal of Dental Research: Official Publication*

- of Indian Society for Dental Research. 2016; 27(3):236-41. [DOI:10.4103/0970-9290.186243] [PMID]
- [2] Ardahan M, Simsek H. Analyzing musculoskeletal system discomforts and risk factors in computer-using office workers. *Pakistan Journal of Medical Sciences*. 2016; 32(6):1425-9. [DOI:10.12669/pjms.326.11436] [PMID] [PMCID]
- [3] Park JH, Park JH. Association among work-related musculoskeletal disorders, job stress, and job attitude of occupational therapists. *Occupational Therapy in Health Care*. 2017; 31(1):34-43. [DOI:10.1080/07380577.2016.1270482] [PMID]
- [4] Andersen JH, Kaergaard A, Frost P, Thomsen JF, Bonde JP, Fallentin N, et al. Physical, psychosocial, and individual risk factors for neck/shoulder pain with pressure tenderness in the muscles among workers performing monotonous, repetitive work. *Spine*. 2002; 27(6):660-7. [DOI:10.1097/00007632-200203150-00017]
- [5] Buckle PW, Devereux JJ. The nature of work-related neck and upper limb musculoskeletal disorders. *Applied Ergonomics*. 2002; 33(3):207-17. [DOI:10.1016/S0003-6870(02)00014-5]
- [6] Davis KG, Kotowski SE. Postural variability: An effective way to reduce musculoskeletal discomfort in office work. *Human Factors*. 2014; 56(7):1249-61. [DOI:10.1177/0018720814528003] [PMID]
- [7] da Costa BR, Vieira ER. Risk factors for work-related musculoskeletal disorders: A systematic review of recent longitudinal studies. *American Journal of Industrial Medicine*. 2010; 53(3):285-323. [PMID]
- [8] Xie Y, Szeto G, Dai J. Prevalence and risk factors associated with musculoskeletal complaints among users of mobile handheld devices: A systematic review. *Applied Ergonomics*. 2017; 59(Pt A):132-42.
- [9] Sethi J, Sandhu JS, Imbanathan V. Effect of body mass index on work related musculoskeletal discomfort and occupational stress of computer workers in a developed ergonomic setup. *Sports Medicine, Arthroscopy, Rehabilitation, Therapy & Technology: SMARTT*. 2011; 3(1):22. [DOI:10.1186/1758-2555-3-22] [PMID] [PMCID]
- [10] Hamberg-van Reenen HH, van der Beek AJ, Blatter BM, van der Grinten MP, van Mechelen W, Bongers PM. Does musculoskeletal discomfort at work predict future musculoskeletal pain. *Ergonomics*. 2008; 51(5):637-48. [DOI:10.1080/00140130701743433] [PMID]
- [11] Genaidy AM, Delgado E, Bustos T. Active micro-break effects on musculoskeletal comfort ratings in meatpacking plants. *Ergonomics*. 1995; 38(2):326-36. [DOI:10.1080/00140139508925107] [PMID]
- [12] Galinsky TL, Swanson NG, Sauter SL, Hurrell JJ, Schleifer LM. A field study of supplementary rest breaks for data-entry operators. *Ergonomics*. 2000; 43(5):622-38. [DOI:10.1080/001401300184297] [PMID]
- [13] McLean L, Tingley M, Scott RN, Rickards J. Computer terminal work and the benefit of microbreaks. *Applied Ergonomics*. 2001; 32(3):225-37. [DOI:10.1016/S0003-6870(00)00071-5]
- [14] Fenety A, Walker JM. Short-term effects of workstation exercises on musculoskeletal discomfort and postural changes in seated video display unit workers. *Physical Therapy*. 2002; 82(6):578-89. [PMID]
- [15] Wahlstrom J. *Ergonomics, musculoskeletal disorders and computer work*. Occupational Medicine (Oxford, England). 2005; 55(3):168-76. [DOI:10.1093/occmed/kqi083] [PMID]
- [16] Hamzat TK, Abdulkareem TA, Akinyinka OO, Fatoye FA. Backpack-related musculoskeletal symptoms among Nigerian secondary school students. *Rheumatology International*. 2014; 34(9):1267-73. [DOI:10.1007/s00296-014-2962-x] [PMID]
- [17] Shabbir M, Rashid S, Umar B, Ahmad A, Ehsan S. Frequency of neck and shoulder pain and use of adjustable computer workstation among bankers. *Pakistan Journal of Medical Sciences*. 2016; 32(2):423-6. [PMID] [PMCID]
- [18] Anareh Z, ZohoorAlinia Z. [Musculoskeletal disorders among computer operators: A study in one of the governmental organizations in Kerman city, Iran (Persian)]. *Journal of Health and Development*. 2016; 5(3):216-25.
- [19] Jacobs K, Foley G, Punnett L, Hall V, Gore R, Brownson E, et al. University students' notebook computer use: Lessons learned using e-diaries to report musculoskeletal discomfort. *Ergonomics*. 2011; 54(2):206-19. [DOI:10.1080/00140130.2010.544764] [PMID]
- [20] D'Silva C. The ergonomic evaluation of laptop use in university students: The development and test-retest reliability of the student laptop use and musculoskeletal posture questionnaire [PhD. dissertation]. Oshawa: University of Ontario Institute of Technology; 2016.
- [21] Gold JE, Driban JB, Yingling VR, Komaroff E. Characterization of posture and comfort in laptop users in non-desk settings. *Applied Ergonomics*. 2012; 43(2):392-9. [DOI:10.1016/j.apergo.2011.06.014] [PMID]
- [22] Rafiee M, Mokhtarinia HR, Hadad O, Reza Soltani P. [Pain and discomfort in laptop users: Prevalence and its relation to adopted posture (Persian)]. *Razi Journal of Medical Sciences*. 2014; 21(122):37-45.
- [23] Straker L, Jones KJ, Miller J. A comparison of the postures assumed when using laptop computers and desktop computers. *Applied Ergonomics*. 1997; 28(4):263-8. [DOI:10.1016/S0003-6870(96)00073-7]
- [24] Bubric K, Hedge A. Differential patterns of laptop use and associated musculoskeletal discomfort in male and female college students. *Work (Reading, Mass)*. 2016; 55(3):663-71. [DOI:10.3233/WOR-162419]
- [25] Mingels S, Dankaerts W, van Etten L, Thijs H, Granitzer M. Comparative analysis of head-tilt and forward head position during laptop use between females with postural induced headache and healthy controls. *Journal of Bodywork and Movement Therapies*. 2016; 20(3):533-41. [DOI:10.1016/j.jbmt.2015.11.015] [PMID]
- [26] Hough R, Nel M. Time and space dimensions of computer laptop use amongst third year students of the University of the Free State. *South African Journal of Occupational Therapy*. 2016; 46(1):27-32. [DOI:10.17159/2310-3833/2016/v46n1a7]
- [27] Jonai H, Villanueva MGB, Takata A, Sotoyama M, Saito S. Effects of the liquid crystal display tilt angle of a notebook computer on posture, muscle activities and somatic complaints. *International Journal of Industrial Ergonomics*. 2002; 29(4):219-29. [DOI:10.1016/S0169-8141(01)00065-8]

- [28] Asundi K, Odell D, Luce A, Dennerlein JT. Changes in posture through the use of simple inclines with notebook computers placed on a standard desk. *Applied Ergonomics*. 2012; 43(2):400-7. [DOI:10.1016/j.apergo.2011.06.013] [PMID]
- [29] Straker L, Pollock C, Burgess-Limerick R, Skoss R, Coleman J. The impact of computer display height and desk design on muscle activity during information technology work by young adults. *Journal of Electromyography and Kinesiology: Official Journal of the International Society of Electrophysiological Kinesiology*. 2008; 18(4):606-17. [DOI:10.1016/j.jelekin.2006.09.015] [PMID]
- [30] Sommerich CM, Starr H, Smith CA, Shivers C. Effects of notebook computer configuration and task on user biomechanics, productivity, and comfort. *International Journal of Industrial Ergonomics*. 2002; 30(1):7-31. [DOI:10.1016/S0169-8141(02)00075-6]
- [31] Lin MY, Barbir A, Dennerlein JT. Evaluating biomechanics of user-selected sitting and standing computer workstation. *Applied Ergonomics*. 2017; 65:382-88. [DOI:10.1016/j.apergo.2017.04.006] [PMID]
- [32] Fogleman M, Lewis RJ. Factors associated with self-reported musculoskeletal discomfort in Video Display Terminal (VDT) users. *International Journal of Industrial Ergonomics*. 2002; 29(6):311-8. [DOI:10.1016/S0169-8141(01)00071-3]
- [33] Moffet H, Hagberg M, Hansson-Risberg E, Karlqvist L. Influence of laptop computer design and working position on physical exposure variables. *Clinical Biomechanics (Bristol, Avon)*. 2002; 17(5):368-75. [DOI:10.1016/S0021-9290(02)00062-3]
- [34] Asundi K, Odell D, Luce A, Dennerlein JT. Notebook computer use on a desk, lap and lap support: Effects on posture, performance and comfort. *Ergonomics*. 2010; 53(1):74-82. [DOI:10.1080/00140130903389043] [PMID]
- [35] Villanueva MB, Jonai H, Sotoyama M, Hisanaga N, Takeuchi Y, Saito S. Sitting posture and neck and shoulder muscle activities at different screen height settings of the visual display terminal. *Industrial Health*. 1997; 35(3):330-6. [DOI:10.2486/indhealth.35.330] [PMID]
- [36] Simoneau GG, Marklin RW. Effect of computer keyboard slope and height on wrist extension angle. *Human Factors*. 2001; 43(2):287-98. [DOI:10.1518/001872001775900940] [PMID]