

Optimizing Growth Conditions of *Kluyveromyces marxianus* for Mannan Production as a Bioemulsifier

Ashraf Hajhosseini¹, Delaram Doroud^{*2}, Anoosheh Sharifan¹, Zohreh Eftekhari³

1-Department of Food Science and Technology, Faculty of Agricultural Sciences and Food Technology, Science and Research Branch, Islamic Azad University, Tehran, Iran.

2- Pasteur Institute, Tehran, Iran.

3- Department of quality control, Pasteur Institute, Tehran. Iran.

Abstract

Background and objective: Mannan which is a linear glycoprotein with β -1,4 links carrying mannose units bind to proteins, includes natural amphiphiles and serves as a bioemulsifier. The aim of this study was optimization of growth and purification of *Kluyveromyces marxianus* for mannan production, which can use as a natural bioemulsifier.

Material and methods: In this study, mannan production by *Kluyveromyces marxianus* was assessed using combinational method of fractional factorial design and response surface methodology optimization. Process variables include concentration of carbon source (15, 30, 45 g l⁻¹) of glucose, and glycerol and methanol at 0, 2.5 and 5 g l⁻¹), nitrogen source (yeast extract and peptone 4, 6 and 8 g l⁻¹), as well as fermentation time (48, 96 and 144 h), pH (4, 6, 8) and agitation speed (150, 200 and 250 rpm).

Results and conclusion: Results showed that four variables of carbon and nitrogen source concentrations, as well as fermentation time and pH included the greatest effects on mannan production. Optimization of the affecting factors using response surface methodology demonstrated appropriate conditions of mannan production by *Kluyveromyces marxianus* as 55.15 g l⁻¹ of glucose, 9.35 g l⁻¹ of yeast extract, pH of 4.99 and fermentation time of 168 h, which led to a mannan yield of 245.98 mg (100 ml)⁻¹ culture media.

Conflict of interest: The authors declare no conflict of interest.

Article Information

Article history:

Received 3 Dec 2019
Revised 17 Dec 2019
Accepted 26 Jan 2020

Keywords:

- Bioemulsifier
- *Kluyveromyces marxianus*
- Mannan
- Purification
- Production

*Corresponding author:

Delaram Doroud,
Production and Research
Complex, Pasteur Institute of
Iran, Tehran, Iran.

Tel: +98 (0263) 6100967
E-mail:
d_doroud@pasteur.ac.ir

How to cite this article

Hajhosseini A, Doroud D, Sharifan A, Eftekhari Z. Optimizing Growth Conditions of *Kluyveromyces marxianus* for Mannan Production as a Bioemulsifier. *Appl Food Biotechnol* 2020; 7(2): 115-126. <http://dx.doi.org/10.22037/afb.v7i2.28055>

1. Introduction

Prebiotics are non-digestible ingredients that help their hosts improve their health by increasing growth and activity of the colon bacteria [1]. Health effects associated with prebiotic consumption decrease acute gastroenteritis and cancer risk and increase mineral absorption and lipid regulation [2]. Mannan, with β -1,4 glucan, is a useful prebiotic on yeast cell surface, acting as a high-affinity ligand. Moreover, the compound is able to make binding sites [3,4].

Bioemulsifiers are active surface compounds with biological origins that can be extracellular or attached to the cell surface. This compounds are produced by bacteria

yeasts and fungi. Regarding synthetic compounds, bioemulsifiers are important due to their natural origins, high biodegradability and benefits for human health. However, some bioemulsifiers described in bacteria include harmful properties that pose limitation for being widely used [1-3]. Yeasts are other sources of bioemulsifiers; however, a majority of them can produce bioemulsifiers only in the presence of water insoluble compounds such as alkanes and oils. In addition, relatively low-yields and difficulties associated with bioemulsifier extraction from growth media are other restrictions [4-6]. The mannoprotein bioemulsifier is a glycoprotein with a molecular weight of 14000-15300

Da. Inside of the cell wall of *Saccharomyces (S.) cerevisiae*, mannoprotein molecules are entrapped in glucan networks and released from cell wall of the yeast by thermal treatment under pressure. This bioemulsifier is able to stabilize oil-in-water (o/w) emulsions. Studies suggest that the chemical can be used in mayonnaise sauces with critical micelle concentration (CMC); thus, decreasing use of xanthan gums.

The baker's yeast is used for the production of this bioemulsifier due to its low costs, availability and non-toxicity. The mannoprotein molecule is composed of a polypeptide chain as well as short and long mannose links. It is stable within a broad pH range of 3-11. Extraction of mannoprotein molecules from baker's yeast cell wall is feasible using two methods of thermal and enzymatic (β 1 & 3 glucanases) processes. If the protein content of mannoprotein molecules is separated by protease enzymes, the emulsifier properties of mannoprotein is lost. This bioemulsifier is active at 5% concentrations of NaCl or higher and its production is practical at industrial scales [7-9].

In the mannoprotein structure, protein shifts into a natural amphiphilic compound through binding hydrophobic groups to mannose units; thereby, enabling its widespread use in development of food, pharmaceutical and cosmetic emulsion systems [10]. Mannoprotein was initially derived from the cell wall of *S. cerevisiae*, by Cameron et al. and Moreira et al., described as an effective polymeric bioemulsifier [11,12]. They reported higher chemical yields with no difficulties in extraction of the emulsifier from culture media as mannoprotein advantages rather than other microbial bioemulsifiers. Dikit et al. extracted and purified mannoprotein from *S. cerevisiae* KA01 by autoclaving in buffer and dialyzing in distilled water, respectively [13]. Liu et al. assessed various methods for the extraction of mannan and glucans from the cell wall of *S. cerevisiae*. They found that hot-water extraction in buffer and purification in hexadecyltrimethylammonium bromide [$C_{16}H_{33}N(CH_3)_3Br$] were the optimum methods with the greatest protective effects on mannoprotein structure [14].

In the current study, a combinational method of fractional factorial design and response surface methodology optimization was used for determining affecting factors on mannan yields in culture media. Results indicated high proportionality and confidence coefficient to predict the yields of mannan under various conditions. The optimum condition for the mannan production, optimum pH and maximum time of the appropriate yeast growth and mannan production were investigated using *Kluyveromyces (K.) marxianus* at various concentrations of carbon and nitrogen sources. Model and synthetic data from this study provide basic and functional information to estimate the substrate use, product formation and operation design, especially in industrial settings.

2. Materials and methods

All chemicals were purchased from Merck, Germany.

2.1. Yeast strains and inoculum stock preparations

K. marxianus IBRC-M 30114 was provided by Iranian Biological Resource Center. After recovery, cells were stored at 4°C on yeast-peptone-dextrose agar slants. To prepare 100 ml of the culture media, peptone (2 g), glucose (2 g) and yeast extract (1 g) were mixed in a tube and then sufficient amount of double distilled water was added to the mixture.

2.2. Inoculum development

A loopful of the yeast colony was transferred into an Erlenmeyer flask containing 100 ml of the inoculum media under sterile conditions. Flask was then incubated with agitation using shaking incubator (Orbital Shaker Incubator Rs 25000, India) at 180 rpm for 24 h at 28°C [15-17].

2.3. Culture media preparation

Briefly, 1600 ml of the yeast-peptone-dextrose culture media were modified in an Erlenmeyer flask using peptone (32 g), glucose (32 g), yeast extract (16 g) and 1600 ml of double distilled water. The final solution was divided into 19 Erlenmeyer flasks. The pH of each flask was adjusted to certain values using concentrated HCl or 4.0 N NaOH solution. Then, flasks were autoclaved at 121°C for 20 min at 1.2 bar.

2.4. Manufacturing condition

Variables, including type and concentration of the carbon and nitrogen sources, pH, time, stirring speed and yeast type, were assessed according to a full factorial design (Table 1).

To prepare culture media, compounds such as KH_2PO_4 (3 g l⁻¹), $(NH_4)_2SO_4$ (4 g l⁻¹), $MgSO_4 \cdot 7H_2O$ (0.5 g l⁻¹), $CaCl_2$ (0.13 g l⁻¹), and NaCl (0.2 g l⁻¹) were used. Totally, 300 μ l of the inoculum were transferred to each flask. Culture media were incubated at 30°C for 48 and 144 h at 150-250 rpm using shaker incubator [18].

2.5. Isolation and preparation of the yeast cells

Cells were isolated using centrifugation of the culture media at various speeds for a certain duration of time. After rinsing with cool deionized water, cells were agitated vigorously using vortex mixer with glass beads in ice. After removing and washing the glass beads, suspensions were centrifuged at appropriate speeds for a certain duration of time. Precipitated cells were rinsed several times using cool deionized water [19,20].

2.6. Extraction and purification of mannan

Yeast cell precipitates from the previous step were resuspended in 20% w w⁻¹ buffer solution, (0.1 M potassium citrate and 0.02 M sodium metabisulfite, pH 7.0) and autoclaved at 121°C for 90 min at 1.2 bar. The mannoprot-

ein was washed with acetic acid (1%) in ethanol (96%) and precipitated using centrifugation at 748 ×g for 10 min. The supernatant was stored at 4°C overnight and then centrifuged at 7656 ×g for 10 min to complete the precipitation [19,21,22]. Hexadecyltrimethylammonium bromide chemical [C₁₆H₃₃N(CH₃)₃Br] was used as solvent for selective precipitation and purification of the mannan from other macromolecules such as proteins. To complete the purification process, the precipitates were dialyzed against deionized water for 48 h [23-25].

2.7. Biomass quantity assessment

Yeast cells were isolated from suspensions using centrifugation at 748 ×g for 10 min. Then, cells were collected and dried at 105°C for 5 h until reaching a constant weight [25,26].

2.8. Extraction and purification yields

The compound yields were calculated as percent of the dry weight of the extracted mannan in 100 ml of the yeast culture media [27-29].

2.9. Experimental design and data analysis

Concentration level and effect range of each variable were assessed as pretreatments using single factor at a time. To select the most important variables, fractional factorial design was used with eight treatments (16 treatments and three center points) (Table 1). Optimization of the significant variables was carried out using response surface methodology based on the central composite design and

Minitab Software v.17 or design of experiments. Microsoft Excel v.16 was used to draw charts.

3. Results and discussion

3.1. Selection of the carbon source

First, various carbon sources (glycerol and methanol) were assessed based on the highest yields of mannan and cell biomass in a limiting solution of saturated glucose (Table 2). Data analysis showed a significant difference in carbon sources. The maximum quantity of mannan produced in the presence of glucose by *K. marxianus* included 205±4.4 mg in 100 ml of culture media with a significant difference, compared to other sources. The mannan yields in the presence of carbon sources (methanol and glycerol) were significantly lower than that in presence of others (P<0.05). Use of glucose as carbon source was reported as the best choice of carbon source (P<0.05) for mannan production.

3.2. Selection of the nitrogen source

A variety of nitrogen sources (organic) were first assessed based on the highest yields of mannan and cell biomass in solutions of carbon sources (Table 3). A significant difference was observed between various sources. The maximum quantity of mannan produced in the presence of yeast extract by *K. marxianus* included 142 ± 3.3 mg per 100 ml. Effects of various nitrogen sources on mannan yields were as follows: the yield of the production when utilizing extract was statistically more than extract and peptone together and peptone alone.

Table 1. Variable levels presented by real data or codes in fractional factorial design

Level variables	Carbon Sources			Nitrogen Sources		Environmental condition		
	Glucose	Glycerol	Methanol	Yeast Extract	Peptone	pH	Time	Agitation Speed
Coded	g l ⁻¹	g l ⁻¹	g l ⁻¹	g l ⁻¹	g l ⁻¹		h	rpm
1	15	0	0	4	4	4	48	150
0	30	2.5	2.5	6	6	6	96	200
-1	45	5	5	8	8	8	144	250

Table 2. Effects of adding different carbon sources on the yields of mannan and biomass of *K. marxianus*

Nitrogen resources	Mannan mg (100 ml) ⁻¹	Biomass (g l ⁻¹)
Glycerol	131±2.3 ^{a*}	12.24±0.15
Methanol	45±3.5 ^b	3.04±0.20
Glycerol and methanol	91±1.6 ^c	8.36±0.11
Glucose	205±4.4 ^d	17.95±0.30

*Results are statistically significant (P<0.05) in each column with non-similar alphabets

Table 3. Effects of adding various sources of nitrogen to culture media on the yields of mannan and biomass of *K. marxianus*

Nitrogen resources	Mannan mg (100 ml) ⁻¹	Biomass (g l ⁻¹)
Yeast Extract	142±3.3 ^{a*}	12.88±0.21
Peptone	112±4.6 ^{bc}	10.74±0.37
Yeast Extract+Peptone	129±2.5 ^b	11.55±0.30

*Results are statistically significant (P<0.05) in each column with non-similar alphabets

Similar findings were reported by Liu et al. for *S. cerevisiae*. They achieved the highest yields when organic nitrogen sources such as soy peptone and yeast extract were available. This revealed that inorganic nitrogen sources included no significant effects on increasing the mannan yields [24]. similar to the results by Demirci and Pometto, organic nitrogen sources enable yeasts to increase cell biomass and cell wall thickness due to easy transfer and adsorption of these sources [30]. Similar to the present results for *K. marxianus*, Mohammadzadeh et al. demonstrated that presence of organic nitrogen sources such as peptone and yeast extract led to the highest yields [31].

3.3. Screening of the process parameters

The major aim of this assessment was to select the most important variables affecting mannan production using fractional factorial design under specific conditions. Analysis of variance demonstrated that the first-order model was appropriate to fit the main effects of the six variables. Estimates of the coefficients and associated P-values within 95% confidence interval (CI) are summarized in Table 4. Mannan production and cell biomass were significantly affected by the concentration of carbon source (glucose), concentration of yeast extract, time and pH, respectively. Glycerol (as an enzyme activating agent), peptone and mixing speed included positive effects on mannan

production. Although their coefficients were as important as other factors, the fixed upper limits were considered for the optimization. Bzducha-Wrobel et al. reported positive effects of these factors as accelerators of the mannan biosynthesis in the cell wall structure of *S. cerevisiae*. Effects of methanol addition on mannan production and cell biomass seemed not significant. Thus, concentration of the carbon sources, fermentation time and pH were used as the major variables for the optimization using response surface methodology [32].

3.4. Optimization of the mannan production

To show possible changes in mannan yields and assess effects of the independent variables, including carbon and nitrogen sources, pH and fermentation time, the most appropriate model should be used to fit data. As for the prediction of yields, linear and second-order polynomial models were fitted to data (Table 5). Based on R^2 (0.996), modified R^2 (0.994) and non-significant results from lack of fit test, the second-order polynomial model was chosen. The fitted second order model was statistically significant ($P < 0.05$) and lack of fit test was non-significant for all attributes at 95% level (after model fitting, the resultant relation was subjected to the backward algorithm to reduce non-significant terms in the model).

Table 4. Estimated effects of the major variables on mannan production resulted from fractional factorial design

Source	DF	Adj SS	Adj MS	F-value	P-value	Effect	Coefficient	SE Coefficient
Model	16	61604.6	3850.3	1898.16	0.001			
Linear	8	35854.3	4481.8	2209.48	0.000		102.361	0.356
Glucose	1	3333.2	3333.2	1643.26	0.001	-28.867	-14.434	0.356
Glycerol	1	155.3	155.3	76.57	0.013	6.231	3.116	0.356
Methanol	1	0.0	0.0	0.01	0.921	0.080	0.040	0.356
Yeast Extract	1	669.9	669.9	330.25	0.003	12.941	6.471	0.356
Peptone	1	300.6	300.6	148.22	0.007	8.670	4.335	0.356
pH	1	3259.8	3259.8	1607.03	0.001	-28.547	-14.274	0.356
Time	1	26110.0	26110.0	12871.98	0.000	80.793	40.396	0.356
Agitation	1	2025.5	2025.5	998.56	0.001	22.503	11.251	0.356
2-Way Interactions	7	3019.7	431.4	212.67	0.005			
Glucose*Glycerol	1	400.8	400.8	197.61	0.005	-10.010	-5.005	0.356
Glucose*Methanol	1	19.3	19.3	9.50	0.091	-2.195	-1.098	0.356
Glucose*Yeast Extract	1	107.6	107.6	53.02	0.018	-5.185	-2.593	0.356
Glucose*Peptone	1	12.2	12.2	6.04	0.133	1.750	0.875	0.356
Glucose*pH	1	255.4	255.4	125.93	0.008	7.991	3.996	0.356
Glucose*Time	1	1277.9	1277.9	629.98	0.002	-17.874	-8.937	0.356
Glucose*Agitation	1	946.4	946.4	466.59	0.002	-15.382	-7.691	0.356
Curvature	1	22730.6	22730.6	11205.97	0.000			
Error	2	4.1	2.0					
Total	18	61608.6						
		S	R-sq	R-sq (adj)				
		1.42423	99.99%	99.94%	*			

Table 5. Central composite design for the independent variables at each run (real data or codes)

Run order	Glucose (g l ⁻¹)	yeast extract (g l ⁻¹)	pH	Time (h)	Mannan mg (100 ml) ⁻¹	Biomass (g l ⁻¹)
1	45	4	4	48	68.35	6.34
2	30	6	6	96	180.44	16.99
3	45	4	8	144	187.27	17.86
4	15	8	8	144	109.20	10.04
5	15	8	4	48	74.06	7.27
6	15	4	8	48	58.88	5.66
7	15	4	4	144	90.37	8.39
8	30	6	6	96	179.25	16.85
9	45	8	4	144	224.80	18.90
10	45	8	8	48	111.27	10.60
11	15	8	8	48	64.55	6.00
12	45	4	8	48	81.65	7.73
13	15	8	4	144	107.22	10.12
14	30	6	6	96	180.70	17.47
15	15	4	4	48	35.51	3.80
16	45	8	8	144	184.70	17.15
17	45	8	4	48	129.10	11.36
18	15	4	8	144	108.48	9.50
19	30	6	6	96	179.21	16.77
20	45	4	4	144	194.20	17.38
21	30	2	6	96	137.66	11.92
22	30	10	6	96	189.71	17.24
23	30	6	10	96	62.68	5.85
24	30	6	2	96	81.42	7.04
25	30	6	6	0	38.50	3.59
26	60	6	6	96	171.75	17.02
27	30	6	6	192	184.24	17.1
28	0	6	6	96	35.30	3.05
29	30	6	6	96	180.83	16.64
30	30	6	6	96	180.15	16.90

Table 6 describes effects of the major factors and second order effect of the factors on the mannan yields. It was found that the mutual interactions of these tested factors such as glucose concentration and yeast extract concentration, time and pH, yeast extract concentration and pH, and time and pH were significant, with carbon source concentration and time, nitrogen source concentration and pH, and nitrogen source concentration and time showing the greatest effects on response. Furthermore, R^2 and modified R^2 were relatively similar, indicating that non-significant variables were not added to the model. High R^2 and modified R^2 values as well as proportionality of these two parameters verified strength of the model to predict outcomes [33]. To determine the surface response model, linear and second-order responses and mutual interactions between the independent variables were used. The following equation demonstrated experimental relationships between the mannan yields and the tested variables within the real data:

$$\begin{aligned} \text{Mannan (mg (100 ml)}^{-1}) &= -473.5 + 5.810 X_1 + 30.03 X_2 \\ &+ 2.088 X_3 + 97.27 X_4 - 0.08383 X_1^2 - 0.955 X_2^2 \\ &- 0.007335 X_3^2 - 6.682 X_4^2 + 0.1179 X_1 X_2 + 0.01895 X_1 X_3 \\ &- 0.1781 X_1 X_4 - 0.0579 X_2 X_3 - 1.770 X_2 X_4 - 0.0236 X_3 X_4 \end{aligned}$$

Equation (1)

3.5. Effects of the carbon source concentration and fermentation time on mannan yields

Effects of the carbon source concentration and time on mannan production yield under condition; pH=6 and yeast extract concentration=6 g l⁻¹, was presented as contour plot (Figure 1).

3.6. Effects of the nitrogen source concentration and pH on mannan yields

Effects of the yeast extract concentration and pH on mannan production yield after 96 h and glucose concentration equal to 30 g l⁻¹ was presented as contour plot (Figure 2).

Increasing yeast extract concentration up to 9.65 g l⁻¹ within a pH range of 5.2 to 5.43 resulted in increasing of the mannan yields. The pH values greater than 5.43 included no positive effects on the mannan yields. Concentrations of more than 9.65 g l⁻¹ were constant. It seemed that yeasts failed to adsorb higher concentrations of the yeast extract because the carbon source was used and growth was suppressed. Low concentrations of the nitrogen source decreased protein biosynthesis and hence cell wall proteins. In contrast, high concentrations increased the growth rate but inhibited mannan production due to the complex nature of such sources. Lizicarova et al. concluded that deficient nitrogen sources decreased effects on cell growth and mannan by 14%, similar to the current findings. Therefore,

yeast extract concentrations between 10 and 11 g l⁻¹ and pH values of 5-5.5 produced the highest mannan yields [34].

Table 6. Analysis of the effect of variables and mutual interactions between independent variable on the responses

Source	DF	Adj SS	Adj MS	F-value	P-value
Model	16	97239.0	6077.4	221.40	0.000
Blocks	2	27.4	13.7	0.50	0.618
Linear	4	62413.8	15603.5	568.43	0.000
Glucose(g l ⁻¹) (X ₁)	1	27066.2	27066.2	986.02	0.000
Yeast Extract (g l ⁻¹) (X ₂)	1	3367.5	3367.5	122.68	0.000
Time (h) (X ₃)	1	31853.7	31853.7	1160.43	0.000
pH (X ₄)	1	126.5	126.5	4.61	0.050
Square	4	29781.6	7445.4	271.24	0.000
Glucose ² (g l ⁻¹) (X ₁ ²)	1	9757.1	9757.1	355.45	0.000
Yeast extract ² (g l ⁻¹) (X ₂ ²)	1	400.4	400.4	14.59	0.002
Time ² (h) (X ₃ ²)	1	7833.4	7833.4	285.37	0.000
pH ² (X ₄ ²)	1	19596.8	19596.8	713.91	0.000
2-Way Interaction	6	5016.2	836.0	30.46	0.000
Glucose(g l ⁻¹)*yeast extract: X ₁ X ₂	1	200.3	200.3	7.30	0.018
Glucose(g l ⁻¹)*Time: X ₁ X ₃	1	2979.2	2979.2	108.53	0.000
Glucose(g l ⁻¹)*pH : X ₁ X ₄	1	457.0	457.0	16.65	0.001
Yeast extract (g l ⁻¹) *Time(h): X ₂ X ₃	1	495.0	495.0	18.03	0.001
Yeast extract (g l ⁻¹)*pH : X ₂ X ₄	1	802.4	802.4	29.23	0.000
Time(h)*pH : X ₃ X ₄	1	82.2	82.2	3.00	0.107
Error	13	356.8	27.4		
Lack-of-Fit	10	354.8	35.5	51.94	0.004
Pure Error	3	2.0	0.7		
Total	29	97595.9			
		S	R-sq	R-sq (adj)	R-sq (pred)
		5.23926	99.63%	99.18%	97.08%




Figure 1. Contour plot of the effects of carbon source and time on the yields of mannan extracted from *Kluyveromyces marxianus*




Figure 2. Contour plot of the effects of nitrogen source and pH on the yields of mannan extracted from *Kluyveromyces marxianus*

3.7. Effects of the nitrogen source concentration and fermentation time on mannan yields

Effects of the yeast extract concentration and time on mannan yields under conditions; pH=6 and glucose concen

tration 30 g l⁻¹ was presented as contour plot (Figure 3).

Increased yeast extract concentrations up to 9.5 g l⁻¹ with time periods of 121 and 161 h were associated to the increased yields. Times greater than 161 h included no positive effects on yields. Concentration more than 9.5 g l⁻¹ did not change. It seemed that yeasts failed to adsorb higher concentrations of the yeast extract because the carbon source was used and growth was suppressed. Low concentrations of the nitrogen source decreased the protein biosynthesis and hence the cell wall proteins. High concentrations increased the growth rate but inhibited mannan production due to the complex nature of such sources.

3.8. Effects of the carbon source concentration and pH on mannan yields




Figure 3. Contour plot of the effects of nitrogen source and time on the yields of mannan extracted from by *Kluyveromyces marxianus*




Figure 4. Contour plot of the effects of carbon source and pH on the yields of mannan extracted from by *Kluyveromyces marxianus*

It was indicated that the mannan yields sharply decreased at low pH values. However, even increases in other factors such as glucose concentration could not increase the mannan yields, needing low rates of glucose consumption as substrate of the fermentation media. Therefore, it can be concluded that the mannan yields were high at glucose ranges of 42–43 g l⁻¹ and pH values of 5.5–5.8. Out of these ranges, increases in pH and glucose concentrations included no positive contributions to the mannan yields. Francois and Aguilar-Uscanga reported that roles of the cell wall and its

Relationships between the factors and the responses can be well understood using contour plot when considering functions of the two variables simultaneously and fixing other variables at midpoints. Effects of the carbon source concentration (e.g. glucose) and pH on mannan yields under the conditions of time of 96 h and yeast extract concentration of 6 g l⁻¹ was represented as contour plot (Figure 4). Since the linear and second-order interactions between the glucose and the pH were significant, curvatures occurred in the plots. Increases in glucose concentrations up to 42.88 g l⁻¹ led to significant increases in yields, particularly at pH values of 5.5–5.7. From this concentration to concentration of 43 g l⁻¹, yields were constant; higher concentrations included decreasing effects on the yields.

of ethanol during biomass expansion, which decreases cellular growth [37,38].

3.9. Effects of pH and fermentation time on the mannan yield

Effects of pH and time on mannan yields under the conditions of yeast extract concentration of 6 g l⁻¹ and glucose concentration of 30 g l⁻¹ was represented as contour plot (Figure 5).

Time periods up to 161 h contributed to improvements of the mannan yields. Decreases in cellular growth and mannan was seen at time periods greater than 161 h. The lower the time periods, the lower the mannan yields. Increases in other factors such as the carbon source concentration could not improve the mannan yields, suggesting low rates of glucose consumption at this condition. Time periods of 149-161 h and pH values of 5.4-6.1 demonstrated the highest mannan yields.

Time is one of the most important environmental factors, affecting growth of the microorganisms and accounting for many changes in biosynthetic and metabolic pathways. Hence, it is expected that time increases or prevents the growth of certain metabolites. Liu et al. demonstrated the maximum yields of mannan in *K. marxianus* at 96 h and 28°C. They reported that the yeast growth and activity of the mannan biosynthetic enzymes were inhibited at time periods greater than 120 h [24]. Aguilar and Francois assessed

effects of pH and temperature on the composition of cell walls, especially mannan. Increases in pH from 3 to 5 significantly increased mannan production, followed by a decrease [35]. Schultz et al. reported the maximum biomass of *K. marxianus* from whey concentrates at pH 5.6 and 33°C. Their results verified the current range of pH and constant temperature (30°C) [39].

3.10. Optimization and validation of the model

The optimal condition for the mannan production by *K. marxianus* included 55.15 g l⁻¹ of glucose concentration, 9.35 g l⁻¹ of yeast extract concentration, pH 4.99, temperature of 30°C and time of 168 h, which resulted in the maxim yield of 245.98 mg (100 ml)⁻¹. To validate the model, Erlenmeyer flask experiments at optimal points were carried out in triplicate and the experimental yields were compared with the predicted yields from the model. The mean yield was 2220 ± 18.0 mg l⁻¹. Considering the predicted yield from the model at optimal point for two yeasts [e.g. 2230 mg l⁻¹], it seemed that the predicted and experimental yields were relatively similar. Moreover, carrying out the experiments in batch fermentation under the optimum yields could reach 2459.8 mg l⁻¹. Compared to Erlenmeyer flask experiments, this improvement in mannan yields in fermentation could be attributed to the increased quantity of available oxygen, uniform distribution of nutrients and better control of pH in culture media [40,41].




Figure 5. Contour plot of the effects of pH and time on yields of mannan extracted from by *Kluyveromyces marxianus*

3.11. Kinetics of cell growth and mannan production at optimal conditions

Kinetic studies are important for optimizing biologic processes. Studying curves of growth and mannan production yeild revealed that glucose was completely consumed and yeast entered the stationary phase after 30 h. The highest quantity of mannan was achieved at the beginning of this phase (Figures 6a, b). When the carbon source was used, the yeast growth was prevented at the end of the process, showing that glucose was appropriate for the yeast and was the only limiting substrate for the microbial growth in the media. During the exponential phase, the ratio of special

growth to time was constant. This suggest that no inhibitory effects such as alcohol were present in the media [42-44].

Mannan can be extracted from several yeast strains, representing 4-13% of the cell dry weight content. Similar findings by Ozmihci and Kargi were reported on kinetic models of the *K. marxianus* growth at various concentration of the carbon source [45].

Galinari et al. and Lukondeh et al. reported kinetic characteristics of *K. marxianus* with yield of the purified bioemulsifier (7-13.3%) of the dry cell weight and concentration of the purified product (12 g l⁻¹) [29,36].

Solis-Pacheco et al. reported the kinetic characteristics of *S. cerevisiae* (CTGM, CTSA) and *Meyerozyma guilliermondii* (CT15, CT25, CT35) with 30 g l⁻¹ of sugars and 5 g l⁻¹ of (NH₄)₂SO₄, they found that the cell wall composition was more variable between the strains than the species and was highly depended on the growth phases. In the stationary phase, only two strains (CT15 and CT25) included high β-glucan and mannan contents in the cell wall (~30 mg g⁻¹) while the others yielded 3-18 mg g⁻¹ as well as control bakery yeast strain [46].

4. Conclusion

The increasing popularity of emulsifier consumptions has resulted in development of novel sources to produce bioemulsifiers. Biological approaches serve as the most effective and appropriate tools for producing these products using microorganisms.

Results of this study have represented a successful model for optimizing conditions that affect the production yields of mannan extracted from *K. marxianus*. Furthermore, data from this study have provided basic information for further investigations of mannan as a safe and effective bioemulsifier for using in food formulations and pharmaceutical products at large scales.




Figure 6a. The growth curve of *Kluyveromyces marxianus* at optimal conditions




Figure 6b. The yield curve of mannan extracted from *Kluyveromyces marxianus* under optimal conditions

Table 7. Conditions of the batch fermentation system for the process of *K. marxianus* growth during 30 h

Time	rpm	O ₂	CO ₂	pH	Do	Temp. (°C)	Biomass (g l ⁻¹)	Mannan mg (100 ml) ⁻¹
12:30	500	20.3	0.05	4.31	32.5	27.3	4.1	42.5
18:00	500	19.7	0.15	4.66	29.1	28.9	6.5	61.3
20:30	500	19.5	0.21	4.59	25.8	29.3	9.2	94.6
22:30	500	19.3	0.25	4.51	20.9	29.4	12.5	131.2
0:30	500	19.5	0.23	4.56	26.5	29.1	16.3	193.6
2:30	500	19.4	0.24	4.57	25.8	29.2	18.5	220.0
4:00	500	19.5	0.26	4.57	26.7	29.1	21.6	234.3
6:30	500	19.5	0.24	4.58	31.2	29.1	26	257.9
8:00	500	19.6	0.23	4.61	32.7	28.9	28.2	287.5
10:30	500	19.6	0.21	4.60	33.2	28.9	31.5	306.5
12:00	500	19.1	0.13	4.60	35.2	29.2	33.2	337.2
14:30	500	19.7	0.06	4.60	36.5	29.1	34	345.1

5. Acknowledgements

The authors are grateful to Pasteur Institute of Iran, Tehran, for supporting this study.

6. Conflict of interest

The authors declare no conflict of interest.

References

- Homayouni Rad A, Akbarzadeh F, Mehrabany EV. Which are more important: Prebiotics or probiotics? *Nutr.* 2012; 28 (12): 1196-1197. doi: 10.1016/j.nut.2012.03.017
- Tabrizi A, Khalili L, Homayouni-Rad A, Dehghan P, Ansari F. Prebiotics, as promising functional food to patients with psychological disorders: A review on mood disorders, sleep, and cognition. *NeuroQuantology* 2019; 17(6): 1-9. doi: 10.14704/nq.2019.17.6.2189
- Mohabbi Z, Azizi M, Hosseini SJ, Alizadeh M, Homayouni A. The effects of prebiotic bread containing oat β-glucan and resistant starch on the glycemic index and glycemic load in healthy individuals. *Nutr Food Sci.* 2019. doi: 10.1108/NFS-10-2018-0292
- Mohebbi Z, Homayouni A, Azizi MH, Hosseini SJ. Effects of beta-glucan and resistant starch on wheat dough and prebiotic bread properties. *J Food Sci Technol.* 2018; 55(1): 101-110. doi: 10.1007/s13197-017-2836-9
- Chaprao MJ, Ferreira IN, Correa PF, Rufino RD, Luna JM, Silva EJ, Sarubbo LA. Application of bacterial and yeast biosurfactants for enhanced removal and biodegradation of motor oil from contaminated sand. *Electron J Biotechnol.* 2015; 18(6): 471-479. doi: 10.1016/j.ejbt.2015.09.005

6. Mohammed MK. Isolation and screening of biosurfactant producing yeast from soil contaminated with oil. *J Wasset Sci Med*. 2018; 11(1): 161-171.
7. Alizadeh-Sani M, Hamishehkar H, Khezerlou A, Azizi-Lalabadi M, Azadi Y, Nattagh-Eshtivani E, Fasihi M, Ghavami A, Aynehchi A, Ehsani A. Bioemulsifiers derived from microorganisms: Applications in the drug and food industry. *Adv Pharm Bull*. 2018; 8(2): 191-199
doi: 10.15171/apb.2018.023
8. Jana UK, Suryawanshi RK, Prajapati BP, Soni H, Kango N. Production optimization and characterization of manno-oligosaccharide generating β -mannanase from *Aspergillus oryzae*. *Bioresource Technol*. 2018; 268:308-314.
doi: 10.1016/j.biortech.2018.07.143
9. Okoro CC, Nwezza EE, Lin J. Persistence of halophilic methanogens and oil-degrading bacteria in an offshore oil-producing facility. *Geomicrobiol J*. 2018; 35(4): 323-333.
doi: 10.1080/01490451.2017.1368742
10. Rahman PK, Mayat A, Harvey JGH, Randhawa KS, Relph LE, Armstrong MC. Biosurfactants and Bioemulsifiers from Marine Algae. *The Role of Microalgae in Wastewater Treatment*: Springer; 2019: pp. 169-188.
doi: 10.1007/978-981-13-1586-2_13
11. Morrissey JP, Etschmann MM, Schrader J, de Billerbeck GM. Cell factory applications of the yeast *Kluyveromyces marxianus* for the biotechnological production of natural flavour and fragrance molecules. *Yeast*. 2015; 32(1): 3-16.
12. Kreling NE, Zapparoli M, Cecchin I, Thome A, Reddy KR, Colla LM. Inactive yeast biomass and mannoproteins influence the retention and biodegradation of biodiesel in soil. *Env Quality Manage*. 2018; 28(2): 105-112.
doi: 10.1002/tqem.21590
13. Dikit P, Maneerat S, H-kittikun A. Mannoprotein from spent yeast obtained from thai traditional liquor distillation: extraction and characterization. *J Food Process Eng*. 2012;35(1):166-177.
doi: 10.1111/j.1745-4530.2010.00592.x
14. Wang J, Li M, Zheng F, Niu C, Liu C, Li Q, Sun G. Cell wall polysaccharides: Before and after autolysis of brewer's yeast. *World J Microbiol Biotechnol*. 2018; 34(9): 137.
doi: 10.1007/s11274-018-2508-6
15. Cameron DR, Cooper DG, Neufeld R. The mannoprotein of *Saccharomyces cerevisiae* is an effective bioemulsifier. *Appl Environ Microbiol*. 1988; 54(6): 1420-1425.
doi: 10.1128/AEM.54.6.1420-1425.1988
16. Moreira TCP, Da Silva VM, Gombert AK, Da Cunha RL. Stabilization mechanisms of oil-in-water emulsions by *Saccharomyces cerevisiae*. *Colloid Surface B*. 2016; 143: 399-405.
doi: 10.1016/j.colsurfb.2016.03.043
17. Dikit P, Maneerat S, Musikasang H, H-kittikun A. Emulsifier properties of the mannoprotein extract from yeast isolated from sugar palm wine. *Sci Asia*. 2010; 36: 312-318.
doi: 10.2306/scienceasia1513-1874.2010.36.312
18. Liu H-Z, Wang Q, He Y. Immunoactivities and antineoplastic activities of *Saccharomyces cerevisiae* mannoprotein. *Carbohydr Polym*. 2011; 83(4): 1690-1695.
doi: 10.1016/j.carbpol.2010.10.026
19. Aktas N, Boyacı İH, Mutlu M, Tanyolac A. Optimization of lactose utilization in deproteinated whey by *Kluyveromyces marxianus* using response surface methodology response surface methodology. *Bioresource Technol*. 2006; 97(18): 2252-2259.
doi: 10.1016/j.biortech.2005.10.039
20. Loser C, Urit T, Gruner E, Bley T. Efficient growth of *Kluyveromyces marxianus* biomass used as a biocatalyst in the sustainable production of ethyl acetate. *Energy Sustain Soc*. 2015; 5(1): 2-15
doi: 10.1186/s13705-014-0028-2
21. Eluk D, Ceruti R, Nagel O, Althaus R. Effect of thermal treatment of whey contaminated with antibiotics on the growth of *Kluyveromyces marxianus*. *J Dairy Res*. 2019; 86(1): 102-107.
doi: 10.1017/S0022029919000098
22. Torija MaJ, Beltran G, Novo M, Poblet M, Rozes N, Guillamon JM, Mas A. Effect of the nitrogen source on the fatty acid composition of *Saccharomyces cerevisiae*. *Food microbiol*. 2003; 20(2): 255-258.
doi: 10.1016/S0740-0020(02)00091-6
23. Hanneh VA, Ali MS, Elnaz M, Arash K, Masoome M. Statistical optimization of culture media and conditions for maximize production of mannan by *Saccharomyces cerevisiae* using response surface methodology. *Annu Res Rev Biol*. 2014: 1927-1940.
doi: 10.9734/ARRB/2014/8712
24. Liu H-Z, Wang Q, Liu Y-Y, Fang F. Statistical optimization of culture media and conditions for production of mannan by *Saccharomyces cerevisiae*. *Biotechnol Bioproc Eng*. 2009; 14(5): 577-583.
doi: 10.1007/s12257-008-0248-4
25. Da Silva Araujo VB, De Melo ANF, Costa AG, Castro-Gomez RH, Madruga MS, De Souza EL, Magnani M. Followed extraction of β -glucan and mannoprotein from spent brewer's yeast (*Saccharomyces uvarum*) and application of the obtained mannoprotein as a stabilizer in mayonnaise. *Innov Food Sci Emerg*. 2014; 23: 164-170.
doi: 10.1016/j.ifset.2013.12.013
26. Al-Manhel A, Niamah A. Mannan extract from *Saccharomyces cerevisiae* used as prebiotic in bioyogurt production from buffalo milk. *Int Food Res J*. 2017; 24(5): 2259-2264.
doi: 10.12944/CRNFSJ.5.3.19
27. Galinari E, Almeida-Lima J, Macedo GR, Mantovani HC, Rocha HAO. Antioxidant, antiproliferative, and immunostimulatory effects of cell wall α -d-mannan fractions from *Kluyveromyces marxianus*. *Int J Biol Macromolecules*. 2018; 109: 837-846.
doi: 10.1016/j.ijbiomac.2017.11.053
28. Dell Anno F, Sansone C, Ianora A, Dell Anno A. Biosurfactant-induced remediation of contaminated marine sediments: Current knowledge and future perspectives. *Mar Environ Res*. 2018; 137: 196-205.
doi: 10.1016/j.marenvres.2018.03.010
29. Galinari E, Sabry DA, Sasaki GL, Macedo GR, Passos FML, Mantovani HC, Oliveira Rocha HA. Chemical structure, antiproliferative and antioxidant activities of a cell wall α -d-mannan from yeast *Kluyveromyces marxianus*. *Carbohydr Polym*. 2017; 157: 1298-305.
doi: 10.1016/j.carbpol.2016.11.015
30. Demirci A, Pometto AL. Enhanced organically bound chromium yeast production. *J Agri Food Chem*. 2000; 48(2): 531-536.
doi: 10.1021/jf990771q

31. Mohammadzadeh J, Tabatabaei F, Mortazavi A, Kadkhodae R, Koocheki A. A study on the effect of mannoprotein extracted from *Kluyveromyces marxianus* on the quality attributes and stability of oil-in-water emulsion. *Sci Infor Database*. 2014; 4(3): 331-346.
32. Bzducha-Wrobel A, Kieliszek M, Błazejak S. Chemical composition of the cell wall of probiotic and brewer's yeast in response to cultivation medium with glycerol as a carbon source. *Eur Food Res Technol*. 2013; 237(4): 489-499. doi: 10.1007/s00217-013-2016-8
33. Campos BE, Ruivo TD, Da Silva Scapim MR, Madrona GS, Bergamasco RdC. Optimization of the mucilage extraction process from chia seeds and application in ice cream as a stabilizer and emulsifier. *LWT Food Sci Technol*. 2016; 65: 874-883. doi: 10.1016/j.lwt.2015.09.021
34. Lizicarova I, Matulova M, Capek P, Machova E. Human pathogen *Candida dubliniensis*: A cell wall mannan with a high content of α -1, 2-linked mannose residues. *Carbohydr Polym*. 2007; 70(1): 89-100. doi: 10.1016/j.carbpol.2007.03.007
35. Aguilar-Uscanga B, Francois J. A study of the yeast cell wall composition and structure in response to growth conditions and mode of cultivation. *Lett Appl Microbiol*. 2003; 37(3): 268-274. doi: 10.1046/j.1472-765X.2003.01394.x
36. Lukondeh T, Ashbolt NJ, Rogers PL. Evaluation of *Kluyveromyces marxianus* FII 510700 grown on a lactose-based medium as a source of a natural bioemulsifier. *J Ind Microbiol Biotechnol*. 2003; 30(12): 715-720. doi: 10.1007/s10295-003-0105-
37. Silva ACS, Santos PND, Andrade RFS, Campos-Takaki GM. Biosurfactant production by fungi as a sustainable alternative. *Arq Inst Biol*. 2018; 85. doi: 10.1590/1808-1657000502017
38. Mirzaeia M, Mirdamadi S, Ehsani M, Aminlari M, Hoseini S. Characterization of yeast protein enzymatic hydrolysis and autolysis in *saccharomyces cerevisiae* and *Kluyveromyces marxianus*. *J Food Biosci Technol*. 2015; 5(2): 19-30. doi: 10.1016/j.jff.2015.09.031
39. Schultz N, Chang L, Hauck A, Reuss M, Syldatk C. Microbial production of single-cell protein from deproteinized whey concentrates. *Appl Microbiol Biotechnol*. 2006; 69(5): 515-520. doi: 10.1007/s00253-005-0012-z
40. Ghaly A, Kamal M, Avery A. Influence of temperature rise on kinetic parameters during batch propagation of *Kluyveromyces fragilis* in cheese whey under ambient conditions. *World J Microbiol Biotechnol*. 2003; 19(7): 741-749. doi: 10.1023/A:1025148022934
41. Kasmir M. Biological processes as promoting way for both treatment and valorization of dairy industry effluents. *Waste Biomass Valor*. 2018; 9(2): 195-209. doi: 10.1007/s12649-016-9795-7
42. Goswami G, Chakraborty S, Chaudhuri S, Dutta D. Optimization of process parameters by response surface methodology and kinetic modeling for batch production of canthaxanthin by *Dietzia maris* NIT-D (accession number: HM151403). *Bioproc Biosyst Eng*. 2012; 35(8): 1375-1388. doi: 10.1007/s00449-012-0726-0
43. Mitra R, Chaudhuri S, Dutta D. Modelling the growth kinetics of *Kocuria marina* DAGII as a function of single and binary substrate during batch production of β -Cryptoxanthin. *Bioproc Biosyst Eng*. 2017; 40(1): 99-113.
44. Nayak M, Dhanarajan G, Dineshkumar R, Sen R. Artificial intelligence driven process optimization for cleaner production of biomass with co-valorization of wastewater and flue gas in an algal biorefinery. *J Clean Prod*. 2018; 201: 1092-2100. doi: 10.1016/j.jclepro.2018.08.048
45. Ozmihi S, Kargi F. Kinetics of batch ethanol fermentation of cheese-whey powder (CWP) solution as function of substrate and yeast concentrations. *Bioresour Technol*. 2007; 98(16): 2978-2984. doi: 10.1016/j.biortech.2006.10.005
46. Solis-Pacheco J, Peter K, Lopez Z, Gonzalez-Reynoso O, Macias-Rodriguez M, Gonzalez-Alvarez V, Lacroix M, Aguilar-Uscanga BR. Differences in the amount of β -glucan and mannan in strains of *Saccharomyces cerevisiae* and *Meyerozyma guilliermondii* isolated from agave must used in tequila production. *Microbial Res Int*. 2013; 2(1): 1-8

بهینه سازی شرایط رشد کلایورومایسس مارکسیانوس برای تولید مانان به عنوان بیومولسیفایر

اشرف حاج حسینی^۱، دلارام درود^{۲*}، انوشه شریفان^۱، زهره افتخاری^۳

۱- گروه علوم و مهندسی صنایع غذایی، دانشکده علوم کشاورزی و صنایع غذایی، واحد علوم و تحقیقات، دانشگاه آزاد اسلامی، تهران، ایران.

۲- انستیتو پاستور، تهران، ایران.

۳- گروه کنترل کیفیت، انستیتو پاستور، تهران، ایران.

چکیده

سابقه و هدف: مانان، گلیکو پروتئینی خطی با اتصالات بتا ۱-۶ متشکل از واحد های مانوز متصل به پروتئین است که دارای ساختار ترکیبات آمفی فیلیک طبیعی بوده و می تواند به عنوان بیومولسیفایر مورد استفاده قرار گیرد. هدف از این مطالعه، بهینه سازی شرایط رشد و خالص سازی کلایورومایسس مارکسیانوس برای تولید مانان است که به عنوان یک بیومولسیفایر طبیعی قابل استفاده است.

مواد و روش ها: در این مطالعه، تولید مانان توسط کلایورومایسس مارکسیانوس با استفاده از ترکیب طرح کسری از فاکتوریل و سطح پاسخ مورد بررسی قرار گرفت. متغیرهای تحقیق شامل غلظت منبع کربن (گلوکز در غلظت های ۱۵، ۳۰، ۴۵ و gl^{-1} و نیز گلیسرول و متانول در غلظت های ۰، ۲/۵ و gl^{-1} ، منبع ازت (عصاره مخمر و پپتون در غلظت های ۰، ۴، ۶ و gl^{-1})، زمان تخمیر (۹۶، ۴۸ h و ۱۴۴ h) و pH (۴، ۶ و ۸)، سرعت همزن (۱۵۰ و ۲۰۰ rpm و ۲۵۰ rpm) بود.

یافته ها و نتیجه گیری: نتایج نشان داد چهار متغیر غلظت منبع کربن و نیتروژن، زمان تخمیر و pH بیشترین تاثیر را بر تولید مانان داشتند. بهینه سازی عوامل موثر با روش سطح پاسخ، شرایط مناسب تولید مانان از کلویورومایسس مارکسیانوس را با غلظت های ۵۵/۱۵ گرم بر لیتر گلوکز، ۹/۳۵ گرم بر لیتر عصاره مخمر، pH برابر ۴/۹۹ و زمان تخمیر ۱۶۸ ساعت نشان داد که به راندمان تولید مانان به میزان $245/98 \text{ mg}(100\text{ml})^{-1}$ محیط کشت منجر شد. بنابراین نتایج این تحقیق می تواند به عنوان یک الگوی موفق در بهینه سازی عوامل مؤثر در راندمان تولید مانان در محیط کشتی از کلایورومایسس مارکسیانوس استفاده شود.

تعارض منافع: نویسندگان اعلام می کنند که هیچ نوع تعارض منافی مرتبط با انتشار این مقاله ندارند.

تاریخچه مقاله

دریافت ۳ دسامبر ۲۰۱۹

داوری ۱۷ دسامبر ۲۰۱۹

پذیرش ۲۶ ژانویه ۲۰۲۰

واژگان کلیدی

• بیومولسیفایر

• کلایورومایسس مارکسیانوس

• مانان

• تولید

*نویسنده مسئول

دلارام درود،

مجتمع تولید و تحقیق، انستیتو

پاستور، تهران، ایران.

تلفن: ۹۸(۲۶۳) ۶۱۰۰۹۶۷

پست الکترونیک:

d_doroud@pasteur.ac.ir