

Open Access



Crescent Journal of Medical and Biological Sciences Vol. 2, No. 2, Spring 2015, 64-68 elSSN: 2148-9696

Honey and Vitamin E Restore the Plasma Level of Gonadal Hormones and Improve the Fertilization Capacity in Noise-Stressed Rats

Asghar Rajabzadeh¹, Mohsen Sagha^{2*}, Mohammad Reza Gholami³, Reza Hemmati³

Abstract

Objective: Noise as a natural teratogenic factor affects the body systems including the reproductive organ to reduce the fertility rate and fetus health. Honey and vitamin E as natural antioxidants protects the sperm released from the reproductive system. This study was conducted to examine the efficacy of honey and vitamin E on fertilization capacity in noise-exposed rats by assessing plasma sexual hormones levels i.e., follicle-stimulating hormone (FSH), luteinizing hormone (LH), and testosterone, altered in relation with noise stress.

Materials and Methods: This study was targeted the 24 male rats that randomly were divided into four equal groups including one control group (unexposed to noise stress) and three experimental groups pre-induced with noise stress for 50 days and then divided as: no treated, honey and vitamin E treated groups, respectively. Then, the blood samples of experimental and control groups were taken, and the serum level of the sexual hormones was analyzed. Finally, to investigate the fertility capacity of rats, the male rats of all groups were coupled with the female ones.

Results: Our results showed that FSH and LH level in noise stressed male rats raised, and the testosterone secretion decreased compared to the control group. Moreover, noise stress injury could reduce weight and the survival rate of the fetus. However, the honey and vitamin E improved the testosterone concentration, declined the plasma FSH and LH level in noise - exposed rats and enhanced the fertility rate.

Conclusion: These findings may also spell out a natural curative approach rather than pharmaceutical drugs to optimize of neuroendocrine gonadal axis and testicular integrity induced by pathogenesis stress, i.e., noise and enhance the male fertility capacity.

Keywords: Gonadal Hormones, Honey, Noise, Vitamin E

Introduction

Stress is defined as a physiological response to the body's homeostasis disorders (1). Among environmental stresses, noise as a natural teratogenic agent is relatively more common and affects body systems including the reproductive organ to reduce the fertility rate and fetus health (2,3). It has been shown that some stresses, e.g., immobilization, heat and water restrictions alter serum sex hormones levels, resulted to decrease in the fertilization rate and implantation (4,5). Furthermore, some researchers demonstrated that exposure to noise can change the stress hormones such as corticosteroid, norepinephrine, and adrenocorticosteroid and disrupts the testis integrity

Received: 22 Dec 2014, Revised: 18 Jan 2015, Accepted: 25 Feb 2015, Available online: 15 Apr 2015

¹ Department of Anatomical Sciences, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

² Research Laboratory for Embryology and Stem Cells AND Department of Anatomical Sciences and Pathology, School of Medicine, Ardabil University of Medical Sciences, Ardabil, Iran

³ Department of Anatomical Sciences, Faculty of Medicine, Lorestan University of Medical Sciences, Khorramabad, Iran

^{*}Corresponding Author: Mohsen Sagha, Research Laboratory for Embryology and Stem Cells AND Department of Anatomical Sciences and Pathology, School of Medicine, Ardabil University of Medical Sciences, Ardabil, Iran

Tel: +98 9131257611, Email: m.sagha@arums.ac.ir

through decreasing the testosterone levels, led to reduce the number of healthy spermatozoa. Moreover, the long-term exposure to noise can highly increase the sperm sensitivity to oxidative stress and in turn affects the sperm capacity and acrosome reaction (6,7).

The testes normally have antioxidant factors, protecting their germ cells from the oxidative damage (8,9). Nevertheless, in the mature spermatozoa when the production of free radicals increases, the physiological system will not respond. Using of antioxidants in everyday life is very important, because many biological problems are caused by free radicals which induce oxidative damage to lipids and proteins in the cell membrane (10).

Natural antioxidants can prevent devastating changes in spermatozoa. Honey and vitamin E are principle antioxidants that are found in abundance in Honey as an antimicrobial nature. and anti-inflammatory food is known as a biological antioxidant and increase sperm number in rats affecting without any on the serum follicle-stimulating hormone (FSH), luteinizing hormone (LH), and testosterone (10-12).

The antioxidant property of honey is due to its phenolic components (11) and others ingredients i.e., catalase, glucose and fructose, and vitamin C and B as well as minerals including magnesium, potassium and calcium (12-14). Vitamin E is another antioxidant that is found in cell membranes and protects the cell membranes against the hydrogen peroxides (15). α -tocopherol is the natural form of vitamin E with high biological activity (16). Some reports indicate the key role of vitamin E in reducing if oxidative stress in the testis (17,18). Indeed, this vitamin has a protective effect on testicular function and increases male fertilization capacity (19).

A series of neurological, paracrine, and endocrine systems are involved in normal physiological functions of the reproductive system. However, less is known about the endocrinological response of the male reproductive organs to noise stress and also the role of the natural antioxidants in reducing stress and increasing the fertility outcome through modulation of sexual hormone level. Hence, this study aimed at evaluating honey and vitamin E effects on the sexual hormones levels and fertilization capacity in noise-exposed male rats. We found that both aforementioned components restore FSH, LH, and testosterone level to an optimum level led to improve fertilization capacity in noised-stressed rats.

Materials and Methods

A total of 24 adult male Sprague-Dawley rats weighing 200 ± 20 g were acclimated to $22 \pm 1^{\circ}$ C and maintained under conditions of 12-h periods of the light and dark with free access to tap water and commercial rat food. All procedures were approved by international guidelines and by the Institute Research Ethics and Animal Care and Use committee of Tabriz University of

Medical Sciences. Every effort was made to minimize the number of animals used and their suffering. The rats randomly divided into four groups (six rats per group) and classified as follows:

Control group that was not exposed to noise and experimental groups 1, 2, and 3 consisting no treated, honey and vitamin E treated groups, respectively. All experimental groups were exposed to 90-130 dB of noise with a frequency of 300-350 Hz daily from 7 pm to 7 am for 50 days. Then, the male rats were transported to a room with dimensions of $3 \text{ m} \times 4 \text{ m} \times 3 \text{ m}$ lagged by wood and acoustic segments (anti-loud voice). In this room, the set of white noise was prepared at 19 O'clock with frequency of 300-350 Hz and intensity of 90-120 dB by the speaker (20) and the timer had been set so that after 1 h of operation, the noise was turned off for a few minutes (from 15 to 60 min) and then re-operated. This situation was led to prevent animal's compatibility. Automatically, intensity and frequency of noise were changed in the district of minimum and maximum every 2-3 min by the noise aided in non-compatibility producing device, condition again (21). For evaluating the amount and intensity of noise, the noise level meter was used (22).

Group 2 and 3 noise-exposed rats received honey (0.2 ml of 5 % honey dissolved in water) and vitamin E (75 mg/kg/day dissolved in corn oil) by gavage with a custom-made instrument containing a tube with an approximate diameter 2 mm, once a day at 18:00 h for 50 days without delaying upon noise pollution.

After 50 days, the blood sample obtained from each rat's tail and serum FSH, LH, and testosterone level were measured using enzyme-linked immunosorbent assay (ELISA) technique (CUSABIO kit, Japan).

Both in experimental and control groups, each male rat was coupled with two females for 1 week. Then, vaginal plug-positive females were considered as being pregnant on day 0. Finally, prior to delivery, the pregnant female rats were killed with an anesthetic drug and then, weight and number of living, dead, and absorbed fetuses were calculated.

Statistical analysis

All experiments were replicated at least three independent times and data were presented as a mean \pm SD and analyzed by one-way ANOVA followed by Tukey's *post-hoc* multiple group comparison test. A difference between groups was considered as statistically significant if P < 0.050.

Results

Sexual hormone level

To investigate whether noise pollution changes sexual hormones levels, male rats were induced by noise for 50 days and our results showed that serum FSH and LH levels demonstrated to be 8.06 ± 0.9 and 7.46 ± 0.8 mIU/ml which was significantly different from control group calculated 2.69 ± 0.9 and 2.35 ± 0.3 mIU/ml, respectively (P < 0.001) (Figure 1A and B). Moreover, measurement of testosterone

level in the control group was shown to be 8.61 ± 0.33 ng/ml but noise stress could significantly decrease it to 3.25 ± 0.28 ng/ml.

Administration of vitamin E and honey diminished gonadotropin hormones levels to 2.97 ± 0.6 and 4.33 ± 0.4 mIU/ml (for FSH) and 3.39 ± 0.36 and 4.6 ± 0.35 mIU/ml (for LH), respectively compared to noise-stress group (P < 0.001) (Figure 1A and B). contrasting to serum gonadotropin hormones, the plasma testosterone secretion was significantly raised to 7.56 ± 0.55 and 5.4 ± 0.75 ng/ml in vitamin E and honey administered rats, respectively, compared to noise-stressed rats (P < 0.001) (Figure 1C).

Male's fertilization outcome

As shown in table 1, noise-exposed male rats found to generate six living and one dead fetuses weighing 3.35 ± 0.33 g upon mating with female rats compared to control group that delivered nine living fetuses weighing 4.51 ± 0.48 g. Whether honey and vitamin E can improve fertility rate in male noised stressed rats, our results showed that these antioxidants could restore fertilization potential to male rats to produce 11 and 9 living fetuses weighing 5.29 ± 0.88 g and 5.29 ± 0.31 g, respectively (Table 1).

Discussion

Normally, FSH and LH secreted from the hypophysis gland control the primary function of the testis through adjusting the spermatogenesis and testosterone production by Leydig's cells. However, some stress factors i.e., noise pollution and forced swimming stress may influence on this balance and sperm parameters (2,23-25).

In present study, we showed that the noise stress changed physiological secretion of sexual hormone in rat plasma and honey and vitamin E restored their levels to an optimum rate affected on animal fertilization potential. In accordance with the current result, Swami et al. (2) also found that the serum FSH and LH levels raised multi-folds in the male rats exposed to noise stress (100 dB) that in turn is more resulting in decreased the plasma testosterone level. In addition, the intermittent scrotal hyperthermia stress was resulted in plasma FSH and LH raising and serum testosterone decreasing (22). Therefore, it concepts that noise pollution may disrupt the steroid hormones concentration and the neuroendocrine gonadal axis resulting in cellular damage of testicular tissue caused to fall sharply secretion of testosterone serum. Furthermore, it seems that every factor that can trigger reduce or even inhibit gonadotropin and sex hormones secretion, normally can lead to induction of deleterious in germinative and somatic cells of testes parenchyma (26). Indeed, reduction in the plasma testosterone level may be back to disorganization of Leydig's cells, as stated in the previous study (12).

Some of these troubles could probably be diminished using the natural curative agents, protecting reproductive system within the stress period. The current study also pointed out that honey and vitamin E improved the serum testosterone concentration while declined the plasma FSH and LH secretion in noise-exposed rats. Some studies also confirmed this finding so that Abdul-Ghani et al. reported that honey augments the spermatogenesis process without any disturbance in steroids hormones secretion, i.e., gonadotropins and estradiol (10).



Figure 1. Sexual hormones levels in different groups; Compared to control group: (A) follicle-stimulating hormone (B) luteinizing hormone levels were increased in noise-exposed rats but honey and vitamin E decreased these gonadotropin hormones levels, (C) compared to control, noise declined significantly testosterone level which was increased again in honey and vitamin E consumed rats ** Versus noise-stressed rats, P < 0.001

Table 1. Number of living and dead fetuses and their weights in different groups

	U	<u> </u>	V	
Groups	Total fetus	Number of living fetus (%)	Number of dead fetus (%)	Fetus weight (g) (mean ± SD)
Control	9	9 (100)	-	4.51 ± 0.48
Noise	7	6 (85)	1(15)	3.35 ± 0.33
Honey	11	11 (100)	-	5.29 ± 0.88
Vitamin E	10	9 (90)	1 (10)	5.29 ± 0.31
SD: Standard deviation				

66 | Crescent J Med & Biol Sci, Vol 2, No. 2, Spring 2015

www.SID.ir

Mahanem et al. declared that if a suitable amount of honey was given to the male rats, the spermatogenic cells lineage and also rates of sperms were increased (27). Naseem et al. reported that vitamin E enhanced the sperm parameters (28). Indeed, with notice to above-mentioned studies, the only thing that can be concluded from this finding is that vitamin E and also with more efficiently honey, due to rich of enzymatic and non-enzymatic antioxidants, done this efficacy, by neutralizing of the injuries patterns of testes integrity (may due to the destructive effects of noise stress) or by regulating of anti-apoptotic patterns. Therefore, in general, each factor that affects the function of interstitial space cells (Leydig and myoid cells) and as well as epithelium germinal cells (sertoli and germ cells), it consequently may lead to suffer a harmful effect on the neuroendocrine gonadal axis and in turn on the spermatogenic cell lineage.

With considering above information, it can suppose that honey and vitamin E by impression on both neuroendocrine gonadal axis and testicular cells may be created condition that the deleterious process (i.e., apoptosis or necrosis) modify or even decrease in the cells. The second assumption is honey and vitamin E: due to enhance steroid hormones levels to sertoli cells, can have a positive effect on nutrition of germ cells. This study also suggests that honey and vitamin E can be useful for keeping the cells in aliveness which suffer from deleterious factors such as noise pollution. The data of this study reinforced the previous studies (29,30). However, Asiyah et al. reported that honey has no more positive influence on the sex hormones concentration in men (31). Therefore, further studies need to elucidate whether honey and vitamin E have a direct effect on sexual hormones production or decreased level of FSH and LH may be as a compensatory mechanism to elevate testosterone level.

In the present study, to show the fertility rate of the noise-exposed rats treated with honey and vitamin E, the male rats were coupled with females. Surprisingly, it was observed that the continuing effects of noise stress injury could reduce the weight of the fetus and the number of the living fetus and survival rate of the fetus. Previously, it has been shown that honey and vitamin E facilitate the recovery of spermatogenesis process by increasing the motility and production of spermatozoa in the epididymis that finally resulted in enhanced fertilization rate in rat (12,18). It was also confirmed that the level of sorbitol dehydrogenase, as a key enzyme in the spermatogenesis, increased following to the honey consumption (10). Honey has also a protective effect on sertoli cells and nourishes them to enhance mature spermatozoa count and affect overall sperm quality. Mahanem et al. indicated that in male rats exposed to cigarette smoking, the serum LH, FSH and testosterone levels changed to optimum levels upon receiving honey (27). In another study, Kimmel et al. (32) observed that in the pregnant rats exposed

to 100 dB noise the number of living fetus and survival rate of fetus were decreased because of increased fetal resorption rate. It seems that the noise pollution has harmful effects on the fertility of male. Also these findings may spell out a natural curative approach rather than pharmaceutical synthetic drugs to optimize of both neuroendocrine gonadal axis and testicular integrity induced by pathogenesis stress, i.e., noise and as well as enhance fertility rate in men.

Conclusion

According to these studies, the noise have a negative impact on male hormones and natural path of sperm development, leading to infertility, and honey and vitamin E as antioxidants reduce the negative effects of noise pollution and enhance the number of living fetuses.

Ethical issues

Ethical of this research work was approved by Ardabil University of Medical Sciences, Iran.

Conflict of interests

We declare that we have no conflict of interests.

Acknowledgments

This study was supported by a research grant (n=PRC-92) from the Ahvaz Jundishapur University of Medical Sciences, Iran. The present study is extracted from master's thesis of Mr. Asghar Rajabzadeh.

References

- 1. von Borell E, Dobson H, Prunier A. Stress, behaviour and reproductive performance in female cattle and pigs. Horm Behav 2007; 52: 130-8.
- 2. Swami CG, Ramanathan J, Charan JC. Noise exposure effect on testicular histology, morphology and on male steroidogenic hormone. Malays J Med Sci 2007; 14: 28-35.
- 3. Cosa M, Cosa G. Annoyance, disturbance and damage caused by noise and vibrations. Ann Ig 1989; 1: 133-56.
- 4. Faldikova L, Diblikova I, Canderle J, Zraly Z, Veznik Z, Sulcova A. Effects of nutrition, social factors and chronic stress on the mouse Leydig cell testosterone production. Veterinární medicína 2001; 46: 160-8.
- 5. Almeida SA, Kempinas WG, Lamano Carvalho TL. Sexual behavior and fertility of male rats submitted to prolonged immobilization-induced stress. Braz J Med Biol Res 2000; 33: 1105-9.
- 6. Chandralekha G. The effect of noise induced stress on the male reproductive endocrine glands of Albino rats [PhD Thesis]. Tamil Nadu, India: Dr. MGR Medical University and Research Chennai; 2002.
- Nawrot PS, Cook RO, Hamm CW. Embryotoxicity of broadband high-frequency noise in the CD-1 mouse. J Toxicol Environ Health 1981; 8: 151-7.
- 8. Bauche F, Fouchard MH, Jegou B. Antioxidant

system in rat testicular cells. FEBS Lett 1994; 349: 392-6.

- 9. Gu W, Hecht NB. Developmental expression of glutathione peroxidase, catalase, and manganese superoxide dismutase mRNAs during spermatogenesis in the mouse. J Androl 1996; 17: 256-62.
- 10. Abdul-Ghani AS, Dabdoub N, Muhammad R, Abdul-Ghani R, Qazzaz M. Effect of Palestinian honey on spermatogenesis in rats. J Med Food 2008; 11: 799-802.
- 11. Estevinho L, Pereira AP, Moreira L, Dias LG, Pereira E. Antioxidant and antimicrobial effects of phenolic compounds extracts of Northeast Portugal honey. Food Chem Toxicol 2008; 46: 3774-9.
- 12. Mohamed M, Sulaiman SA, Jaafar H, Sirajudeen KN. Antioxidant protective effect of honey in cigarette smoke-induced testicular damage in rats. Int J Mol Sci 2011; 12: 5508-21.
- 13. Gheldof N, Wang XH, Engeseth NJ. Identification and quantification of antioxidant components of honeys from various floral sources. J Agric Food Chem 2002; 50: 5870-7.
- 14. Mato I, Huidobro JF, Simal-Lozano J, Sancho MT. Significance of nonaromatic organic acids in honey. J Food Prot 2003; 66: 2371-6.
- 15. Kessopoulou E, Powers HJ, Sharma KK, Pearson MJ, Russell JM, Cooke ID, et al. A double-blind randomized placebo cross-over controlled trial using the antioxidant vitamin E to treat reactive oxygen species associated male infertility. Fertil Steril 1995; 64: 825-31.
- 16. Shalaby MA, Zorba EH. Protective effect of celery oil, Vitamin E and their combination against testicular damage in male rats. Global Veterinaria 2010; 5: 122-8.
- 17. Kutlubay R, Oguz EO, Can B, Guven MC, Sinik Z, Tuncay OL. Vitamin E protection from testicular damage caused by intraperitoneal aluminium. Int J Toxicol 2007; 26: 297-306.
- Momeni HR, Oryan S, Eskandari N. Effect of vitamin E on sperm number and testis histopathology of sodium arsenite-treated rats. Reprod Biol 2012; 12: 171-81.
- 19. Sahinturk V, Guclu C, Baycu C. Protective effects of vitamin E on ethane dimethane sulfonate-induced testicular toxicity in rats. Asian J Androl 2007; 9: 117-24.
- 20. Erat M, Ciftci M, Gumustekin K, Gul M. Effects of nicotine and vitamin E on glutathione reductase activity in some rat tissues in vivo and in vitro. Eur J Pharmacol 2007; 554: 92-7.
- 21. Noguchi J, Yoshida M, Ikadai H, Imamichi T,

Watanabe G, Taya K. Age-related changes in blood concentrations of FSH, LH and testosterone and testicular morphology in a new rat sterile mutant with hereditary aspermia. J Reprod Fertil 1993; 97: 433-9.

- 22. Heidary F, Ahmadi R, Lotfi A. The effects of Cigarette or Hookah smoking on serum levels of LH, FSH or Testosterone in male rats Biological and Pharmaceutical Sciences. Proceedings of the 2nd International Conference on Medical, Biological and Pharmaceutical Sciences (ICMBPS'2012); 2012 Apr 28-29; Singapore.
- 23. Jalali M, Saki G, Sarkaki AR, Karami K, Nasri S. Effect of noise stress on count, progressive and non-progressive sperm motility, body and genital organ weights of adult male rats. J Hum Reprod Sci 2012; 5: 48-51.
- 24. Saki G, Rahim F, Alizadeh K. Effect of forced swimming stress on count, motility and fertilization capacity of the sperm in adult rats. J Hum Reprod Sci 2009; 2: 72-5.
- 25. Saki G, Rahim F, Vaysi OA. Effect of forced swimming stress on in-vivo fertilization capacity of rat and subsequent offspring quality. J Hum Reprod Sci 2010; 3: 32-4.
- 26. Moustafa MH, Sharma RK, Thornton J, Mascha E, Abdel-Hafez MA, Thomas AJ, et al. Relationship between ROS production, apoptosis and DNA denaturation in spermatozoa from patients examined for infertility. Hum Reprod 2004; 19: 129-38.
- 27. Mahanem M, SitiAmrah S, Yatiban M, Hasnan J. Effect of 'Tualang Honey' on spermatogenesis in rats. Malay J Med Sci 2007; 14: 126.
- 28. Naseem M, Goh YM, Hafandi A, Amal NM, Kufli CN, Rajion MA. Effects of vitamin E and soybean oil supplementation on sperm parameters in male Sprague-Dawley rats. Trop Biomed 2007; 24: 45-8.
- 29. Saki G, Jasemi M, Sarkaki AR, Fathollahi A. Effect of administration of vitamins C and E on fertilization capacity of rats exposed to noise stress. Noise Health 2013; 15: 194-8.
- 30. Zakem HB, Alliston CW. The effects of noise level and elevated ambient temperatures upon selected reproductive traits in female Swiss-Webster mice. Lab Anim Sci 1974; 24: 469-75.
- 31. Asiyah HA, Syazana NS, Hashida NH, Sharifah D, Kamaruddin MY. Effects of nicotine and Gelam honey on testis parameters and sperm qualities of juvenile rats. Scientific Research and Essays 2011; 6: 5471-4.
- 32. Kimmel CA, Cook RO, Staples RE. Teratogenic potential of noise in mice and rats. Toxicol Appl Pharmacol 1976; 36: 239-45.

Citation: Rajabzadeh A, Sagha M, Gholami MR, Hemmati R. **Honey and Vitamin E Restore the Plasma Level of Gonadal Hormones and Improve the Fertilization Capacity in Noise-Stressed Rats**. Crescent J Med & Biol Sci 2015; 2(2): 64-8.