



Antioxidants Can Mitigate Decreased Neurogenesis of Neural Stem Cells During Noise Stress

Mahtab Teimouri^{1,*} and Tayebeh Sanchooli^{2,**}

¹Department of Anatomical Sciences, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

²Department of Anatomical Sciences, School of Medicine, Zahedan University of Medical Sciences, Zahedan, Iran

*Corresponding author: Department of Anatomical Sciences, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran. Email: mahtab.teymuri@yahoo.com

**Corresponding author: Department of Anatomical Sciences, School of Medicine, Zahedan University of Medical Sciences, Zahedan, Iran. Email: tasanchooli@gmail.com

Received 2020 January 20; Revised 2020 February 24; Accepted 2020 February 24.

Keywords: Oxidative Stress, Antioxidant Neurogenesis, Neural Stem Cells, Noise, Free Radical

Dear Editor,

Nowadays, neural stem cells (NSCs) are known as an interesting study topic. These ectodermal progenitor cells are generated in the developing central nervous system (CNS) during the fetal period (1). The potential of NSCs to differentiate into neurons (neurogenesis), and glial cells (gliogenesis) will continue for the entire life in the specific areas of brain cortex (2). This potential can be damaged by maternal and environmental stress. Noise stress as an environmental stressor can cause adverse effects in the CNS (3). An experimental study showed that neurogenesis in the dentate gyrus of the hippocampus as the main source of NSCs could be affected by noise stress during the prenatal period (4). Another previous study showed that exposure to high-intensity noise during the postnatal period could suppress neurogenesis in the rat hippocampus, resulting in impaired memory performance and loss of spatial learning (5). Noise stress can cause oxidative stress, leading to the production of free radicals in cells. Embryonic and adult NSCs as the vulnerable cells in the CNS can be damaged by noise stress (6). Antioxidants are molecules that help neutralize the adverse effects of produced free radicals (7). A previous study showed that antioxidant drugs such as N-acetyl-L-cysteine, acetyl-L-carnitine, and phenyl N-tert-butyl nitron (PBN) were impressive in preventing or treating the hearing loss due to noise exposure (8). Also, vitamin E, selenium, manganese, zinc, omega-3, omega-6, and carotenoids provided by foods or supplements could reduce the destructive effects of oxidative stress (9). Due to the decrease in the proliferation ability of NSCs during noise stress, pregnant women and their newborns in the early postnatal period must be preserved from noise stress. This strategy is useful but not possible to imple-

ment in large cities where urban or occupational noise exposure will eventually happen. It is considered that the use of antioxidants is not recommended for pregnant women, and antioxidant supplementary must be used in recommended dietary allowances (RDAs) (10). In conclusion, plant-based foods containing antioxidants are very useful and safe for pregnant women and their offspring against noise stress-mediated damage.

Footnotes

Authors' Contribution: Mahtab Teimouri and Tayebeh Sanchooli equally contributed to the writing of the letter.

Conflict of Interests: There is no potential conflict of interest.

Ethical Approval: There is no ethical consideration for this letter.

Funding/Support: There is no funding/support for this letter.

References

- Martínez-Cerdeño V, Noctor SC. Neural progenitor cell terminology. *Frontiers in neuroanatomy*. 2018;**12**:104.
- Koutmani Y, Karalis KP. Neural stem cells respond to stress hormones: distinguishing beneficial from detrimental stress. *Frontiers in physiology*. 2015;**6**:77.
- Samson J, Sheeladevi R, Ravindran R. Oxidative stress in brain and antioxidant activity of *Ocimum sanctum* in noise exposure. *Neurotoxicology*. 2007;**28**(3):679-85.
- Kim H, Lee M, Chang H, Lee T, Lee H, Shin M, et al. Influence of prenatal noise and music on the spatial memory and neurogenesis in the hippocampus of developing rats. *Brain and Development*. 2006;**28**(2):109-14.

5. Liu L, Shen P, He T, Chang Y, Shi L, Tao S, et al. Noise induced hearing loss impairs spatial learning/memory and hippocampal neurogenesis in mice. *Scientific reports*. 2016;**6**:20374.
6. Golbidi S, Li H, Laher I. Oxidative stress: a unifying mechanism for cell damage induced by noise,(water-pipe) smoking, and emotional stress—therapeutic strategies targeting redox imbalance. *Antioxidants & redox signaling*. 2018;**28**(9):741-59.
7. Azman KF, Zakaria R, Abdul Aziz CB, Othman Z. Tualang honey attenuates noise stress-induced memory deficits in aged rats. *Oxidative medicine and cellular longevity*. 2016;**2016**.
8. Choi SH, Choi C. Noise-induced neural degeneration and therapeutic effect of antioxidant drugs. *Journal of audiology & otology*. 2015;**19**(3):111.
9. Fetoni AR, De Bartolo P, Eramo SLM, Rolesi R, Paciello F, Bergamini C, et al. Noise-induced hearing loss (NIHL) as a target of oxidative stress-mediated damage: cochlear and cortical responses after an increase in antioxidant defense. *Journal of Neuroscience*. 2013;**33**(9):4011-23.
10. Vetrani C, Costabile G, Di Marino L, Rivellese AA. Nutrition and oxidative stress: a systematic review of human studies. *International journal of food sciences and nutrition*. 2013;**64**(3):312-26.