

# The Simultaneous Effects of Aerobic Training and Octopamine on the Mitophagy of the Brown Adipose Tissue after the Induction of Intoxication by Deeply Heated Oil in Male Wistar Rats

Sadegh Abdollahi <sup>1</sup>, Khalid Mohamadzadeh Salamat <sup>2\*</sup>, Kamal Azizbeigi <sup>3</sup>, Zaher Etemad <sup>2</sup>

1. PhD Student, Department of Physical Education, Sanandaj Branch, Islamic Azad University, Sanandaj, Iran.

2. Assistant Professor, Department of Physical Education, Sanandaj Branch, Islamic Azad University, Sanandaj, Iran.

3. Associate Professor, Department of Physical Education, Sanandaj Branch, Islamic Azad University, Sanandaj, Iran.

ARTICLE INFO	ABSTRACT
<p><i>Article type:</i> Review Article</p> <hr/> <p><i>Article History:</i> Received: 27 Mar 2020 Accepted: 18 May 2020 Published: 18 Jul 2020</p> <hr/> <p><i>Keywords:</i> Octopamine LAMP2 PINK Aerobic training</p>	<p><b>Introduction:</b> Deep heated oils produce toxins that pose significant risk to the human health. The present study aims to investigate the simultaneous effects of aerobic training and octopamine consumption on the mitophagy of the brown adipose tissue after the induction of intoxication by deeply heated oil in male Wistar rats.</p> <p><b>Methods:</b> This study was conducted on 40 male Wistar rats after four weeks of feeding with heated oil. The animals were randomly assigned to the groups of control-intoxication (CI; n=8), exercise training-intoxication (ETI; n=8), supplement-intoxication (SI; n=8), supplement-exercise-intoxication (SEI; n=8), and healthy control (HC; n=8). Exercise training was performed for four weeks at the intensity of 50-65% VO<sub>2</sub>max for 20 minutes per session. Octopamine was used as the supplement for four weeks. In addition, intraperitoneal injection was carried out with the dose of 81 μmol/kg five days per week in the CI, SI, and SEI groups. Real-time polymerase chain reaction (RT-PCR) was applied to evaluate the expression of the PTEN-induced kinase 1 (<i>PINK1</i>) and lysosome-associated membrane protein 2 (<i>LAMP2A</i>) genes.</p> <p><b>Results:</b> No significant difference was observed in the <i>PINK1</i> gene expression between the CI, ETI, and SEI groups (P&gt;0.05), while a significant difference was observed between the CI, ETI, and SEI groups with the HC in this regard (P≤0.05). The <i>LAMP2</i> gene expression significantly was higher in the HC group compared to the other groups (P≤0.05). In addition, the <i>LAMP2</i> gene expression in the SEI group was significantly higher than the HC group (P=0.001).</p> <p><b>Conclusion:</b> According to the results, the combination of aerobic training and octopamine supplementation led to mitophagy induction through intoxication with heated oils by increasing the expression of the <i>LAMP2A</i> gene and induction of chaperones by the activation of the chaperones with binding to the LAMP2 receptor protein in the lysosome.</p>

► Please cite this paper as:

Abdollahi S, Mohamadzadeh Salamat Kh, Azizbeigi K, Etemad Z. The Simultaneous Effects of Aerobic Training and Octopamine on the Mitophagy of the Brown Adipose Tissue after the Induction of Intoxication by Deeply Heated Oil in Male Wistar Rats. *J Nutrition Fasting Health*. 2020; 8(3): 151-158. DOI: 10.22038/jnfh. 2020.47446.1256

## Introduction

Obesity is associated with several metabolic diseases, such as type II diabetes, dyslipidemia, hypertension, cardiovascular diseases, and some cancers(1). Lifestyle changes are a consequence of modernity that has caused obesity-related illnesses (1). Such changes not only lead to weight gain, but they could also produce toxins that lead to various diseases due to fast food consumption and the repeated use of heated oils; some of the consequent disorders include cancers and mutagens, increased lipid peroxidation, antioxidant degradation, liver fatty liver tissue damage, and increased apoptosis due to oxidative stress(2,3).

While cooking with heated oils, acrylamide is released from starch at higher doses than the recommended limits (4). Heterocyclic amine is another toxin that is produced in this process, which is formed through the heating and cooking process form the protein substances when amino acids and creatine simultaneously react at high cooking temperatures (barbecued/fried foods) (3).The findings in this regard have indicated that acrolein-induced oxidative stress directly influences the cardiovascular health of rats due to apoptosis in the cardiomyocyte cells. Furthermore, acrolein affects the mitochondrial ability as a component of cellular respiration (4).Other reports have shown that acrolein may

\* Corresponding author: Khalid Mohamadzadeh Salamat, Assistant Professor, Department of Physical Education, Sanandaj Branch, Islamic Azad University, Sanandaj, Iran. Tel: +989128330422. Fax: +988733886661 Email: kh.mohamadzadeh@gmail.com .

© 2020 mums.ac.ir All rights reserved.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

exert inhibitory effects on some cellular respiratory components, such as complex I and II, pyruvate dehydrogenase, and Alpha-Ketoglutaric dehydrogenase in the liver cells of rats(5).

Recently, attempts have been made to diminish the adverse effects of acrolein in the heated oils through exercise training and its synergistic effects with supplementations(6). Today, herbal supplements are considered highly effective in the reduction of oxidative stress and inflammation (7).

Octopamine supplementation has been reported to have antioxidant properties. Octopamine is used as a food ingredient, as well as medicinal and dietary supplementation (8). Octopamine is an adrenergic substance with a sympathetic function. Weight loss and adrenergic actions are associated with metabolic changes (9). Some of the main functions of octopamine are in antioxidant and anti-inflammatory processes, weight loss, fat burning, and anticancer treatments(10). On the other hand, recent findings have indicated that exercise training could decrease oxidative stress and inflammation and improve antioxidant enzymes (11,12). Meanwhile, some studies have reported that exercise training has no significant impact on inflammation and oxidative stress (13). In this study, we used an antioxidant supplement during exercise training in order to reduce inflammation and control diseases (14).

According to the literature, the increased production of reactive oxygen species (ROS) and respiratory chain dysfunction might be associated with decreased PTEN-induced kinase 1(PINK1) activity (15). The reduction of PINK1 expression leads to higher oxidative stress (16), which in turn could lead to apoptosis. The PINK1 protein is composed of a serine/threonine kinase activity, which is located in the mitochondrial membrane and protects cells from ectopic apoptosis (17). On the other hand, the lysosome-associated membrane protein 2 (LAMP2) is a lysosome-associated membrane glycoprotein, which plays a key role in the incidence of cancer and is expressed on the surface of cancerous tumors, especially the cells of highly metastatic cancer, such as colon cancer and melanoma(18). A mitophagy mechanism involves functioning through PINK1 and LAMP2a(10).

Data is scarce regarding the effects of octopamine consumption in the form of an herbal

extract on the induction of the mitophagy caused by the use of heated oils as the use of such oils has been on the rise in various food products.

The present study aimed to determine whether aerobic exercise training and octopamine supplementation could affect the mitophagy of the brown adipose tissue after the induction of intoxication by deeply heated oil.

## Materials and Methods

### Experimental Animals

This study was conducted on 40 male Wistar rats (age: 20 weeks, weight: 300-350 g), which were provided by Pasteur Institute of Iran. The rats were randomly assigned to the groups of control-intoxication (CI; n=8), exercise training-intoxication (ETI; n=8), supplement-intoxication (SI; n=8), supplement-exercise-intoxication (SEI; n=8), and healthy control (HC; n=8). The animals were kept in standard laboratory conditions at the approximate temperature of 25°C and relative humidity of 45% within a 12-hour dark-light cycle until completing the experiment and exercise training. All the animals had free access to water and feed standard for laboratory animals.

The study protocol was in accordance with the Principles of Laboratory Animals and approved by Kurdistan University of Medical Sciences (code: IR.MUK.REC.1398/5006).

### Preparation of the Rations with Hot Oil

Initially, eight liters of sunflower oil was heated for four consecutive days eight hours per day at the temperature of 190-200°C (3), and every 30 minutes, foods such as chicken nuggets, potatoes, chicken, and protein products (e.g., sausages), were immersed in the oil. On the fourth day, the oil was preserved until the start of the experiment to be used as the intoxication agent and administered orally to the animals via gavage as the feed for four weeks.

### Aerobic Exercise Training

The exercise training program was conducted for four weeks at moderate intensity every other day. The intensity of exercise training in the first week reached 50% of the maximum consumed oxygen, while it reached 65% of the maximum consumed oxygen in the last week. In order for the animals to adapt to the exercise protocol, an adaptation exercise training session was implemented at the speed of nine meters per

minute for 20 minutes during one week before the main exercise training program. The duration of exercise training was 20 minutes, with the intensity set at 16 meters per minute on the first day and 26 meters per minute on the last day of the intervention. To initiate the exercise training program, the rats warmed-up for five minutes at the speed of seven meters per minute, and after the main workout, the animals cooled-down at the speed of five meters per minute.

#### **Supplementation and Blood and Tissue Sampling**

Octopamine (Sigma-Aldrich, USA) was used as the supplement for four weeks and was administered via intraperitoneal injection at the dose of 81  $\mu\text{mol/kg}$  five days per week (IP solution with 9% normal saline). At the next stage and 48 hours after the intervention, all the rats had breakfast for 10-8 hours and were weighed before tissue biopsy. Afterwards, anesthesia was performed via inhalation using chloroform. After completing the anesthesia, pain test, and assurance of unconsciousness, blood sampling was carried out from the left ventricle of the heart. Following that, the small intestinal tissues were immediately removed and washed with mucosal PBS. Blood and additional materials were also removed, and the tissues were encoded in two-milliliter microtubes, which were transferred into a nitrogen tank and stored at the temperature of  $-80^{\circ}\text{C}$  until cell analysis.

#### **Biochemical Analysis**

After blood sampling, the serum containing normal saline entered the left ventricle, and the corner of the right atrium was cut with thin scissors to prevent the return of the blood to the heart. After approximately 20 minutes and based on the weight of the rats, complete whitening of the eyes, and returned blood from the organs, the fixative serum (4% aldehyde paraffin) was replaced with normal saline. After 20-30 minutes and the complete fixation of the organs, the serum was removed from the heart, and the brown adipose tissue was excreted from the body, and the tissues were transferred to a secondary fixative container (10% formalin). After 3-5 days, the tissues were transferred to the tissue process apparatus for dehydration and paraffin molding. In this process, the real-time

polymerase chain reaction RT-PCR) was used to evaluate the expression of the *PINK1* and *LAMP2* genes.

#### **RT-PCR**

RT-PCR was used to investigate the gene expression in the brown adipose tissue. For the molecular investigations at the gene expression level, RNA was initially extracted from the tissues of the animals in all the groups in accordance with the instructions of the manufacturer (Qiagen, Germany). To this end, 200  $\mu\text{L}$  of RNeasy Lysis Buffer was added to the samples, and the samples were incubated at the temperature of  $-80^{\circ}\text{C}$  for 24 hours. The plaque available on the cryotube was crushed in a semi-frozen state to lubricate the samples, and 100  $\mu\text{L}$  of chloroform was added to the samples for one minute. The resulting solution was centrifuged at 12,000 rpm for 10 minutes. The transparent liquid on top of the tube containing the RNA was slowly removed and placed in a DEPC microtube. Afterwards, 1 cc of isopropanol was poured on the transparent RNA and stirred manually for one minute.

At the next stage, the samples were centrifuged at 12,000 rpm for 10 minutes, the top liquid was discarded, and 1 cc of 70% alcohol was added to the sediment. After vortexing, the mixture was centrifuged at 7,500 rpm for 10 minutes. Following that, the liquid was drained, and the plaque dried inside the microtube. In addition, 20  $\mu\text{L}$  of distilled water ( $60^{\circ}\text{C}$ ) was poured on the plaque and placed on a plaque ( $60^{\circ}\text{C}$ ) for five minutes. After RNA extraction from all the studied samples with high purity and concentration, cDNA synthesis was performed in accordance with the instructions of the manufacturer (Fermentas, USA), and the synthesized cDNA was used for the reverse transcription reaction. The *PINK1* expression was measured using the quantitative model of RT-PCR. The primers were designed based on the data of *PINK1* and glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) available in the NCBI gene bank, as well as the data obtained by Macrogen Company. The *GAPDH* gene was used as the control gene, and its expression rate was calculated using the  $\Delta\Delta\text{CT}-2$  formula. Table 1 shows the sequence of the applied primers.

**Table 1.**Primer Sequence

Gene	Oligo Sequence 5'-3'
<i>PINK1</i>	F5' CAAAGGGAGTAGATGGGGAG3' R5' GATGGATGGGAATAGGTAGTGA3'
<i>LAMP2</i>	F5' GCAGTGCAGATGAAGACAAC3' R5' AGTATGATGGCGCTTGAGAC3'
<i>GAPDH</i>	F5' CAT ACT CAG CAC CAG CAT CAC C3' R5'AAG TTC AAC GGC ACA GTC AAG G 3'

### Immunohistochemical (IHC) Assay

At this stage, 20 grams of paraformaldehyde (Merck, Germany) was added to 500 cc of distilled water, and the mixture was placed on a hot plate until its temperature reached 60°C. Afterwards, NaOH was added to the solution until the pH of the solution reached the range of 7.2-7.4. Finally, aPBS tablet was added to the solution. In the next step, 2 cc of 37.5% hydrochloric acid was added to 100 cc of distilled water. In addition, 100 Landa of fetal bovine serum was added to 900 Landa of PBS, and 10% bovine serum was obtained. Following that, 3 Landa of Triton 100x was added to 997 Landa of PBS, and 0.3% Triton was obtained.

The samples were washed with PBS in four steps at five-minute intervals between each step. In order to restore the antigens, normal hydrochloric acid was poured on the samples for 30 minutes. In order to neutralize the acid, Borate buffers were added for five minutes, and the cells were washed with PBS. In addition, 0.3% Triton was used for 30 minutes to make the cell membranes permeable, and the cells were washed with PBS. In order to block the reaction of the secondary antibody, the 10% bovine serum was added in the form of an additional background color for 30 minutes. Following that, the primary antibody (1-100) was diluted with PBS and added to the samples. After creating a moist environment for the prevention of the tissues from drying, the samples were placed in a refrigerator at the temperature of 2-8°C overnight. On the following day, the tissue container was taken out of the refrigerator and washed four times with PBS (five minutes each time). In the next step, the secondary antibodies with the dilutions of 1-150 were added to the samples and incubated at the temperature of 37°C for one hour and 30 minutes in the dark


using an incubator. Afterwards, the samples were transferred from the incubator to a dark room, and after they were washed four times, DAPI was added, immediately removed, and poured on the PBS samples. Finally, the samples were observed using the Olympus fluorescence microscope with a 400 lens to confirm the markers.

### Statistical Analysis


Data analysis was performed in SPSS version 22, and the data were expressed as mean and standard deviation. In order to determine the effects of intoxication by heated oil, t-test was used for the comparison of the findings between the independent groups of HC and intoxication-control. Furthermore, two-way analysis of variance (ANOVA) was used to assess the main effects of exercise training and octopamine, as well as the simultaneous effects of octopamine. Bonferroni's trial test was also applied in case significant differences were observed. In all the statistical analyses, the significance level was considered at  $P \leq 0.05$ .

### Results

The findings indicated that the expression of the *PINK* gene significantly increased due to intoxication by deeply heated oil ( $P=0.001$ ). On the other hand, exercise training could significantly reduce the *PINK* gene expression ( $F=11.49$ ;  $P=0.003$ ;  $\mu=0.365$ ). Octopamine supplementation had no significant effect on the *PINK* gene expression ( $F=2.13$ ;  $P=0.160$ ;  $\mu=0.096$ ). Although the lowest *PINK* gene expression was observed in the octopamine and exercise-training group, the interactive effects between exercise and octopamine supplementation on the *PINK* gene expression were not considered significant ( $F=0.42$ ;  $P=0.514$ ;  $\mu=0.022$ ) (Figure 2).



**Figure 1.** Comparison of *PINK* Gene Expression in Healthy Controls and Controls Poisoned with Deeply Heated Oil (\*Significant changes compared to healthy control)




**Figure 2.** Expression of *PINK* Gene in Study Groups (Data expressed as mean and standard deviation)


According to the obtained results, the expression of the *LAMP2A* gene significantly decreased ( $P=0.001$ ) due to intoxication by deeply heated oil. On the other hand, exercise training significantly increased the *LAMP2A* gene expression ( $F=9.63$ ;  $P=0.006$ ;  $\mu=0.325$ ). Octopamine supplementation also significantly increased the *LAMP2A* gene expression ( $F=21.25$ ;  $P=0.001$ ;  $\mu=0.515$ ). The interaction of exercise training and octopamine

supplementation had a significant effect on the *LAMP2A* gene expression ( $F=4.80$ ;  $P=0.040$ ;  $\mu=0.194$ ). Considering the amount of *LAMP2A* gene expression in the exercise-training and supplementation groups, it was expected that the expression of the *LAMP2A* gene would be higher than the other interventions. This was confirmed by the findings as the expression of the *LAMP2A* gene was higher in the mentioned groups compared to the other interventions (Figure 4).





**Figure 3.** Comparison of *LAMP2A* Gene Expression in Healthy Controls and Controls Poisoned with Deeply Heated Oil (\*Significant changes compared to control)



**Figure 4.** Expression of *LAMP2A* Gene in Study Groups (Data expressed as mean and standard deviation)

### Discussion

The present study aimed to investigate the simultaneous effects of aerobic exercise training and octopamine consumption on the mitophagy of the brown adipose tissue. The obtained results indicated a significant difference in the *PINK1* gene expression between the HC and intervention groups, which could be due to the fact that heated oils increase the expression of the *PINK* gene. According to the literature, the consumption of heated oils significantly increases the formation of metastatic lung tumors (19). Moreover, the toxic compounds produced in burnt oils and assimilation of oil in the Wistar rats increased the lipid peroxidation and liver enzymes (20).

The findings of the current research indicated no significant difference between the SI, SEI, and EIT groups in terms of the *PINK1* gene expression. In other words, the combination of octopamine with exercise training has not more advantages than exercise training or octopamine supplementation alone. Although the difference in this regard was not considered significant, octopamine supplementation resulted in the reduction of the *PINK1* gene expression. Therefore, it could be concluded that as an emulgor of adrenergic compounds, octopamine could exert antioxidant effects on the brown adipose tissue of rats. These findings support a new model for the elimination of damaged mitochondria, where *PINK1* utilizes parkin for malfunctioning mitochondria to stimulate

mitochondrial deletion by mitophagy (21). *PINK1* reduction is associated with increased mitophagy (22). Through increasing the total antioxidant capacity and activity of the red blood cells in the body, this interaction could prevent oxidative stress and inappropriate production of the creatine kinase indicator of muscle damage and malondialdehyde and carbonylated protein indicators of lipid peroxidation.

According to the results of the present study, exercise training and octopamine supplementation had significant effects on the expression of the *LAMP2A* gene in the Wistar rats intoxicated by deeply heated oil, and the maximum effect could be achieved when the two interventions are combined. In this regard, Han et al. (2017) reported that aerobic exercise training could change the gene expression of *LAMP2A* through a high-fat diet, thereby modulating oxidative stress and improving the oxidative indicators of the brown adipose tissue in obese rats (23).

In another study, Park et al. (2016) observed a significant difference in the antioxidant indicators after a period of aerobic exercise training (24). Furthermore, the mentioned study indicated that intoxication by heated oils blocked the regulation of the *LAMP2A* gene expression by affecting its receptors and moving the mitophagy regulation in a negative direction, thereby decreasing the *LAMP2A* gene expression. *LAMP2a* is a protein receptor located in the lysosomal membrane, inside which the lysosomes break down with the autophagosome, causing the inner membrane to become lubricious and the inner components of the autophagosome to decompose. The toxicities that are induced by ROS could reduce the expression of the *LAMP2A* gene through inactivating the hsc70 or mutagen receptors in these receptors, which in turn increases the production of ROS in damaged mitochondria. Therefore, the continued consumption of burned fats may adversely affect the content of lipid mutagens, cause cellular damage to the liver and kidneys, increase urinary mutagenicity, and increase the cell proliferation of the *esophagus*(19).

The findings of the current research demonstrated that exercise training, octopamine supplementation, and their interactive effects increased the expression of the *LAMP2A* gene, thereby activating oxidative stress as a result of

intoxication with oil through inducing mitophagy by the chaperones (25). Furthermore, they are moved to the lysosome surface by hsc70 with the help of the chaperones, and after binding to the 2A-LAMP receptor protein on the lysosome surface, the substrates cross the membrane width through the 2A-LAMP-dependent displacement complex and rapidly decompose into the lumen(22) According to the results of the present study, it seems that the interactive effects of aerobic exercise training and octopamine consumption could induce mitophagy and decrease the ROS produced through intoxication by the heated oils.

### Conclusion

According to the results, exercise training and octopamine supplementation might have significant effects on the improvement of mitophagy and prevention of the damages caused by the use of heated oils in the brown adipose tissue. Given the scarcity of data in this regard and limitations of our study regarding the loss of the laboratory rats, it is recommended that the current study be replicated using more samples in order to obtain more generalizable findings in the future.

### Acknowledgments

This article was extracted from a doctoral dissertation in the field of sports physiology. The study was conducted at the Histogenetic Research Laboratory of Tehran in 2019 and funded by the researchers. We would like to thank all those who assisted us in this research project.

### References

- Huang CJ, McAllister MJ, Slusher AL, Webb HE, Mock T, Acevedo EO. Obesity-related oxidative stress: the impact of physical activity and diet manipulation. *Sports Med Open*. 2015. 1(1): 32.
- Joseph N, Nelliyanil M, Rai S, Babu R, Kotian SM, Ghosh T, et al. Fast food consumption pattern and its association with overweight among high school boys in Mangalore city of southern India. *J Clin Diagn Res*. 2015. 9(5): LC13-7.
- Kim TH, Choi D, Kim JY, Lee JH, Koo SH. Fast food diet-induced non-alcoholic fatty liver disease exerts early protective effect against acetaminophenintoxication in mice. *BMC Gastroenterol*. 2017. 17(1): 124.
- Ekiz E, Oz F. The effects of different frying oils on the formation of heterocyclic aromatic amines in meatballs and the changes in fatty acid compositions

- of meatballs and frying oils. *J Sci Food Agric*. 2019; 99(4): 1509-18.
5. Wang L, Sun Y, Asahi M, Otsu K. Acrolein, an environmental toxin, induces cardiomyocyte apoptosis via elevated intracellular calcium and free radicals. *Cell Biochem Biophys*. 2011; 61(1): 131-6.
  6. Carvalho VH, Oliveira AHS, de Oliveira LF, da Silva RO, Mascio PD, Gualano B, et al. Exercise and  $\beta$ -alanine supplementation on carnosine-acrolein adduct in skeletal muscle. *Redox Biol*. 2018; 18: 222-8.
  7. Almasi J, Azizbeigi K, Mohammadzadeh Salamat K, Naghshtabrizi B, Azarbaijani MA. The effect of taking resveratrol supplement during heart rehabilitation exercises on the lipid profile of patients undergoing coronary artery bypass graft surgery: randomized clinical trial. *Journal of Knowledge & Health*. 2018; 13(2): 8-14. (Persian)
  8. Sun Y, Ito S, Nishio N, Tanaka Y, Chen N, Isobe KI. Acrolein induced both pulmonary inflammation and the death of lung epithelial cells. *Toxicol Lett*. 2014. 229(2): 384-92.
  9. Thevis M, Koch A, Sigmund G, Thomas A, Schänzer W. Analysis of octopamine in human doping control samples. *Biomed Chromatogr*. 2012. 26(5): 610-5.
  10. de Oliveira AL, de Paula MN, Comar JF, Vilela VR, Peralta RM, Bracht A. Adrenergic metabolic and hemodynamic effects of octopamine in the liver. *Int J Mol Sci*. 2013; 14(11): 21858-72.
  11. Atashak S, Stannard SR, Azizbeigi K. Cardiovascular risk factors adaptation to concurrent training in overweight sedentary middle-aged men. *J Sports Med Phys Fitness*. 2016; 56(5): 624-30.
  12. Azizbeigi K, Stannard SR, Atashak S. Green Tea Supplementation During Resistance Training Minimally Affects Systemic Inflammation and Oxidative Stress Indices in Obese Men. *Jundishapur J Nat Pharm Prod*. 2019; 14(1): e61419.
  13. Libardi CA, De Souza GV, Cavaglieri CR, Madruga VA, Chacon-Mikahil MPT. Effect of resistance, endurance, and concurrent training on TNF- $\alpha$ , IL-6, and CRP. *Med Sci Sports Exerc*. 2012; 44(1): 50-6.
  14. Alishahi A, Azizbeigi K, Mohammadzadeh Salamat K, Yektayar M. The Effect of Aerobic Training with Vitamin C Supplementation on Myeloperoxidase, Asymmetric Dimethyl Arginine and Blood Pressure in Middle-Age Hypertensive Overweight Men, *J Clin Res Paramed Sci*. Online ahead of Print ; 8(2): e86843.
  15. Wood-Kaczmar A, Gandhi S, Yao Z, Abramov AY, Miljan EA, Keen G, et al. PINK1 is necessary for long term survival and mitochondrial function in human dopaminergic neurons. *PLoS One*. 2008; 3(6): e2455.
  16. Valente EM, Salvi S, Ialongo T, Marongiu R, Elia AE, Caputo V, et al. PINK1 mutations are associated with sporadic early-onset parkinsonism. *Ann Neurol*. 2004; 56(3): 336-41.
  17. Sarafian V, Jadot M, Foidart JM, Letesson JJ, Van den Brûle F, Castronovo V, et al. Expression of Lamp-1 and Lamp-2 and their interactions with galectin-3 in human tumor cells. *Int J Cancer*. 1998; 75(1): 105-11.
  18. Xin-Fang L, Jumat S, Rais MM, Kamsiah J. Effect of Repeatedly Heated Palm Olein on Blood Pressure—Regulating Enzymes Activity and Lipid Peroxidation in Rats. *Malays J Med Sci*. 2012; 19(1): 20-9.
  19. Adraskela K, Veisaki E, Koutsilieris M, Philippou A. Physical exercise positively influences breast cancer evolution. *Clin Breast Cancer*. 2017; 17(6): 408-17.
  20. Vives-Bauza C, Przedborski S. Mitophagy: The latest problem for Parkinson's disease. *Trends Mol Med*. 2011; 17(3): 158-65.
  21. Yao Z, Gandhi S, Burchell VS, Plun-Favreau H, Wood NW, Abramov AY. Cell metabolism affects selective vulnerability in PINK1-associated Parkinson's disease. *J Cell Sci*. 2011; 124(Pt 24): 4194-202.
  22. Burenjargal M, Totani N. Cytotoxic compounds generated in heated oil and assimilation of oil in Wistar rats. *J Oleo Sci*. 2009; 58(1): 1-7.
  23. Park SY, Kwak YS. Impact of aerobic and anaerobic exercise training on oxidative stress and antioxidant defense in athletes. *J Exerc Rehabil*. 2016; 12(2): 113-7.
  24. Deas E, Piipari K, Machhada A, Li A, Gutierrez-del-Arroyo A, Withers DJ, et al. PINK1 deficiency in  $\beta$ -cells increases basal insulin secretion and improves glucose tolerance in mice. *Open Biol*. 2014; 4(5): 140051.