

# Estimating greenhouse gas emissions using emission factors from the Sugarcane Development Company, Ahvaz, Iran

Amir Zahedi<sup>1</sup>, Sorour Ghodrati<sup>2</sup>, Mehdi Ahmadimoghaddam<sup>3</sup>, Nematollah Jaafarzadeh<sup>4\*</sup>

<sup>1</sup>Committee Research, Department of Environmental Health Engineering, School of Health, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

<sup>2</sup>Department of Environmental Assessment and Land Use Planning, School of Environment and Energy, Islamic Azad University, Science and Research Branch, Tehran, Iran

<sup>3</sup>Environmental Technologies Research Center, Department of Environmental Health Engineering, School of Health, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

<sup>4</sup>Environmental Technologies Research Center, Department of Environmental Health Engineering, School of Health, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

## Abstract

**Background:** Greenhouse gas (GHG) emissions are increasing worldwide. They have harmful effects on human health, animals, and plants and play a major role in global warming and acid rain.

**Methods:** This research investigated carbon dioxide (CO<sub>2</sub>) and CH<sub>4</sub> emissions obtained from different parts of the Hakim Farabi, Dobar Khazaei, and Ramin factories which produce ethanol and yeast. Seasonal rates of CO<sub>2</sub> at the soil surface at the studied sites were estimated from measurements made on location and at intervals with manual chambers. This study aimed to assess the production rate of GHG emissions (CH<sub>4</sub>, CO<sub>2</sub>) in the sugar production units of Hakim Farabi, Dobar Khazaei, and Ramin factories.

**Results:** Mean concentrations of CO<sub>2</sub> and CH<sub>4</sub> emissions are respectively 279 500.207 and 3087.07 tons/year from the Hakim Farabi agro-industry, 106 985.24 and 1.14 tons/year at the Dobar Khazaei ethanol producing factory, and 124 766.17 and 1.93 tons/year at the Ramin leavening producing factory.

**Conclusion:** Sugar plant boilers and the burning of sugarcane contributed the most CO<sub>2</sub> and CH<sub>4</sub> emissions, respectively. Moreover, lime kilns and diesel generators showed the least carbon dioxide and methane emissions, respectively.

**Keywords:** Carbon Dioxide, Methane, Ethanol, Farms, Global Warming

**Citation:** Zahedi A, Ghodrati S, Ahmadimoghaddam M, Jaafarzadeh N. Estimating greenhouse gas emissions using emission factors from the Sugarcane Development Company, Ahvaz, Iran. Environmental Health Engineering and Management Journal 2018; 5(1): 9–14. doi: 10.15171/EHEM.2018.02.

## Article History:

Received: 23 April 2017

Accepted: 21 August 2017

ePublished: 3 January 2018

## \*Correspondence to:

Nematollah Jaafarzadeh

Email: Jaafarzadeh-n@ajums.ac.ir

## Introduction

Greenhouse gases (GHGs) cause a significant amount of pollution in the atmosphere (1). They are harmful to human health, animals, and plants, and they play a major role in global warming and acid rain (2). During recent decades, public concern has arisen regarding the presence of air pollutants both outdoors and indoors, and their potential to damage human health has increased (3,4). Carbon dioxide (CO<sub>2</sub>) and CH<sub>4</sub> are the main GHGs today; CO<sub>2</sub> emissions from fossil fuels are the main cause of global warming (5,6). Other studies showing the importance of CO<sub>2</sub> and CH<sub>4</sub> rates in global warming have been conducted in some industrial and agricultural countries (7-11). The great development of the sugarcane and ethanol production industries, including agricultural industries and other technologies, seen in recent decades has created a need for recalculating the emission of

pollutants and energy balances. Methane and CO<sub>2</sub> are the main emissions of the agricultural/industrial levels (12). In the last 10 years, atmospheric concentrations of CO<sub>2</sub> have increased by almost 40%, from a preindustrial concentration of approximately 280 ppmv (parts per million volume) to almost 384 ppmv in 2007 (13). Large reductions in GHG emissions are expected to prohibit serious climate destabilization, however, GHG emissions from fixed and transport sources are growing more quickly than other energy-using sectors and are predicted to increase GHG emissions by 80% between 2007 and 2030 (14). CH<sub>4</sub> is produced during the production and transport of coal, natural gas, oil, and other natural resources such as fertilizer and landfills with open-path technology and other anaerobic waste treatment systems (15,16). Municipal landfills are great resources of CH<sub>4</sub> emissions; in municipal solid waste landfills, CH<sub>4</sub> is about



**Table 1.** CO<sub>2</sub> and CH<sub>4</sub> emission factors in sugarcane factories

Emission resource	Gas	Emission factor	Unit	Reference
Sugarcane straw Burning in the field	CO <sub>2</sub>	1.66	g/kg	CEPA (23)
	CH <sub>4</sub>	2.7	g/kg	CEPA
Trash burning in the field	CO <sub>2</sub>	1.3 – 1.4	kg/tones	EPA (24)
	CH <sub>4</sub>	2.7	kg/tones	IPCC (24)
Boiler emission	CO <sub>2</sub>	1891	g/m <sup>3</sup> fuel	CAPP (25)
	CH <sub>4</sub>	0.03	g/m <sup>3</sup> fuel	CAPP
Limestone burning emission	CO <sub>2</sub>	3.14	kg/m <sup>3</sup> fuel	Canada can met energy diversification research laboratory (26)
	CH <sub>4</sub>	0.12	Ton CO <sub>2</sub> /1000 m <sup>3</sup> fuel	CAPP
Diesel generator	CO <sub>2</sub>	2.68	Kg CO <sub>2</sub> /L fuel	SEIA (27)
	CH <sub>4</sub>	0.13	g CH <sub>4</sub> /L fuel	Canada national inventory report

12000 hectares in the Farabi farm. The main sources of GHG emissions in this complex are open burning in the field, boilers, limestone burning, diesel generators, and electrical equipment. Emission rates for the different parts were calculated using the following formula. Table 1 indicates the emission factor for CO<sub>2</sub> and CH<sub>4</sub> gases in sugarcane factories as different guidelines. Emission factor and emission rate are common and simple indices used to show GHG emissions. Table 2 indicates the CO<sub>2</sub> and CH<sub>4</sub> emission rates at different units of the Dobal Khazaei factory.

Emission rate = emission factor \* fuel type (22)

The emission factors in different parts of the factory based on type of fuel are as follows:

Emission factor = Emission rate / Fuel type

Figure 2 shows the carbon dioxide and methane emission rates during harvest; this emission factor obtained from bagasse burning that occurred as open-field burning at the Hakim Farabi farm.




Dobal Khazaei factory is one of the main factories in the region. It produces the largest amounts of ethanol and leaven in the country. In this factory, there are 3 sources of CO<sub>2</sub> emissions (boilers, the ethanol unit, and electrical equipment) and two sources of CH<sub>4</sub> emissions (boilers and electrical equipment). Figure 3 shows a comparison of CO<sub>2</sub> and CH<sub>4</sub> emission rates from different emission sources at the Hakim Farabi factory. Figure 4 shows the CO<sub>2</sub> and CH<sub>4</sub> emission rates at the Dobal Khazaei factory.

### Methane and carbon dioxide emissions

Methane and carbon dioxide are emitted from different parts of the Hakim Farabi and Dobal Khazaei complexes, including the burning of sugarcane fields before harvest,

**Table 2.** CO<sub>2</sub> and CH<sub>4</sub> emission rates from Dobal Khazaei factory

Emission resource	Gas	Emission (ton/year)
Boiler (ethanol production )	CO <sub>2</sub>	10892.16
	CH <sub>4</sub>	0.20
Leaven production process	CO <sub>2</sub>	14000
Boilers (Leaven production process)	CO <sub>2</sub>	7412.72
	CH <sub>4</sub>	0.14


**Figure 2.** Open-field burning at Hakim Farabi farm.**Figure 3.** Comparison of CO<sub>2</sub> and CH<sub>4</sub> emissions from different sources at Hakim Farabi Sugar factory

using stillage as fertilizer, boilers, and burning ethanol in ethanol engines. Results of this study indicate the emission factor for  $\text{CH}_4$  and  $\text{CO}_2$  in Hakim Farabi farm. During the burning of sugarcane trash in the field,  $\text{CH}_4$  is the main gas produced and emitted;  $\text{CO}_2$  emissions are insignificant. Therefore, the bulk of  $\text{CH}_4$  emissions are produced in field by trash burning, whereas  $\text{CO}_2$  emissions resulting from open burning in the field are related to bagasse and cane waste. Figure 5 shows the comparison of  $\text{CO}_2$  and  $\text{CH}_4$  emissions from different sources. A recent study that analyzed emission factors for sugarcane and simulated burning in a wind tunnel indicated that GHG emissions were 0.32 kg/t (dry fuel) in the case of a spreading fire and 0.59 kg/t (dry fuel) for fire in a pile. In transient or disturbed boiler processes,  $\text{CH}_4$  emissions are produced only from unburnt organics.


### Discussion

In this study, GHG emissions from five main factory sources were investigated, including the sugarcane farm, boilers, limestone burning, diesel generators, and electrical equipment. The Farabi cane crop was transferred from the farm to the Farabi, Dobal Khazaei, and Ramin factories to become the final products such as ethanol and sugarcane. Therefore, sugarcane mills produce sugar, ethanol, Leaven, which are splatted and sent to respond to market needs. Straw and sugarcane that are cut and burned on the Hakim Farabi farm produce the greatest amounts of  $\text{CO}_2$  and  $\text{CH}_4$  emissions. In addition, bagasse, the residue of

sugarcane sap dewatered from sugarcane, is combusted in sugar factories to produce steam and electricity. Nguyen and Gheewala showed in their study that the methane emission rate from sugarcane burning is 3.5 kg/ton. They also revealed that open burning on the farm caused a  $\text{CH}_4$  emission rate of 2.2 kg/ton. Their results were obtained by the emission factor (28). The Hakim Farabi and Dobal Khazaei factories are major producers of sugar, leaven, and ethanol, and their activities produce a large quantity of  $\text{CO}_2$  and  $\text{CH}_4$  emissions. Ordinarily, GHG emissions are produced in winter, which is the harvest season. The GHGs emissions from these factories along with particulate matter (PM) are considered synergistic agents. The California Environmental Protection Agency revealed that the  $\text{CO}_2$  and  $\text{CH}_4$  emission rates from the open burning of sugarcane for the production of ethanol are 315.973 and 514.1 g/ton, respectively (29), but in the current study,  $\text{CO}_2$  and  $\text{CH}_4$  emission rates of 1660 and 2700 g/ton were determined (Figure 2). Boilers are another part of the factories that release high concentrations of  $\text{CO}_2$  and  $\text{CH}_4$  emissions into the atmosphere. Boilers, lime furnaces, diesel generators, and electrical equipment are internal units in the Hakim Farabi Complex and are used in the production of sugar and ethanol. The GHG emissions in these processes are caused by the internal burner. This unit of the factory uses natural gas and gasoline; therefore, the emission and energy rates related to gasoline are affected by the class of the crude oil, the petroleum refinery configuration, and



**Figure 4.** Comparison of  $\text{CO}_2$  and  $\text{CH}_4$  emissions from different sources at Dobal Khazaei factory



**Figure 5.** Comparison of  $\text{CO}_2$  and  $\text{CH}_4$  emissions from different sources in the whole complex



such factors as natural gas, petrol, and gasoline quality (11). The lime furnace uses natural gas, and therefore produces a high amount of  $\text{CH}_4$ . The power plant unit is one of the main parts of the Hakim Farabi complex. It uses six diesel generators to supply electricity to the entire Hakim Farabi complex. In this unit, the use of gasoline fuel could not be determined as local reliable documents were not available. Thus, international consolidated data about energy consumption and GHG emissions in the production of oil-derived fuels was used in this study. The electrical equipment is another main unit in the Hakim Farabi factory. This unit supplies electrical energy for the entire Hakim Farabi complex.

Total bagasse production is almost 1129222 tons/year (2014) at 50% moisture. It is the sum of bagasse which percentage of bagasse will be lost in different ways, and this is used as fuel in some industries such as paper pulp and food industries. However, the main part of bagasse is used in sugar mills as an energy resource to produce sugar or ethanol. Also some of bagasse are used to produce sugar and ethanol. The bagasse to sugar ratio is 1:1, and the bagasse to ethanol ratio is 0.5:1.

Sugarcane farming correlates with a significant amount of GHG emissions such as  $\text{CO}_2$  and  $\text{CH}_4$ , from both upstream processes such as fertilizer production and from the field itself. The nitrogen in bagasse (i.e. straw) on the farm is also the N in fertilizer and emits  $\text{N}_2\text{O}$ . Open field burning was done in farm, by physical harvesting of sugarcane, and transportation logistics. These two activities are the main GHG emission sources in the sugarcane ethanol life cycle. In the last decade, mechanical harvesting increased the speed of operations. This technology also increased fuel consumption by 20%; however, the use of bigger equipment and new technology is replacing these methods. Automatic harvesting will increase to 50% in the next 8 years (10). However, mechanical harvesting will increase GHG emissions in the atmosphere. Moreover, an increase in green cane reaping will produce a huge volume of trash. This could increase GHG emissions due to open burning in the field. Most boilers do not have wet scrubbers, and  $\text{CH}_4$  emissions have not been reported. The most significant pollutants emitted by bagasse boilers are  $\text{CO}_2$  and PM, which can travel long distances in the atmosphere and be transferred to Ahvaz city atmosphere. Due to the higher volume of gases emitted from this agro-industry complex, it is suggested that this factory should use scrubbers and filters to control gases and use biofilters to remove gases through the use of microorganisms.

### Conclusion

The Hakim Farabi, Dobal Khazaei, and Ramin are the most important sugar and ethanol factories in Iran, and they release a great amount of GHG emissions into the atmosphere. In these factories, GHG emissions come from five main sources: the sugarcane farm boilers, limestone burning, diesel generators, and electrical equipment. In

these factories, straw and bagasse are burned in the field, and open burning released significant concentrations of  $\text{CO}_2$  and  $\text{CH}_4$  into the atmosphere. Ordinarily, GHG emissions are produced in winter, which is the harvest season. The GHG emissions from these factories along with PMs are considered synergistic agents. The emission rates for the different parts of the factories were calculated. The results showed that in the Hakim Farabi factory, boiler emissions, electrical equipment, diesel generator, limestone burning emissions, sugarcane straw burning in the field, and trash burning in the field released the highest concentrations of  $\text{CO}_2$ , respectively. Furthermore, sugarcane straw burning and trash burning in the field produced the biggest percentage of methane emissions. In open burning, the  $\text{CH}_4$  emissions are greater than the  $\text{CO}_2$  emissions. In the Dobal Khazaei factory, electrical equipment, ethanol, and boilers released the highest concentrations of  $\text{CO}_2$  into the atmosphere. Also, the electrical equipment produced the highest rate of  $\text{CH}_4$  emissions into the atmosphere.

### Acknowledgments

This research project was financially supported by the Environmental Technologies Research Center, Ahvaz Jundishapur University of Medical Sciences (grant No. ETRC 9012).

### Ethical issues

It is confirmed that this manuscript is the original work of the authors. It has not been published, nor is it under review in another journal, and it is not being submitted for publication elsewhere.

### Competing interests

The authors declare that they have no competing interests.

### Authors' contribution

All authors contributed equally and participated in the collection, analysis, and interpretation of the data. All authors critically reviewed, refined, and approved the manuscript.

### References

1. Samaras C, Meisterling K. Life cycle assessment of greenhouse gas emissions from plug-in hybrid vehicles: implications for policy. *Environ Sci Technol* 2008; 42(9): 3170-6. doi: 10.1021/es702178s.
2. Weisser D. A guide to life-cycle greenhouse gas (GHG) emissions from electric supply technologies. *Energy* 2007; 32(9): 1543-59. doi: 10.1016/j.energy.2007.01.008.
3. Bastianoni S, Pulselli FM, Tiezzi E. The problem of assigning responsibility for greenhouse gas emissions. *Ecol Econ* 2004; 49(3): 253-7. doi: 10.1016/j.ecolecon.2004.01.018.
4. Jaafarzadeh N, Amiri H, Ahmadi M. Factorial experimental design application in modification of volcanic ash as a natural adsorbent with Fenton process for arsenic removal. *Environ Technol* 2012; 33(2): 159-65. doi:

- 10.1080/09593330.2011.554887.
5. Davis SJ, Caldeira K. Consumption-based accounting of CO<sub>2</sub> emissions. *Proc Natl Acad Sci U S A* 2010; 107(12): 5687-92. doi: 10.1073/pnas.0906974107.
  6. Stocker TF, Qin D, Plattner GK, Tignor M, Allen SK, Boschung J, et al. *Climate change 2013: The physical science basis*. Intergovernmental Panel on Climate Change, Working Group I Contribution to the IPCC Fifth Assessment Report (AR5). New York: Cambridge Univ Press; 2013.
  7. Smith KR, Desai MA, Rogers JV, Houghton RA. Joint CO<sub>2</sub> and CH<sub>4</sub> accountability for global warming. *Proceedings of the National Academy of Sciences* 2013; 110(31): E2865-74. doi: 10.1073/pnas.1308004110.
  8. Taylor M, Tam C, Gielen D. Energy efficiency and CO<sub>2</sub> emissions from the global cement industry. *Korea* 2006; 50(2-2): 61-7.
  9. de Carvalho Macedo I. Greenhouse gas emissions and energy balances in bio-ethanol production and utilization in Brazil (1996). *Biomass and Bioenergy* 1998; 14(1): 77-81. doi: 10.1016/S0961-9534(97)00038-X.
  10. Denmead OT, Macdonald B, Bryant G, Naylor T, Wilson S, Griffith DWT, et al. Emissions of methane and nitrous oxide from Australian sugarcane soils. *Agric For Meteorol* 2010; 150(6): 748-56. doi: 10.1016/j.agrformet.2009.06.018.
  11. Macedo IC, Seabra JE, Silva JE. Green house gases emissions in the production and use of ethanol from sugarcane in Brazil: The 2005/2006 averages and a prediction for 2020. *Biomass Bioenergy* 2008; 32(7): 582-95. doi: 10.1016/j.biombioe.2007.12.006.
  12. Rafiei M, Gadgil AS, Ghole VS, Gore SD, Jaafarzadeh N, Mirkazemi R. Assessment of air pollution and its effects on the health status of the workers in beam rolling mills factory (Iran National Steel Industrial Group) from Ahvaz-Iran. *Indian J Occup Environ Med* 2009; 13(1): 20-2. doi: 10.4103/0019-5278.50719.
  13. Doney SC, Fabry VJ, Feely RA, Kleypas JA. Ocean acidification: the other CO<sub>2</sub> problem. *Ann Rev Mar Sci* 2009; 1: 169-92. doi: 10.1146/annurev.marine.010908.163834.
  14. Woodcock J, Edwards P, Tonne C, Armstrong BG, Ashiru O, Banister D, et al. Public health benefits of strategies to reduce greenhouse-gas emissions: urban land transport. *Lancet* 2009; 374(9705): 1930-43. doi: 10.1016/s0140-6736(09)61714-1.
  15. Burba G, Anderson T, Ediger K, von Fischer J, Gioli B, Ham J, et al, eds. *Latest on Mobile Methane Measurements with Fast Open-Path Technology: Experiences, Opportunities & Perspectives*. Vienna: EGU General Assembly; 2016.
  16. Ge S, Liu L, Xue Q, Yuan Z. Effects of exogenous aerobic bacteria on methane production and biodegradation of municipal solid waste in bioreactors. *Waste Manag* 2016; 55: 93-8. doi: 10.1016/j.wasman.2015.11.024.
  17. Jorgenson AK. *Global Warming and the Neglected Greenhouse Gas: A Cross-National Study of the Social Causes of Methane Emissions Intensity*, 1995. *Social Forces* 2006; 84(3): 1779-98. doi: 10.1353/sof.2006.0050.
  18. Howarth RW, Santoro R, Ingraffea A. Methane and the greenhouse-gas footprint of natural gas from shale formations. *Clim Change* 2011; 106(4): 679-90. doi: 10.1007/s10584-011-0061-5.
  19. Searchinger T, Heimlich R, Houghton RA, Dong F, Elobeid A, Fabiosa J, et al. Use of U.S. Croplands for Biofuels Increases Greenhouse Gases Through Emissions from Land-Use Change. *Science* 2008; 319(5867): 1238-40. doi: 10.1126/science.1151861.
  20. Satterthwaite D. Cities' contribution to global warming: notes on the allocation of greenhouse gas emissions. *Environ Urban* 2008; 20(2): 539-49. doi: 10.1177/0956247808096127.
  21. Newell RG, Raimi D. Implications of Shale Gas Development for Climate Change. *Environ Sci Technol* 2014; 48(15): 8360-8. doi: 10.1021/es4046154.
  22. Shahsavani A, Naddafi K, Jafarzade Haghhighifard N, Mesdaghinia A, Yunesian M, Nabizadeh R, et al. The evaluation of PM<sub>10</sub>, PM<sub>2.5</sub>, and PM<sub>1</sub> concentrations during the Middle Eastern Dust (MED) events in Ahvaz, Iran, from April through September 2010. *J Arid Environ* 2012; 77: 72-83. doi: 10.1016/j.jaridenv.2011.09.007.
  23. Crilly D, Zhelev T. Emissions targeting and planning: An application of CO<sub>2</sub> emissions pinch analysis (CEPA) to the Irish electricity generation sector. *Energy* 2008; 33(10): 1498-507. doi: 10.1016/j.energy.2008.05.015.
  24. Venkatram A. A critique of empirical emission factor models: a case study of the AP-42 model for estimating PM<sub>10</sub> emissions from paved roads. *Atmos Environ* 2000; 34(1): 1-11. doi: 10.1016/S1352-2310(99)00330-1.
  25. Chambers A, Strosher M, Wootton T, Moncrieff J, McCready P. DIAL measurements of fugitive emissions from natural gas plants and the comparison with emission factor estimates. 15th Annual Emission Inventory Conference, US Environmental Protection Agency New Orleans; 2006.
  26. Ordorica-Garcia G, Croiset E, Douglas P, Elkamel A, Gupta M. Modeling the Energy Demands and Greenhouse Gas Emissions of the Canadian Oil Sands Industry. *Energy Fuels* 2007; 21(4): 2098-111. doi: 10.1021/ef0700984.
  27. Murray SN, Walsh BP, Kelliher D, O'Sullivan DTJ. Multi-variable optimization of thermal energy efficiency retrofitting of buildings using static modelling and genetic algorithms – A case study. *Build Environ* 2014; 75: 98-107. doi: 10.1016/j.buildenv.2014.01.011.
  28. Nguyen TLT, Gheewala SH. Life cycle assessment of fuel ethanol from cane molasses in Thailand. *Int J Life Cycle Assess* 2008; 13(4): 301. doi: 10.1007/s11367-008-0011-2.
  29. Detailed C. GREET pathway for California reformulated gasoline blendstock for oxygenate blending (CARBOB) from average crude refined in California, Version 2.1. Sacramento, CA: California Air Resources Board; 2009.