



DOI: 10.32768/abc.201854187-189

Warfarin Induced Necrosis of The Breast: A Case Report

Sanaz Zand^a, Roghaye Jalili^b, Seyed Hasan Emami Razavi^b, Massoome Najafi^{*b,c}, Ahmad Kaviani^{a,b}

^a Department of Research, Kaviani Breast Disease Institute (KBDI), Tehran, Iran

^b Department of Surgery, Tehran University of Medical Sciences (TUMS), Tehran, Iran

^c International Center for Surgical Safety (ICSS), St. Michael Hospital, University of Toronto, Canada

ARTICLE INFO

Received:

18 October 2018

Revised:

28 October 2018

Accepted:

08 November 2018

Key words:

Warfarin,
breast necrosis,
breast gangrene,
oral anticoagulant therapy

ABSTRACT

Background: Warfarin induced breast necrosis is a rare complication of oral anticoagulant therapy. Although it can be related to protein C, S, and antithrombin III deficiency; the pathogenesis of necrosis is still unknown.

Case presentation: We report a case of a 38-year-old woman with extensive left breast necrosis after receiving warfarin for treatment of deep vein thrombosis. Simple mastectomy was performed and the wound was closed secondarily with an abdominal advancement flap. Rivaroxaban was prescribed after discontinuation of warfarin.

Conclusion: Although breast necrosis following warfarin usage is uncommon, it should be considered in women presenting with breast symptoms after initiation of warfarin. Early diagnosis and appropriate management are essential to prevent extensive loss of breast tissue.

Introduction

Warfarin is one of the most commonly prescribed oral anticoagulant medications, and it is used for the treatment of deep vein thrombosis (DVT) and pulmonary embolism (PE); and for thromboembolism prophylaxis in patients with valvular heart disease, mechanical heart valves, and atrial fibrillation.¹

The most common complication of warfarin is bleeding,¹ which can be prevented by adjusting international normalized ratio (INR). Necrosis, osteoporosis, calcification, and purple toe syndrome are other side effects. Skin and subcutaneous tissue necrosis is an extremely rare and serious complication of warfarin with a prevalence of 0.1- 0.01 percent.² It was first described by Flood *et al.* in 1943.³ Since then, there has been more than 300 reported cases in the literature,² but less than 40 of the reported cases involved the breast tissue.⁴

We present a case of warfarin induced necrosis of the breast in a 38-year-old patient, treated with simple mastectomy.

Case presentation

A 38-year-old female presented to the breast clinic of Imam Khomeini Hospital with the complaint of a necrotic ulcer on her left breast for 10 days. She had a history of admission for deep vein thrombosis (DVT) of the left lower limb one month earlier. The deep vein thrombosis occurred following long-term use of oral contraceptive pills (OCP). She received intravenous heparin followed by warfarin. She was discharged from the hospital with a daily dose of 10 mg of warfarin and with an INR of 2.1.

On admission for breast necrosis, no fever was detected but the patient had a foul-smelling necrosis involving most of her left breast (Figure 1). Blood work showed INR, activated thromboplastin time (aPTT), and platelets of 1.32, 38 seconds, and 460,000 per cubic milliliter, respectively. Hematology consultation recommended discontinuation of warfarin and administration of Rivaroxaban.

Considering the extent of breast necrosis (Figure 1), mastectomy was recommended. Before surgery, Rivaroxaban was discontinued, and heparin and

Address for correspondence:

Massoome Najafi, MD
Address: Imam Khomeini Hospital Complex, Keshavarz Blvd.,
1419733141, Tehran, Iran.
Tel: +98 21 61192509
Fax: +98 21 66935063
Email: mdmasoume@yahoo.com



Figure 1. A, B: Left breast necrosis, C: Removed specimen

broad-spectrum antibiotics were started. The patient underwent left mastectomy. There were purulent secretions under the necrotic tissue and the wound was left open. Intravenous antibiotics were continued for 7 days. The wound irrigation and dressing were performed daily. The patient was discharged from the hospital with Rivaroxaban 7.5 mg/ twice a day for 21 days, then 20 mg/ day, and daily irrigation of the wound was arranged at a local clinic. Healing of the wound was monitored by the surgeon on a weekly basis. After one month, the wound was closed with abdominal advancement flap. Anticoagulation was continued for 6 months.

Further hematologic evaluation showed protein C deficiency. Pathology report confirmed breast parenchyma with diffuse and extensive necrosis and acute inflammation.

Discussion

Skin and subcutaneous tissue necrosis is a rare complication of warfarin, occurring in 0.1-0.01% of patients receiving oral anticoagulation.² Women are more commonly affected with a female to male ratio of 3:1.⁵ The areas with increased subcutaneous fat (abdomen, thighs, buttocks, legs, and breasts) are more susceptible to necrosis.^{6,9} Breast is affected in about 10-15% of the patients.⁴ It usually affects one breast, but bilateral breast necrosis is also reported.^{4,10}

The symptoms start 1 to 10 days after initiation of warfarin, and the majority of cases occur within 3 to 6 days.^{4,6} However, there are reports of late-onset symptoms several years after commencement of therapy.¹¹⁻¹³ The initial manifestations of breast necrosis are paresthesia, pressure, and discomfort in the affected area. Later, erythematous or hemorrhagic skin lesions appear and will develop into blisters and bullae, leading to skin necrosis and scar formation.^{4,8}

It is important to consider other conditions that can mimic warfarin necrosis in differential diagnosis, such as micro-embolization (septic emboli, cholesterol emboli), heparin-induced skin necrosis as a result of the thrombocytopenia and thrombosis syndrome, disseminated intravascular coagulation, purpura fulminans, necrotizing fasciitis, cryoglobulinemia, inflammatory breast cancer, decubitus ulcers, snake venom induced skin necrosis, cellulitis, venous gangrene, necrotizing fasciitis, cellulitis, venous gangrene, calciphylaxis, and hematoma.^{1,14}

The exact pathogenesis of necrosis following warfarin usage is still unclear, however protein C deficiency, hypersensitivity, and direct toxic effect of warfarin are possible mechanisms of necrosis.^{1,6} The most popular pathogenesis is acute protein C deficiency. Warfarin blocks vitamin K epoxide reductase enzyme and inhibits synthesis of clotting factors II, VII, IX, and X. It also inhibits synthesis of



anticoagulant protein C and protein S. Protein C and protein S have shorter half-life than vitamin K dependent clotting factors. Consequently, the patient develops a transient hypercoagulable state at the beginning of warfarin therapy and needs another form of anticoagulation, such as heparin, for 48-72 hours to prevent clot formation.^{1,7,8,15}

Management of breast necrosis following oral anticoagulation therapy with warfarin, starts with discontinuation of warfarin. However, withdrawing warfarin will not affect the course of the established skin lesions.^{4,5,16} Vitamin K and fresh-frozen plasma are administered to replenish protein C levels,⁸ and anticoagulation should be continued by administering intravenous heparin or the non-vitamin K antagonist oral anticoagulants (NOACs) such as Dabigatran or Rivaroxaban.¹⁷ The necrotic lesions should be managed surgically with debridement and appropriate wound care.

In conclusion, fulminant painful purpura and erythematous skin lesions in patients receiving warfarin should be suspected as warfarin induced necrosis. Immediate discontinuation of warfarin, administration of vitamin K, and fresh-frozen plasma is recommended. The skin lesion should be managed with appropriate surgical debridement and wound care.

Ethical Consideration

The patient announced her consent for using her data for this case report.

Conflict of Interest

The authors have no conflict of interest to disclose.

References

1. Simman R, Gould N, Jackson S. Warfarin-Induced Tissue Necrosis (WITN): Case Report and Literature Review. *J Surg Oper Care*. 2015;1(1):101.
2. Chan YC, Valenti D, Mansfield AO, Stansby G. Warfarin induced skin necrosis. *Br J Surg*. 2000;87(3):266-72.
3. Flood E. Thrombophlebitis migrans disseminata: report of a case in which gangrene of abreast occurred: observations on the therapeutic use of dicumarol (3, 3'-methylenebis (4-hydroxycoumarin)). *NY State J Med*. 1943;43:1121-4.
4. Ejzenberg D, Neusquen LP, Rolnik DL, Lozinsky AC, Piato JR. Breast necrosis induced by the use of coumadin: case report and review of literature. *Einstein (Sao Paulo)*. 2015;13(3):417-9.
5. Fred HL. Skin Necrosis Induced by Coumarin Congeners. *Tex Heart Inst J*. 2017;44(4):233-6.
6. Nazarian RM, Van Cott EM, Zembowicz A, Duncan LM. Warfarin-induced skin necrosis. *J Am Acad Dermatol*. 2009;61(2):325-32.
7. Khalid K. Warfarin-induced necrosis of the breast: case report. *J Postgrad Med*. 2004;50(4):268-9.
8. Au AF, Fosnot J, Wu LC. Coumadin-induced skin necrosis of the breasts: case report. *Ann Plast Surg*. 2012;69(1):109-10.
9. Davis CE, Jr., Wiley WB, Faulconer RJ. Necrosis of the female breast complicating oral anticoagulant treatment. *Ann Surg*. 1972;175(5):647-56.
10. Lopez Valle CA, Hebert G. Warfarin-induced complete bilateral breast necrosis. *Br J Plast Surg*. 1992;45(8):606-9.
11. Essex DW, Wynn SS, Jin DK. Late-onset warfarin-induced skin necrosis: case report and review of the literature. *Am J Hematol*. 1998;57(3):233-7.
12. Goldberg SL, Orthner CL, Yalisove BL, Elgart ML, Kessler CM. Skin necrosis following prolonged administration of coumarin in a patient with inherited protein S deficiency. *Am J Hematol*. 1991;38(1):64-6.
13. Sorensen DM, Eilbert W. Images in emergency medicine. Man with painful skin lesion. Warfarin sodium skin necrosis. *Ann Emerg Med*. 2010;55(3):302, 5.
14. Eby CS. Warfarin-induced skin necrosis. *Hematol Oncol Clin North Am*. 1993;7(6):1291-300.
15. Fawaz B, Candelario NM, Rochet N, Tran C, Brau C. Warfarin-induced skin necrosis following heparin-induced thrombocytopenia. *Proc (Bayl Univ Med Cent)*. 2016;29(1):60-1.
16. Gelwix TJ, Beeson MS. Warfarin-induced skin necrosis. *Am J Emerg Med*. 1998;16(5):541-3.
17. Bakoyiannis C, Karaolanis G, Patelis N, Maskanakis A, Tsaples G, *et al*. Dabigatran in the Treatment of Warfarin-Induced Skin Necrosis: A New Hope. *Case Rep Dermatol Med*. 2016;2016:3121469.