

Hospital-Acquired Infections in Elderly Versus Younger Patients in an Acute Care Hospital

Perla Sarai Solis-Hernandez,¹ Melissa Vidales-Reyes,¹ Elvira Garza-Gonzalez,² Guillermo Guajardo-Alvarez,³ Susana Chavez-Moreno,⁴ and Adrian Camacho-Ortiz^{1,*}

¹Hospital Epidemiology, Hospital Universitario Dr Jose Eleuterio Gonzalez, Monterrey, Mexico

²Gastroenterology Service, Hospital Universitario Dr Jose Eleuterio Gonzalez, Monterrey, Mexico

³Geriatric Service, Hospital Universitario Dr Jose Eleuterio Gonzalez, Monterrey, Mexico

⁴School of Medicine, University of Monterrey, Monterrey, Mexico

*Corresponding author: Adrian Camacho-Ortiz, Hospital Epidemiology, Hospital Universitario Dr Jose Eleuterio Gonzalez, Monterrey, Mexico. Tel: +52-8183482767, Fax: +52-8183482767, E-mail: acamacho_md@yahoo.com

Received 2015 August 26; Accepted 2015 September 2.

Abstract

Background: A growing number of elderly patients are hospitalized for various causes and age has been described inconsistently as a risk factor for acquiring nosocomial infections with a subsequent higher mortality rate compared to younger patients.

Objectives: To describe the incidence, type, and microbiological characteristics of nosocomial infections in elderly and non-elderly patients.

Patients and Methods: Retrospective analysis of all hospital-acquired infections (HAIs) in an academic community hospital. Patients were stratified into two groups: non-elderly (18 - 64 years) and elderly (> 65 years).

Results: A total of 18469 patients were included (108555 hospital days) in this study. About 79.6% of HAI were infected non-elderly and 20.3% elderly ($P < 0.0001$) patients. Higher infection rates for ventilator-associated pneumonia (VAP), secondary bacteremia, and catheter-associated urinary tract infections were noted for elderly as well as a higher frequency of positive cultures for Gram-negative bacteria, particularly *Klebsiella* spp. and fungal infections ($P > 0.05$). Hospital mortality increased with every HAI diagnosed per patient.

Conclusions: The study showed that HAIs were more frequent in elderly population predominately with respect to VAP and infections by Gram-negative pathogens. Overall mortality was greater in the elderly group although the odds ratio for death was higher in younger patients and increased with every HAI diagnosed.

Keywords: Elderly, Gram-Negative Bacteria, Cross Infection, Mortality, Ventilator-Associated, Pneumonia

1. Background

A growing number of the global population is aging; accordingly a higher number of elderly patients are hospitalized for various causes. According to several studies (1-3) conducted in this population, age is a risk factor for acquiring nosocomial infections with a subsequent higher mortality rate compared to younger patients. Meanwhile another study (4) found no significant difference between elderly patients and younger population.

The available studies on this subject are mostly based on European population (1, 2, 5-7), and those studies on Latin ethnicity (4) were conducted using a smaller number of patients.

2. Objectives

This study aimed to describe the incidence, type, and microbiological characteristics of nosocomial infections, among elderly and non-elderly patients.

3. Patients and Methods

3.1. Setting

This retrospective study was performed from January 2013 to December 2013 at the University Hospital "Dr. Jose Eleuterio Gonzalez" tertiary care hospital with 450 beds that provides medical attention to an open population in Monterrey, Northeastern region of Mexico.

3.2. Study Population

During the study period, all hospitalized patients were included in the nosocomial surveillance for hospital-acquired infections. We eliminated from the study all patients under the age of 18. Patients were stratified in two groups; non-elderly, including all patients with an age range of 18 - 64 years and elderly, including all patients older than 64 years.

All patients with diagnosis of a hospital-acquired infection were further studied for type of the infection, microbial diagnosis, hospital stay, and mortality.

3.3. Definitions

A nosocomial infection was defined by the site of infection using diagnostic criteria from the Center for Disease Control (CDC) (8). Hospital epidemiological surveillance was performed based on CDC and WHO recommendations (9).

3.4. Case Surveillance and Diagnosis of HAI

The infection control unit includes infectious disease specialists, a hospital epidemiologist, doctors and nurses trained in infection control. All cases of suspected HAI were detected by experienced personnel from the infection control unit and were later discussed with the rest of the unit's staff for a definitive classification.

3.5. Statistical Analysis

A descriptive analysis was performed using basic parametric test. The Student t-test was used for numerical variables and Chi-square or Fisher exact test for analyzing categorical variables. $P < 0.05$ was considered statistically significant.

4. Results

Over the study period, 18469 patients were hospitalized with a total of 108555 hospital days. Of those patients, 15796 patients belonged to non-elderly and 2673 to elderly group.

In the elderly population, the primary diagnoses for hospitalization were as follow: cardiovascular (coronary artery disease, cerebral-vascular disease), 17.8%; malignant tumor related illness, 12.9%; trauma and orthopedic related diagnoses, 8.1%; diabetes related complications, 4.8%; and respiratory diseases, 3.6%. For the non-elderly patients, the primary diagnoses were as follows: obstetrics and or pregnancy related: 39.3%; abdominal surgeries: 10.1%; accidents related diagnoses: 8.3%; malignant tumor related illness: 6.8%; and respiratory tract diseases: 2.2%.

We recorded a total of 862 HAIs (79.6% in non-elderly patients and 20.3% in elderly patients) with an overall higher infection rate in the elderly group (6.54 HAI per 100 admissions) versus the non-elderly (4.34 HAI per 100 admissions) group (Table 1).

When comparing both groups with a specific type of hospital acquired infection, we found that ventilator-associated pneumonia (VAP), secondary bacteremia (SB), and catheter associated urinary tract infections (CA-UTI) had a higher incidence in the elderly group compared to


the non-elderly group (0.67 vs. 1.20 per 1000 ventilator-days for VAP, $P = 0.023$; 0.23 vs. 0.67 per 1000 catheter-days for SB, $P = 0.002$; and 1.74 vs. 2.66 per 1000 catheter-days for CA-UTI, $P = 0.010$) (Table 2).

Rates of VAP, CA-UTI, and central line associated bloodstream infection (CLABSI) had an increasing tendency as aged groups progressed and surgical site infection (SSI) had a decreasing tendency. Although in the 75 - 84 age group HAI rates had an increase, the rates were lower in the > 85 years group (Figure 1).

More than two-thirds (67.3%) of the isolates in the whole study group were Gram-negative bacteria while Gram-positive bacteria were reported in only 25.9% of the isolates. The elderly group had a higher frequency of positive cultures for Gram-negative bacteria, particularly *Klebsiella* spp. and fungal infections ($P > 0.05$). *Clostridium difficile* infection rate was not different between both age groups (Table 3).

Overall, in-hospital mortality in patients with HAI was statistically lower in the non-elderly population 20.5% vs. 38.5% in elderly group (OR 0.41, 95% CI; 0.27 - 0.62, $P < 0.0001$). In-hospital mortality increased in association with the number of HAIs diagnosed per patient, but it only reached statistical significance when only 1 HAI was diagnosed. In patients diagnosed with only 1 HAI, the mortality rate was 14.2% in the non-elderly vs. 30.8% in the elderly group (OR 2.69, 95% CI; 1.60 - 4.52, $P = 0.0002$). When 2 HAIs were diagnosed in the non-elderly group the mortality rose to 37.8% vs. 54.5% in the elderly group (OR 1.97, 95% CI; 0.75 - 5.15, $P = 0.166$). Finally, when 3 or more HAIs were diagnosed, the mortality rose to 52.5% vs. 72.2% in the elderly group (OR 2.4, 95% CI; 0.55 - 10.4, $P = 0.23$). Unlike the patients without HAI (by age group), all groups showed differences in higher probability of in-hospital deaths (Table 4).

Figure 1. Rate of Hospital-Acquired Infections by Age Groups



VAP: ventilator associated pneumonia; CLABSI: central line associated bloodstream infections; CA-UTI: catheter-associated urinary tract infections; SSI: surgical site infection.

Table 1. Infection Rate of Hospital-Acquired Infections According to the Age Groups

Age groups, y	No. of Patients	Patients-Day	No. of HAIs	Infection Rate Per 100 Admissions	Infection Rate Per 1000 Patient-Days
18 - 64	15,796	89,427	687	4.3492 ^a	7.6822 ^b
> 65	2,673	19,128	175	6.5470 ^a	9.1489 ^b
Total	18,469	108,555	862	4.6673	7.9407

^aDifference between this two groups $P < 0.0001$.

^bDifference between this two groups $P < 0.0442$.

Table 2. Number, Percentage and Rate of Hospital-Acquired Infections by Sites and by Age Group

Variable	Non-Elderly		Elderly		Total		P
	No. (%)	Infection Rate Per 1000 p-d	No. (%)	Infection Rate Per 1000 p-d	No. (%)	Infection Rate Per 1000 p-d	
Ventilator-associated pneumonia	60 (8.7)	0.670	23 (13.1)	1.202	83 (9.6)	0.764	0.023
Pneumonia	20 (2.9)	0.223	5 (2.9)	0.261	25 (2.9)	0.230	0.960
Other respiratory tract infections	4 (0.6)	0.044	1 (0.6)	0.052	5 (0.6)	0.046	0.888
Primary bacteremia	46 (6.7)	0.514	8 (4.6)	0.418	54 (6.3)	0.497	0.717
Secondary bacteremia	20 (2.9)	0.223	13 (7.4)	0.679	33 (3.8)	0.304	0.002
Central line-associated bloodstream infections	62 (9)	0.693	16 (9.1)	0.836	78 (9)	0.718	0.602
Intravascular catheter insertion site infection	49 (7.1)	0.547	11 (6.3)	0.575	60 (7)	0.552	0.884
Catheter-associated urinary tract infection	156 (22.7)	1.744	51 (29.1)	2.666	207 (24)	1.906	0.010
Urinary tract infection	15 (2.2)	0.167	6 (3.4)	0.313	21 (2.4)	0.193	0.302
Surgical site infection	196 (28.5)	2.191	28 (16)	1.463	224 (26)	2.063	0.054
Skin and soft tissues infections	27 (3.9)	0.301	5 (2.9)	0.261	32 (3.7)	0.294	0.948
Eye infections	1 (0.1)	0.011	2 (1.1)	0.104	3 (0.3)	0.027	0.141
Peritonitis	3 (0.4)	0.033	0 (0)	0.000	3 (0.3)	0.027	0.965
Gastrointestinal infections	23 (3.3)	0.257	6 (3.4)	0.313	29 (3.4)	0.267	0.849
Other	5 (0.7)	0.055	0 (0)	0.000	5 (0.6)	0.046	0.654
Total	687 (100)	7.682	175 (100)	9.148	862 (100)	7.940	0.044

Table 3. Distribution of Microorganisms Isolated From Hospital-Acquired Infections by Age Groups^a

Microorganisms	Non-Elderly	Elderly	Total	P
Gram-negative	420 (67.4)	90 (66.7)	510 (67.4)	
<i>Escherichia coli</i>	59 (9.5)	15 (11.1)	74 (9.8)	0.209
<i>Pseudomonas</i> spp.	103 (16.5)	23 (17)	126 (16.6)	0.278
<i>Klebsiella</i> spp.	62 (10)	20 (14.8)	82 (10.8)	0.016
<i>Acinetobacter</i> spp.	107 (17.2)	18 (13.3)	125 (16.5)	0.981
<i>Proteus</i> spp.	10 (1.6)	0 (0)	10 (1.3)	0.394
<i>Enterobacter</i> spp.	24 (3.9)	6 (4.4)	30 (4.0)	0.547
<i>Stenotrophomonas maltophilia</i>	7 (1.1)	1 (0.7)	8 (1.1)	0.874
<i>Citrobacter</i> spp.	8 (1.3)	0 (0)	8 (1.1)	0.508
<i>Morganella morganii</i>	8 (1.3)	2 (1.5)	10 (1.3)	0.962
<i>Providencia</i> spp.	21 (3.4)	3 (2.2)	24 (3.2)	0.783
<i>Serratia</i> spp.	8 (1.3)	2 (1.5)	10 (1.3)	0.962
Other Gram-negative	3 (0.5)	0 (0)	3 (0.4)	0.476
Gram-positive	164 (26.3)	32 (23.7)	196 (25.9)	
<i>Enterococcus</i> spp.	66 (10.6)	14 (10.4)	80 (10.6)	0.540
Coagulase-negative staphylococci	20 (3.2)	2 (1.5)	22 (2.9)	0.678
<i>Staphylococcus aureus</i>	47 (7.5)	10 (7.4)	57 (7.5)	0.637
<i>Streptococcus</i> spp.	8 (1.3)	0 (0)	8 (1.1)	0.508
<i>Clostridium difficile</i>	23 (3.7)	6 (4.4)	29 (3.8)	0.436
Yeast	39 (6.3)	13 (9.6)	52 (6.9)	
<i>Candida</i> spp.	39 (6.3)	13 (9.6)	52 (6.9)	0.049
Total	623 (100)	135 (100)	758 (100)	

^aValues are presented as No. (%)

Table 4. Twenty-Eight Day-Mortality Associated With the Number of HAI Diagnosed During the Same Hospitalization^a

	Non-Elderly					Elderly				
	28 Day-Mortality ^b	Alive at Discharge ^b	OR	95% CI	P	28 Day-Mortality ^b	Alive at Discharge ^b	OR	95% CI	P
1 HAI vs. No HAI	14.20	85.70	6.39	4.75 to 8.59	0.0001	30.80	69.10	4.21	2.66 to 6.64	0.0001
2 HAI vs. No HAI	37.80	62.10	23.47	14.51 to 37.96	0.0001	54.50	45.40	11.32	4.84 to 26.47	0.0001
> 3 HAI vs. No HAI	52.50	47.50	42.62	22.73 to 79.94	0.0001	72.70	27.20	25.15	6.63 to 95.45	0.0001

^aAbbreviation: HAI: hospital-acquired infection.

^bValues are presented based on %.

5. Discussion

In the present study, the HAI rate per 100 admissions and per 1000 patient days was greater in the elderly group ($P < 0.0001$ and $P = 0.04$, respectively); these results are similar to other reports assessing the same topic (2, 3). Whereas, a Brazilian study showed discordant data (4) in which no significant difference was found between both groups, which was probably due to the small number of patients analyzed.

We found that the most common types of HAIs were SSI, CA-UTI, VAP, and CLABSI. Similar to our results, Avci et al. (2) reported a higher incidence of urinary tract infections, respiratory tract infections, surgical site, skin and soft tissue infections, per 1000 patient days in the elderly group. A study on critically ill patients found that bloodstream infections were less frequent in patients aged 85 and older compared to the younger patients and there were no significant difference in the other types of infection (10). Another study on postoperative patients reported no statistically significant difference between young and elderly patients (7).

Stephan et al. found an increased incidence of pneumonia, SSI, ITU, bacteremia, and catheter-associated infection in the 60 - 75 age group (the younger group of the population analyzed) and a decreased incidence in the age group of older than 75 years. These results were similar to those obtained in our study, in which we observed that in 85 years and older group, the major HAIs in our hospital (VAP, CLABSI, CR-UTI, SSI) had a lower incidence, which was probably due to fewer number of invasive procedures in this age group (7).

The most frequent microorganisms isolated were Gram-negative bacteria with a greater incidence in the elderly group ($P = 0.03$). This finding was also described by Dimopoulos et al. (10) reporting that Gram-negative microorganisms were more frequent in patients aged 85 and older than in all groups younger than 75. They found no significant statistical differences in the frequencies of Gram-positive microorganisms and fungi infections, opposite to our study in which we observed a greater incidence in the fungi infections in the elderly ($P = 0.0085$).

In this study, the 28 day-mortality in patients with HAIs was higher compared to patients that did not developed

HAI in both the elderly and non-elderly groups; this issue has been reported in other studies (11).

According to several studies, elderly patients are associated with an increased mortality rate (2, 3, 5-7, 10). Similar to these, in our hospital the overall mortality in patients with HAIs was statistically higher in the elderly group ($P < 0.0001$). We acknowledge limitations to our study such as the difference in the primary diagnosis and comorbidities between the groups which made the comparison cumbersome.

In sum, we conclude that HAIs were more frequent in the elderly population predominately with respect to VAP and Gram-negative isolates. Mortality was higher in the elderly population although the associated risk was higher in the younger patients with every diagnosed HAI.

Footnote

Authors' Contribution: Study concept and design: Elvira Garza-Gonzalez and Adrian Camacho-Ortiz; acquisition of data: Perla Sarai Solis-Hernandez and Melissa Vidales-Reyes; analysis and interpretation of data: Elvira Garza-Gonzalez; drafting of the manuscript: Adrian Camacho-Ortiz; critical revision of the manuscript for important intellectual content: Elvira Garza-Gonzalez, Guillermo Guajardo-Alvarez, and Adrian Camacho-Ortiz; statistical analysis: Adrian Camacho-Ortiz; administrative, technical, and material support: Adrian Camacho-Ortiz; study supervision: Adrian Camacho-Ortiz.

References

1. Ellidokuz H, Ucku R, Uysal U, Abacioglu H. Hospital-acquired infections in elderly patients: results of a West Anatolian University Hospital surveillance. *Arch Gerontol Geriatr.* 2003;**37**(3):259-63. [PubMed: 14511851]
2. Avci M, Ozgenc O, Coskuner SA, Olut AI. Hospital acquired infections (HAI) in the elderly: comparison with the younger patients. *Arch Gerontol Geriatr.* 2012;**54**(1):247-50. doi: 10.1016/j.archger.2011.03.014. [PubMed: 21529974]
3. Laurent M, Bories PN, Le Thuaut A, Liuu E, Ledudal K, Bastuji-Garin S, et al. Impact of comorbidities on hospital-acquired infections in a geriatric rehabilitation unit: prospective study of 252 patients. *J Am Med Dir Assoc.* 2012;**13**(8):760 e7-12. doi: 10.1016/j.jamda.2012.07.002. [PubMed: 22885408]

4. Ribas RM, Gontijo Filho PP. Comparing hospital infections in the elderly versus younger adults: an experience in a Brazilian University Hospital. *Braz J Infect Dis.* 2003;7(3):210-5. [PubMed: 14499044]
5. Blot S, Cankurtaran M, Petrovic M, Vandijck D, Lizy C, Decruyenaere J, et al. Epidemiology and outcome of nosocomial bloodstream infection in elderly critically ill patients: a comparison between middle-aged, old, and very old patients. *Crit Care Med.* 2009;37(5):1634-41. doi: 10.1097/CCM.0b013e31819da98e. [PubMed: 19325489]
6. Beaujean DJ, Blok HE, Vandenbroucke-Grauls CM, Weersink AJ, Raymakers JA, Verhoef J. Surveillance of nosocomial infections in geriatric patients. *J Hosp Infect.* 1997;36(4):275-84. [PubMed: 9261757]
7. Stephan F, Cheffi A, Bonnet F. Nosocomial infections and outcome of critically ill elderly patients after surgery. *Anesthesiology.* 2001;94(3):407-14. [PubMed: 11374598]
8. Garner JS, Jarvis WR, Emori TG, Horan TC, Hughes JM. CDC definitions for nosocomial infections, 1988. *Am J Infect Control.* 1988;16(3):128-40. [PubMed: 2841893]
9. Ducloux G, Fabry J, Nicolle L. WHO/CDS/CSR/EPH/ Prevention of hospital acquired infections: a practical guide. 2002.
10. Dimopoulos G, Koulenti D, Blot S, Sakr Y, Anzueto A, Spies C, et al. Critically ill elderly adults with infection: analysis of the extended prevalence of infection in intensive care study. *J Am Geriatr Soc.* 2013;61(12):2065-71. [PubMed: 24479140]
11. Vrijens F, Hulstaert F, Devriese S, Van de Sande S. Hospital-acquired infections in Belgian acute-care hospitals: an estimation of their global impact on mortality, length of stay and healthcare costs. *Epidemiol Infect.* 2012;140(1):126-36. doi: 10.1017/S0950268811000100. [PubMed: 21320376]

Archive of SID