

Soil Contamination with Soil Transmitted Helminthes in Schools and Play Areas of Kermanshah City, West of Iran

Mohammad Ali Mohaghegh,¹ Mohammad Reza Vafaei,¹ Mehdi Azami,² Nooshin Hashemi,³ Seyed Hossein Hejazi,^{1,2} Farzaneh Mirzaei,^{1,4} Hamed Kalani,¹ Mohammad Falahati,¹ Somayeh Jahani,⁵ and Mohsen Ghomashlooyan^{1,*}

¹Department of Parasitology and Mycology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, IR Iran

²Skin Diseases and Leishmaniasis Research Center, Isfahan University of Medical Sciences, Isfahan, IR Iran

³North Khorasan University of Medical Sciences, Bojnurd, IR Iran

⁴School of Paramedicine, Shahid Sadoughi University of Medical Sciences, Yazd, IR Iran

⁵Infectious Diseases and Tropical Medicine Research Center, Zahedan University of Medical Sciences, Zahedan, IR Iran

*Corresponding author: Mohsen Ghomashlooyan, Department of Parasitology and Mycology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, IR Iran. Tel: +98-935530151, E-mail: mgn1370@yahoo.com

Received 2016 April 08; Revised 2016 April 26; Accepted 2016 April 30.

Abstract

Background: Geohelminth infections are common infections worldwide. These helminthes are parasitic nematodes, which could infect humans and animals by contact with eggs or infective larvae and in some cases cause serious disorders, especially in children.

Objectives: The present study was designed to evaluate the prevalence of geohelminth eggs in the soil of schools and play areas of Kermanshah city, western Iran.

Materials and Methods: One hundred and ninety-two soil samples were collected from 24 primary schools and 24 play areas from four different parts of each area. Diagnosis of geohelminth was confirmed by the recovery of their eggs by the flotation technique using sucrose solution and examined under light microscope using 10X and 40X objectives.

Results: Out of 192 soil samples, soil of play areas from four regions was contaminated with geohelminth eggs. The rate of soil contamination with *Trichuris* spp. eggs was 3.12% while this was 2.1% for hookworm eggs, yet no *Ascaris* eggs were found in the examined soil samples.

Conclusions: Based on the results of this study there is a risk of infection with soil transmitted helminthes among children of Kermanshah, Iran.

Keywords: Geohelminth, Soil Contamination, School, Play Area, Nematodes

1. Background

Geohelminth infections are major intestinal pathogens, which affect underprivileged communities with incomplete admission to innocent water and appropriate health facilities. Soil Transmitted Helminth (STH) such as *Ascaris lumbricoides*, *Trichuris trichiura* and hookworms are among the most important geohelminths, which have been estimated to infect approximately two billion humans all over the world (1). Infection with geohelminths effect feeding, growth and fitness and also cause chronic infections in the population of endemic areas (1, 2). Soil provides appropriate conditions for conversion of the eggs to the infective stage. Also presence of eggs in the superficial layers of the soil can be considered as a factor affecting their transition to humans in each region (3). One of the main pathways for human infection

is ingestion of infective eggs by consumption of raw vegetables and fruits contaminated with soil containing eggs (2, 4). Thus, it seems that soil is the main resource for human infection and can be a direct indicator for the risk of infection among human populations, especially children. Evidences indicate that STH infections affect children in developing and developed countries and are directly associated with growth retardation, physical activity reduction and disorders in learning abilities (5, 6).

Basically, the endemicity of geohelminth depends on the presence of infected individuals, continuous contamination of the soil with feces and suitable soil condition and frequent contact between the soil and healthy individuals. Therefore, geohelminth infection rate is affected by both natural and cultural factors (3, 6, 7). Due to limited resources in most developing countries, prioritising the

control of such infections seems very difficult and it could be an important reason for the presence of the infection in these countries. These kinds of infections are considered as neglected diseases, which in case of suitable conditions can cause serious problems (4, 8).

2. Objectives

The present study aimed to determine the prevalence of soil contamination by geohelminth eggs in schools and play areas of Kermanshah city, West of Iran.

3. Materials and Methods

3.1. Study Area

Kermanshah city is located in the center of Kermanshah province in the West of Iran, and the city is divided to six regions. Its' population is estimated around 850000, where most of its population are engaged in agricultural activities. Kermanshah is one of the western agricultural centers of Iran that produces grain, rice, vegetable, fruits and oilseeds. Kermanshah has cold winters and usually has rainfalls in fall and spring.

3.2. Sample Collection

During the period from September to December 2014, a total of 192 soil samples were collected from 24 primary school and 24 play areas in Kermanshah city. In each of the six regions, the samples were collected from four play areas and four primary schools. The samples were collected from four different parts of each area, including northern, eastern, western and southern part. Samples were collected from loose soil, mainly from [U+FB02]owerbeds. Approximately 200 grams of soil was obtained from an approximate depth of 3 cm, and all samples were then placed in plastic containers, labeled with a number and were exposed to air to dry.

3.3. Parasitological Procedures

Two grams of each sample was isolated using 150 μ m mesh sieves and transferred into the test tubes. Next 10-mL Tween-80 with 0.05% concentration was added to the mixture and shook vigorously. The tubes were centrifuged at 1500 rpm for five minutes. In the next step, the supernatants were discarded and the tubes containing the sediments were filled with approximately 1 cm from the top with sucrose solution (1.2 g/cm³) and the sediments were suspended. The tubes were then centrifuged for 15 minutes at 1500 rpm. Finally, the tubes were filled with sucrose solution and a cover slip was placed on each tube and left stationary for 30 minutes and then the cover slips were placed on the microscopic slides and examined under a light microscope using 10X and 40X objectives.

4. Results

A total of 192 soil samples were examined and low rate of geohelminth egg contamination was observed. Although, the prevalence of free-living nematodes was considerably high in the studied areas (Table 1), *Trichuris* spp. eggs were not observed in any of the studied schools, but were found in three play areas (3.12%). Additionally, no *Ascaris* eggs were found in the examined soil samples.

5. Discussion

Public sites such as schools and play areas are considered an important source for human infection, especially for children. The places where children play are important for public health (9). In many studies, it has been demonstrated that parasitic infections have a considerable negative influence on health factors such as weight and height (10, 11).

Recent studies evaluated STH prevalence in the soil from parks, playgrounds, schools, etc. (12, 13). Reports of soil contamination with geohelminth eggs have been published in Iran and other countries, as summarized in Table 2. According to the results of the present study, soil contamination with geohelminth eggs (2.6%) is low in Kermanshah compared to the other studied areas in Iran and other countries. Environmental factors such as soil type, pre-processing sieving, suspension of sediment, washing and flotation and specific gravity of flotation solution seem to influence the recovery of geohelminth eggs in the soil (14, 15). Furthermore, some factors like climate, herbage and presence of animals could affect soil contamination (16). Increase in the number of stray dogs and cats plays an important role in soil contamination with hook worm eggs, one of the main causes of cutaneous larva migrans in humans, which were found in two play areas in separate regions of Kermanshah. The eggs are excreted by infected host to the soil and develop to third stage larva (L3), which is infective and can infect children by direct contact with contaminated soil in schools and play areas (17). During childhood, infection with hookworms plays an important role in moderate to severe anemia in preschool and school-aged children (18-20).

Trichuris spp. is another STH, which were found in another region in Kermanshah. Human infections with this parasite are associated with some clinical disorders, which are similar to that of inflammatory bowel disease, such as chronic abdominal pain, diarrhea and other disorders such as impaired growth, anemia and finger clubbing (21).

The results of the present cross-sectional study indicate that the current status of contamination of soil with

Table 1. Soil Contamination With Geohelminth Eggs in Different Parts of Kermanshah^a

Region	Collected Soil Samples		Free Living Nematodes		<i>Trichuris</i> Spp.		Hook Worms	
	School	Play Areas	School	Play Areas	School	Play Areas	School	Play Areas
1	16	16	2 (12.5)	13 (81.25)	0	0	0	1 (6.25)
2	16	16	0	14 (87.5)	0	1 (6.25)	0	0
3	16	16	2 (12.5)	9 (56.25)	0	0	0	1 (6.25)
4	16	16	3 (18.75)	14 (87.5)	0	0	0	0
5	16	16	3 (18.75)	8 (50)	0	0	0	0
6	16	16	7 (43.75)	6 (37.5)	0	2 (12.5)	0	0
Total	96	96	17 (17.7)	64 (66.6)	0	3 (3.12)	0	2 (2.1)

^aValues are expressed as No. (%).**Table 2.** Reports of Soil Contamination With Geohelminth Eggs Have Been Published in Iran and Other Countries

Country/city	Location of Study	Sample. No. (%)
Poland/Lodz	Park playground	168 (7.7)
	School sports fields	216 (15.7)
Poland/Lublin	Beach	215 (18.6)
Egypt/Sharkyia	Rural and urban areas	120 (28.3)
Philippines/Barangay Bayog	Rural areas	120 (54)
India/Titagarh	Soil of wastewater-irrigated areas	35 (68.6)
Nigeria/Abuja	Public parks	764 (67.5)
Iran/Tehran	Public places	150 (6.7)
Iran/shiraz	Public places	112 (10.7)
Iran/Kermanshah	School/play areas	192 (2.6)

soil transmitted helminth is fairly good, but some preventive measures such as controlling stray dogs and cats and preventing their entry into public places to keep schools and play areas and other public places uncontaminated seems necessary.

Acknowledgments

We thank the Parasitology and Mycology department of the school of medicine of Isfahan University of Medical Sciences, Isfahan, Iran.

Footnote

Authors' Contribution: Mohammad Ali Mohaghegh, Mohsen Ghomashlooyan and Mehdi Azami were respon-

sible for the experimental design of the study. Mohammad Ali Mohaghegh, Mohsen Ghomashlooyan, Mohammad Reza Vafaei, Nooshin Hashemi and Mohammad Falahati were responsible for the execution techniques and parasitological examination of soils. Farzaneh Mirzaei, Hamed Kalani, Somayeh Jahani and Seyed Hossein Hejazi were responsible for the statistical analysis and Mohammad Reza Vafaei for the collection of the samples. All authors reviewed and contributed to the writing of this manuscript.

References

- de Silva NR, Brooker S, Hotez PJ, Montresor A, Engels D, Savioli L. Soil-transmitted helminth infections: updating the global picture. *Trends Parasitol.* 2003;**19**(12):547-51. [PubMed: [14642761](#)].
- Savioli L, Albonico M. Soil-transmitted helminthiasis. *Nat Rev Microbiol.* 2004;**2**(8):618-9. doi: [10.1038/nrmicro962](#). [PubMed: [15303271](#)].
- Mizgajski H. The distribution and survival eggs of *Ascaris* suum in six different natural soil profiles. *Acta Parasitol.* 1993;**38**(4):170-4.
- Bethony J, Brooker S, Albonico M, Geiger SM, Loukas A, Diemert D, et al. Soil-transmitted helminth infections: ascariasis, trichuriasis, and hookworm. *Lancet.* 2006;**367**(9521):1521-32. doi: [10.1016/S0140-6736\(06\)68653-4](#). [PubMed: [16679166](#)].
- Brooker S. Estimating the global distribution and disease burden of intestinal nematode infections: adding up the numbers—a review. *Int J Parasitol.* 2010;**40**(10):1137-44. doi: [10.1016/j.ijpara.2010.04.004](#). [PubMed: [20430032](#)].
- Hotez PJ, Brindley PJ, Bethony JM, King CH, Pearce EJ, Jacobson J. Helminth infections: the great neglected tropical diseases. *J Clin Invest.* 2008;**118**(4):1311-21. doi: [10.1172/JCI34261](#). [PubMed: [18382743](#)].
- Petney TN. Environmental, cultural and social changes and their influence on parasite infections. *Int J Parasitol.* 2001;**31**(9):919-32. [PubMed: [11406141](#)].
- Umar A, Bassey S. Incidence of *Strongyloides stercoralis* infection in Ungogo, Nassarawa, Dala and Fagge local government areas of Kano state, Nigeria. *Bayero J Pure Appl Sci.* 2010;**3**(2):76-80.
- Gazzinelli A, Correa-Oliveira R, Yang GJ, Boatín BA, Kloos H. A research agenda for helminth diseases of humans: social ecology, environmental determinants, and health systems. *PLoS Negl Trop Dis.* 2012;**6**(4):eee1603. doi: [10.1371/journal.pntd.0001603](#). [PubMed: [22545168](#)].

10. Adams EJ, Stephenson LS, Latham MC, Kinoti SN. Physical activity and growth of Kenyan school children with hookworm, *Trichuris trichiura* and *Ascaris lumbricoides* infections are improved after treatment with albendazole. *J Nutr*. 1994;**124**(8):199–206. [PubMed: 8074755].
11. Ahmed A, Al-Mekhlafi HM, Al-Adhroey AH, Ithoi I, Abdulsalam AM, Surin J. The nutritional impacts of soil-transmitted helminths infections among Orang Asli schoolchildren in rural Malaysia. *Parasit Vectors*. 2012;**5**:119. doi: [10.1186/1756-3305-5-119](https://doi.org/10.1186/1756-3305-5-119). [PubMed: 22704549].
12. Zibaei A, Abdollahpour F, Birjandi M, Firoozeh F. Soil contamination with *Toxocara* spp. eggs in the public parks from three areas of Khorram Abad, Iran. *Nepal Med Coll J*. 2010;**12**(2):63–5. [PubMed: 2122397].
13. Khademvatan S, Abdizadeh R, Tavalla M. Molecular characterization of *Toxocara* spp. from soil of public areas in Ahvaz southwestern Iran. *Acta Trop*. 2014;**135**:50–4. doi: [10.1016/j.actatropica.2014.03.016](https://doi.org/10.1016/j.actatropica.2014.03.016). [PubMed: 24695242].
14. Santarém V, Dias A, Felix A, Rodenas R, da Silva A. Contaminação por ovos de *Toxocara* spp. em praças públicas das regiões central e periurbana de Mirante do Paranapanema, São Paulo, Brasil. *Vet Zoot*. 2012;**17**(1):47–53.
15. Santarem VA, Magoti LP, Sichieri TD. Influence of variables on centrifuge-flotation technique for recovery of *Toxocara canis* eggs from soil. *Rev Inst Med Trop Sao Paulo*. 2009;**51**(3):163–7. [PubMed: 19551291].
16. Santarém V, Rubinsky-Elefant G, Ferreira M. Soil-transmitted helminthic zoonoses in humans and associated risk factors. Rijeka: InTech; 2011. pp. 43–66.
17. Bowman DD, Montgomery SP, Zajac AM, Eberhard ML, Kazacos KR. Hookworms of dogs and cats as agents of cutaneous larva migrans. *Trends Parasitol*. 2010;**26**(4):162–7. doi: [10.1016/j.pt.2010.01.005](https://doi.org/10.1016/j.pt.2010.01.005). [PubMed: 20189454].
18. Brooker S, Peshu N, Warn PA, Mosobo M, Guyatt HL, Marsh K, et al. The epidemiology of hookworm infection and its contribution to anaemia among pre-school children on the Kenyan coast. *Trans R Soc Trop Med Hyg*. 1999;**93**(3):240–6. [PubMed: 10492749].
19. Stoltzfus RJ, Chwaya HM, Montresor A, Albonico M, Savioli L, Tielsch JM. Malaria, hookworms and recent fever are related to anemia and iron status indicators in 0- to 5-y old Zanzibari children and these relationships change with age. *J Nutr*. 2000;**130**(7):1724–33. [PubMed: 10867043].
20. Magalhaes RJ, Clements AC. Mapping the risk of anaemia in preschool-age children: the contribution of malnutrition, malaria, and helminth infections in West Africa. *PLoS Med*. 2011;**8**(6):e1000438. doi: [10.1371/journal.pmed.1000438](https://doi.org/10.1371/journal.pmed.1000438). [PubMed: 21687688].
21. Bundy DA, Cooper ES. Trichuris and trichuriasis in humans. *Adv Parasitol*. 1989;**28**:107–73. [PubMed: 2683614].