

Antibacterial Properties of Essential Oil of *Heracleum persicum* (Golpar) and Foodborne Pathogens



Nabi Shariatifar¹, Toktam Mostaghim², Afshin Afshar³, Issa Mohammadpourfard^{4*}, Mehran Sayadi⁵, Mohammad Rezaei⁴

¹Department of Environmental Health, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

²Department of Food Science and Technology, Shahr-e-Qods Branch, Islamic Azad University, Tehran, Iran

³Department of Food and Drug, Alborz University of Medical Sciences, Karaj, Iran

⁴Department of Food Hygiene and Quality Control, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

⁵Department of Food Hygiene and Quality Control, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

*Corresponding Author:

Issa Mohammadpourfard,
Tel: +989149080532;
Email:
Isamohammadpour@yahoo.com

Published Online May 20, 2017

Keywords: Antimicrobial activity, Essential oil, Food Quality, *Heracleum persicum*, Pathogenic bacteria



Abstract

Background: Antimicrobial resistance is a serious health threat to human society. Antibiotics and preservatives are also the most common modalities to increase the shelf life of foods.

Objectives: Antibacterial activity of essential oil of *Heracleum persicum* (Golpar) fruit against some of the main foodborne bacteria was determined.

Materials and Methods: Antibacterial activity of essential oil was evaluated against the bacteria (*Staphylococcus aureus*, *Escherichia coli*, *Salmonella typhi*, *Vibrio cholera*, and *Yersinia enterocolitica*) using disc diffusion method. Broth micro-dilution method was used to determine their minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC). The data was expressed as mean and standard deviation and analyzed using analysis of variance (ANOVA) in SPSS software ($P < 0.05$).

Results: Comparing with *S. typhi* strains (inhibition zone = 18 ± 0.46 mm), essential oil was found to be more effective against *V. cholera* strains (inhibition zone = 34 ± 0.2 mm). The MIC value (%) of the essential oil against *V. cholera* and *S. typhi* were 8 and 32 respectively. Essential oil of *H. persicum* (Golpar) showed a good antimicrobial activity against foodborne pathogens.

Conclusion: The results revealed that the essential oil of *H. persicum* can be used in food preservation systems to inhibit the growth of *V. cholera* strains and improve food quality and safety.

Received February 26, 2017; Revised May 7, 2017; Accepted May 17, 2017

Background

Foodborne pathogens belong to a group of micro-organisms that cause foodborne diseases and it is essential to find effective drugs against the infection.¹ Prevalence of foodborne diseases has increased worldwide and become a major cause of mortality in individuals with impaired immune systems in the developing countries.² There is an urgent need to monitor antimicrobial resistance by improving the ways the antibiotics may be used and reduce hospital cross-infection. New antibiotics need to be developed continuously to maintain the effectiveness of antimicrobial therapy.³⁻⁵ The recent decades have demonstrated an increase in interests in the use of herbal products.⁶ Plants play an important role in human health. It is estimated that 25% of modern medicines are derived directly or indirectly from herbs.^{7,8} Many plant compounds have been proven to be effective insecticidal, antifungal, anti-bacterial, anti-viral, anti-parasitic, and anti-oxidant agents.^{3,9} Aromatic essential oils are often

used in food industry to add flavor and prevent growth of microbial and mold contaminants.¹⁰ *Heracleum*, which is used for medicinal purposes, is an annual herbaceous plant. It is a genus of about 60 species, which mainly grows in Iran and Europe.^{11,12} *Heracleum persicum* is known as *Heracleum* in Farsi and as tromso palm in Norwegian. Fruits of this plant are used as spices and as anti-infection and analgesic in traditional medicine and the stems are used in pickles.¹³ Many studies have been conducted on the antibacterial effect of *Heracleum* and most of them have been on the leaves and stems; still, there has been no study on antibacterial effect of essential oil of its fruits. Literature review showed a paucity of studies on the antibacterial effects of chemical composition of the oil of *H. persicum* fruit on foodborne pathogens.

Objectives

This study aimed at determining the antibacterial effects of essential oil of *Heracleum* fruit on some of the

foodborne pathogens.

Materials and Methods

Plant Materials

Heracleum fruits were collected in May 2013 from the heights in Shemiranat-Tehran and then identified and confirmed in Herbarium Center of School of Pharmacy, Tehran University of Medical Sciences. The collected plants were dried in the shade and then the fruits were separated from the stems and ground by mechanical means.

Chemicals and Reagents

All chemicals and standards were obtained from Merck (Darmstadt, Germany).

Extraction of Essential Oil

The essential oils of the *Heracleum* were obtained using hydro distillation and a Clevenger apparatus; 300 g of fruits was placed in the Clevenger with sufficient distilled water to cover the material. Extraction continued for 3 consecutive hours when the water started to boil.¹⁴

Antimicrobial Activity Test

Disk Diffusion Test

The essential oils were tested against *Staphylococcus aureus* ATCC 25913, *Escherichia coli* ATCC 8739, *Salmonella enterica* PTCC 1709, *Vibrio cholera* PTCC 1611 and *Yersinia enterocolitica* PTCC 1477. The microorganisms were cultured in BHI (brain heart infusion) for 18 hours at 37°C, compared with 0.5 McFarland standard (5 × 10⁸ CFU/mL), and inoculated directly in boards with Müller-Hinton agar (Merck). After inoculation of each microorganism, the diffusion was carried out; so that 10 µL of the essential oil was placed on paper discs (6 mm of diameter) at 37°C for 24 hours. Afterward, the zone of inhibition was measured.³

Minimum Inhibitory Concentration and Minimum Bactericide Concentration

To obtain minimum inhibitory concentration (MIC) and minimum bactericide concentration (MBC), quantitative tests were performed using 1%-9% concentrations of the essential oil in Müller-Hinton broth medium. The concentrations were added by 1×10⁶ CFU/mL of the bacteria. Positive control (cultured along with the bacterium without the essential oil) and negative control (cultured without the bacterium) were also prepared. Then, the samples were placed in an incubator (37°C, 24 hours). Before reading the results, the last concentration without opaqueness (no growth) was adopted as MIC for each sample. The samples without turbidity were cultured in Müller-Hinton broth medium. After incubation (24 hours, 37°C), the last concentration suitable enough to be used as bactericide was adopted as MBC.

Statistical Analysis

Each experiment, from sample preparation to analysis,

was repeated for three times, and the data were analyzed using one-way analysis of variance (ANOVA) and Tukey tests in SPSS (v. 16).

Results

The results of disc diffusion method showed antimicrobial activity of essential oil of *H. persicum* against all of the bacterial strains; including *S. aureus* (inhibitory zone: 32.16±0.76 mm), *S. enterica* (inhibitory zone: 18±0.46 mm), *E. coli* (inhibitory zone: 19±1.32 mm), *V. cholera* (inhibitory zone: 34±2 mm), and *Y. enterocolitica* (inhibitory zone: 22.16±0.76 mm) (Table 1).

The results showed that yield of essential oil of *Heracleum* fruit was about 11%. Impact of the essential oil on each bacterium (*Staphylococcus*, *E. coli*, *Yersinia*, *Vibrio*, and *Salmonella*) is listed in Table1. The highest and lowest zones of growth inhibition at concentration of 100% were observed in *Vibrio* and in *Salmonella* respectively. In addition, minimum and maximum MIC against bacteria were in *Vibrio* and *Yersinia* respectively. As listed in Table 2, *Yersinia* and *Salmonella* (64%) and *Staphylococcus* (16%) have the highest and lowest values of MBC. There was no significant difference (*P*<0.05) between the bacterial strains in terms of inhibitory halo diameter except for *E. coli* and *Salmonella*.

Discussion

Many antimicrobial drugs have been found and there are still more drugs in the nature not found yet. In light of this, the past decade has witnessed an increase in research on plants as medicine source.^{15,16}

Essential oils are potential sources of novel antimicrobial compounds especially against bacterial pathogens. Essential oil plants may also be applied for the treatment of infectious diseases, thanks to its antimicrobial activity.^{15,16} Results from in vitro studies in this paper showed that the essential oils can inhibit bacterial growth; although, their effectiveness varied.

Nazemi et al indicated that aqueous extract of *Heracleum* did not have any antibacterial effect; however, *Heracleum* methanol extract had significant effect on the bacteria *Bacillus polymyxa*, *Bacillus subtilis*, *Enterococcus faecalis*, *Nocardia*, and *S. aureus*.^{17,18}

Kousha and Bayat indicated that extracts of leaves and

Table 1. Average Inhibitory Halo Diameter (mm) of Various Bacterial Strains for *Heracleum persicum* Essential Oil

Bacterial Strain	Samples	Range	Mean ± SD
<i>Staphylococcus aureus</i> ATCC 25913	3	32.5-33	32.16 ± 0.76 ^a
<i>Escherichia coli</i> ATCC 8739	3	17.5-20	19 ± 1.32 ^b
<i>Salmonella enterica</i> PTCC 1709	3	18-19.5	18 ± 0.46 ^b
<i>Vibrio cholera</i> PTCC 1611	3	32.5-34	34 ± 2 ^a
<i>Yersinia enterocolitica</i> PTCC 1477	3	22-22.5	22.16 ± 0.76 ^a

Table 2. MIC and MBC Value (%) for *Heracleum persicum* Essential Oil Against Pathogenic Bacterial Strains

	<i>Staphylococcus aureus</i>	<i>Vibrio cholera</i>	<i>Yersinia enterocolitica</i>	<i>Escherichia coli</i>	<i>Salmonella</i>
MIC	11	8	18	30	32
MBC	16	16	64	32	64

flowers of *H. persicum* had antimicrobial activity against 4 gram-negative and 4 gram-positive bacteria.¹⁹

Manikandan et al²⁰ reported that the methanolic extract of leaves of different plants were more active than the aqueous extract in presence of *Bacillus subtilis*, *Bacillus pumilus*, *Micrococcus luteus* and *S. aureus*.

Habibi et al²¹ showed that the highest inhibitory effect of *H. rechingeri* species was against gram-positive bacteria and *B. subtilis* in particular. Consistent results were obtained at concentration level of 50% ($P < 0.05$).

Hemati et al indicated that essential oil and hydro-alcoholic extract of *H. persicum* had analgesic, anti-inflammatory, and antioxidant effects.²²

Inhibition effects of *H. persicum* (57.16%) on *Agrobacterium tumefaciens* was reported by Dehghan Noudeh et al.²³ They reported that *H. persicum* (57.16%) had inhibiting effect on all tested gram-positive and gram-negative strains. Our results were consistent with the results of that study.

Frey and Meyers²⁴ indicated that the traditional medicines were capable of creating inhibition zone against the bacteria *S. typhimurium*, *S. aureus*, *E. coli* and *Streptococcus lactis*. Moreover, their data supported the hypothesis that the selection and use of these plants for disease treatment was not random.

Comparison of our results with those obtained by Dehghan Nodeh et al indicated that 4 fractions from *Heracleum* had antimicrobial activity against *A. tumefaciens*, *E. coli*, *S. aureus*, *B. subtilis* and *P. aeruginosa* and the highest and lowest MIC were observed in *E. coli* and *B. subtilis* respectively.²³

Several studies have been conducted on antibacterial activity of herbs, which are rich in monoterpenoids. Using drop diffusion method, these studies have shown significant inhibition zones for all microorganisms under study such as gram-positive bacteria (*S. aureus*) and fungus (*Candida albicans*).^{25,26} Results of the present study were consistent with other studies as to gram-positive (*S. aureus*) and gram-negative bacteria (*E. coli*, *S. enterica*, *V. cholera*, and *Y. enterocolitica*).

Inconsistencies between the present study and other works might be because of different genotypes of *H. persicum* utilized, growth ecosystem (habitat, temperature, height), and setting of the experiments (pH and temperature). Several factors influence growth and performance of the plants in an ecosystem including type, habitat, soil, height, and geographical position. Some of these factors have notable effects on the results quantitatively and qualitatively.

Medical herbs as immunostimulants have been widely used as food additives for years and the procedures have

been recorded by several research works. Moreover, in vivo, in vitro, and ex vivo surveys of the effects of *Heracleum* sp. used as an immunostimulant agent in terrestrial and aquatic animals or human require deeper investigations. In addition, more studies are needed on antiviral effects of this family.

Given the information obtained from these studies and the present study, anti-bacterial properties of *Heracleum* against the pathogens transmitted by food is supported.

Conclusion

Notable antimicrobial activity against foodborne pathogenic bacteria was demonstrated by the essential oil of *Heracleum*. The essential oil may be effective on other gram-positive and gram-negative bacteria. The practical application of essential oils is to increase shelf life of food. Therefore, it can be a good alternative for artificial preservatives used in the food industry today.

Authors' Contributions

This article was extracted from the MSc project of IM where NS supervised these project and suggested the problem; TM, AA and MS participated in determination of sample points, sample preparation procedure and extraction of essential oil; MR wrote and edited the manuscript. All authors read and approved the final manuscript.

Ethical Approval

This study was approved by Ethics Committee of Tehran University of Medical Sciences.

Conflict of Interest Disclosures

The authors declare that they have no conflict of interests.

Financial Support

This study was supported by Research Committee of Tehran University of Medical Sciences.

References

1. Scallan E, Hoekstra RM, Angulo FJ, et al. Foodborne illness acquired in the United States—major pathogens. Emerg Infect Dis. 2011;17(1):7-15. doi:10.3201/eid1701.P11101.
2. Lund BM, O'Brien SJ. The occurrence and prevention of foodborne disease in vulnerable people. Foodborne Pathog Dis. 2011;8(9):961-973. doi:10.1089/fpd.2011.0860.
3. Shariatifar N, Fathabad AE, Khaniki GJ, Gandomi H. Evaluation of the antibacterial activity of essential oil and aqueous and ethanolic extracts of *Quercus infectoria* leaves on food-borne pathogenic bacteria. Int J Pharm Sci Res. 2014;5(10):709-713.
4. Ultee A, Kets E, Smid E. Mechanisms of action of carvacrol on the food-borne pathogen *Bacillus cereus*. Appl Environ Microbiol. 1999;65(10):4606-4610.
5. Shetty K, Labbe RG. Food-borne pathogens, health and role of dietary phytochemicals. Asia Pac J Clin Nutr. 1998;7:270-276.
6. Cowan MM. Plant products as antimicrobial agents. Clin Microbiol Rev. 1999;12(4):564-582.
7. Fowler MW. Plants, medicines and man. J Sci Food Agric. 2006;86(12):1797-1804. doi:10.1002/jsfa.2598

8. Kamkar A, Tooryan F, Basti A, Misaghi A, Shariatifar N. Chemical composition of summer savory (*Satureja hortensis* L.) essential oil and comparison of antioxidant activity with aqueous and alcoholic extracts. J Vet Re. 2013;68(2):183-190.
9. Boukraâ L, Abdellah F, Ait-Abderrahim L. Antimicrobial Properties of Bee Products and Medicinal Plants. <http://www.formatex.info/microbiology4/vol2/960-970.pdf>. Published 2013.
10. Holley RA, Patel D. Improvement in shelf-life and safety of perishable foods by plant essential oils and smoke antimicrobials. Food Microbi. 2005;22(4):273-292.
11. Shahrani M. The effects of methanol extracts of garlic and *Heracleum persicum* on basal and pentagastrin-stimulated gastric acid and pepsin secretions in rat. Kerman University of Medical Sciences, Kerman, Iran; 2009.
12. Chauhan NS. Medicinal and Aromatic Plants of Himachal Pradesh. Indus Publishing; 1999.
13. Shahrani M, Nabavizadeh F, Shirzadeh H, Yousefi H, Taghi Moradi M, Moghaddasi J. IC Journal Master List. Journal of Shahrekord University of Medical Sciences. 2006;8(1):35-41. [Persian].
14. Rezaei M, Karimi F, Shariatifar N, Mohammadpourfard I, Malekabad ES. Antimicrobial Activity of the essential oil from the leaves and seeds of *Coriandrum sativum* toward food-borne pathogens. West Indian Medical Journal. 2016;65(1):8-12. doi: 10.7727/wimj.2014.162.
15. Lam KS. New aspects of natural products in drug discovery. Trends Microbiol. 2007;15(6):279-289.
16. Koehn FE, Carter GT. The evolving role of natural products in drug discovery. Nat Rev Drug Discov. 2005;4(3):206-220. doi:10.1038/nrd1657.
17. Nazemi A, Khatami M, Purshamsiyan K, Hashemi M. First survey of Antimicrobial activity aqueous and ethanolic extracts of the plant *Heracleum persicum*. Medical Science Journal of Islamic Azad University. 2005;15(2):91-94. [Persian].
18. Nazemi A, Hashemi M, Khataminejad M, Pourshamsian K. Study on antibacterial effect of methanol and aquatic extract of *Heracleum persicum* plant. Islamic Azad Univ Med Sci. 2005;15:91-94.
19. Kousha A, Bayat M. Bactericidal and fungicidal activity of methanolic extracts of *Heracleum persicum* Desf. ex fischer against some aquatic and terrestrial animal pathogens. Int Jf Pharm. 2012;8(7):652-656.
20. Manikandan T, Neelakandan T, Rani GU. Antibacterial activity of *Salicornia brachiata*, a halophyte. J Phytol. 2009;1(6):441-443.
21. Habibi Z, Eshaghi R, Mohammadi M, Yousefi M. Chemical composition and antibacterial activity of essential oil of *Heracleum rechingeri* Manden from Iran. Nat Prod Res. 2010;24(11):1013-1017. doi:10.1080/14786410902885138.
22. Hemati A, Azarnia M, Angaji A. Medicinal effects of *Heracleum persicum* (Golpar). Middle-East J Sci Res. 2010;5(3):174-176.
23. Dehghan Noudeh G, Sharififar F, Dehghan Noudeh A, et al. Antitumor and antibacterial activity of four fractions from *Heracleum persicum* Desf. and *Cinnamomum zeylanicum* Blume. Journal of Medicinal Plants Research. 2010;4(21):2176-2180.
24. Frey FM, Meyers R. Antibacterial activity of traditional medicinal plants used by Haudenosaunee peoples of New York State. BMC Complement Altern Med. 2010;10:64. doi:10.1186/1472-6882-10-64.
25. Oliveira DR, Leitao GG, Bizzo HR, Alviano DS, Alviano CS, Leitão SG. Chemical and antimicrobial analyses of essential oil of *Lippia origanoides* HBK. Food Chem. 2007;101(1):236-240.
26. Kuljanabagavad T, Sriubolmas N, Ruangrunsi N. Chemical composition and antimicrobial activity of the essential oil from *Heracleum siamicum*. J Health Res. 2010;24(2):55-60.